

# There Are No Maximal d.c.e. *wtt*-degrees

Guohua Wu<sup>1</sup>(✉) and Mars M. Yamelev<sup>2</sup>

<sup>1</sup> Division of Mathematical Sciences, School of Physical and Mathematical Sciences,  
Nanyang Technological University, Singapore 637371, Singapore  
guohua@ntu.edu.sg

<sup>2</sup> Institute of Mathematics and Mechanics, Kazan Federal University,  
18 Kremlyovskaya Street, Kazan 420008, Russia  
mars.yamaleev@ksu.ru

## 1 Introduction

In this article, we will study the weak-truth-table (*wtt*, for short) degrees of d.c.e. sets and show that there is no maximal d.c.e. *wtt*-degree.

**Theorem 1.** *For any d.c.e.  $wtt$ -degree  $\mathbf{d}_{wtt} < \mathbf{0}'_{wtt}$ , there is a d.c.e.  $wtt$ -degree  $\mathbf{c}_{wtt}$  strictly between  $\mathbf{d}_{wtt}$  and  $\mathbf{0}'_{wtt}$ .*

Here  $\mathbf{0}'_{wtt}$  is the *wtt*-degree of  $K$ , the halting problem. Theorem 1 says that for any d.c.e. set  $D$ , if  $D$  is *wtt*-incomplete, then we can split  $K$  into c.e. sets  $B$  and  $C$  such that  $K$  cannot be *wtt*-reducible to any of  $B \oplus D$  and  $C \oplus D$ . As  $K$  is *wtt*-equivalent to  $B \sqcup C$ , we have that  $B \oplus C \oplus D$  is *wtt*-equivalent to  $K$ . Thus,  $B \oplus D$  and  $C \oplus D$  are not *wtt*-reducible to each other, so they are strictly above  $\mathbf{d}_{wtt}$ . Our current work shows that d.c.e. *wtt*-degrees can always split above less ones (in progress), an analogue of Ladner and Sasso's result for c.e. *wtt*-degrees in [19].

Before giving a proof of the theorem above, we first review some well-known facts of density/nondensity of Turing degrees of c.e. sets and d.c.e. sets. Recall a set  $A \subseteq \mathbb{N}$  is computably enumerable (c.e. for short) if  $A$  is a domain of some partial computable function, and  $D \subseteq \mathbb{N}$  is d.c.e. if  $D$  is the difference of two computably enumerable sets, i.e.  $D = A - B$  for some c.e. sets  $A$  and  $B$ . The research on the structures of the c.e. degrees and the d.c.e. degrees has shown many nice properties and also many pathological properties, which are always accompanied with new techniques of constructions.

For the c.e. degrees, Sacks proved that this structure is dense and every nonzero element splits, and Lachlan proved that the density and splitting above

---

Wu is partially supported by AcRF Tier 2 grants MOE2011-T2-1-071 (ARC 17/11, M45110030) and MOE2016-T2-1-083 from Ministry of Education of Singapore, and by AcRF Tier 1 grants, RG29/14, M4011274 and RG32/16, M4011672 from Ministry of Education of Singapore.

Yamaleev is supported by Russian Foundation for Basic Research (projects 15-41-02507, 15-01-08252), by research grant of Kazan Federal University, and by the subsidy allocated to Kazan Federal University for the project part of the state assignment in the sphere of scientific activities (project 1.2045.2014).

cannot be combined, where Lachlan developed the  $0'''$  argument for the first time, an argument being called “monstrous” construction in the 1980s.

Cooper initiated the study of the structure of d.c.e. degrees in his PhD thesis [3] in 1971. Lachlan observed that the d.c.e. degrees are downwards dense and Cooper [5] proved that each nonzero d.c.e. degree splits, an analogue of Sacks splitting. Recall that a Turing degree is properly d.c.e. if it contains a d.c.e. set, but no c.e. sets. As c.e. degrees are also d.c.e., what Lachlan and Cooper needed to do in their proofs is to consider the case when the given degree is properly d.c.e. As pointed in Downey and Stob [12], and Cooper and Li [8], it is necessary to have the cases separated, as no uniform way working for both cases exists. Cooper [4] even proved that the  $\text{low}_2$  d.c.e. degrees are dense.

Cooper and Yi considered the interaction between c.e. degrees and d.c.e. degrees in [9] and introduced the notion of isolation. The existence of isolated degrees can be obtained from a result in Kaddah’s paper [17], where she proved that low c.e. degrees branch in the d.c.e. degrees. Using this interaction phenomenon, Wu [24] provided another proof of Downey’s diamond theorem, where Wu used the isolation to connect the cupping and the capping parts of the diamond embeddings. See Wu and Yamaleev’s survey [25] on this topic.

Even though these two degree structures share several algebraic properties, these two structures are not elementarily equivalent. This was first proved in the 1980s by Arslanov in [2] and Downey in [11]. As to the density, Cooper, Harrington, Lachlan, Lempp and Soare proved that the d.c.e. degrees are not dense, where they constructed a maximal d.c.e. degree  $\mathbf{d}$  below  $\mathbf{0}'$ . Obviously,  $\mathbf{0}'$  does not split above  $\mathbf{d}$ .

We will consider d.c.e. *wtt*-degrees in this paper, i.e. the weak-truth-table degrees of d.c.e. sets. For  $A, B \subseteq \mathbb{N}$ , say that  $A$  is *weak-truth-table reducible* to  $B$ , denoted as  $A \leq_{wtt} B$ , if there is a partial computable functional  $\Phi_e$  and a computable function  $f$  such that (i)  $A = \Phi_e^B$ , and (ii) for every  $x$ ,  $f(x) \geq u(B; e, x)$ , where  $u(B; e, x)$  is the use of the computation  $\Phi_e^B(x)$ . We use  $\varphi_e$  to denote the use function of  $\Phi_e$ . The *wtt*-reduction was proposed by Friedberg and Rogers in 1959 in [15], and is now also called *bounded Turing* reduction. Lachlan proved that the upper semi-lattice of c.e. *wtt*-degrees is distributive, providing a crucial structural difference between c.e. *wtt*-degrees and c.e. Turing degrees. Ladner and Sasso then gave another difference in [19] by showing that the splitting and density can be combined for the c.e. *wtt*-degrees. Technically, weak-truth-table degrees can be handled much easier than Turing degrees. For instance, density of the c.e. *wtt*-degrees can be proved by a finite injury argument, whereas the analogous result for c.e. Turing degrees requires an infinite injury priority proof.

On the other hand, some structural properties of Turing degrees can be obtained from those of *wtt*-degrees via the so-called contiguous degrees. Here, a c.e. Turing degree  $\mathbf{c}$  is *contiguous* if  $\mathbf{c}$  contains exactly one c.e. *wtt*-degree. That is, any two c.e. sets  $A, B$  in a contiguous degree  $\mathbf{c}$  are *wtt*-equivalent. Ladner and Sasso proved in [19] that any nonzero *wtt*-degree  $\mathbf{c}$  has the anticupping property in the c.e. *wtt*-degrees. Thus, when  $\mathbf{c}$  is contiguous,  $\mathbf{c}$  also has the anticupping property in the Turing degrees, a result first proved by Yates

by direct construction. This kind of “transfer” phenomenon has been further developed by Ambos-Spies in [1], Stob in [23], and Downey in [10].

In this paper, we consider the *wtt*-degrees of d.c.e. sets, and the remainder of this paper will be devoted to the proof of Theorem 1: there are no maximal d.c.e. *wtt*-degrees.

Our notation and terminology are standard and generally follow Soare [22] and Odifreddi [20]. The readers can refer Cooper’s paper [6] for d.c.e. Turing degrees and Ambos-Spies’ paper [1], Stob’s paper [23] and Downey’s paper [10] for the general idea on c.e. *wtt*-degrees.

## 2 Requirements and Construction

Let  $K = \{e : \varphi_e(e)\}$ , Turing’s halting problem, and let  $\{K_s : s \in \omega\}$  be a recursive enumeration of  $K$ . Note that  $K$  is *wtt*-complete among all d.c.e. sets, and we will assume that for each  $s$ ,  $|K_{s+1} \setminus K_s| = 1$ . Let  $D$  be any d.c.e. set in  $\mathbf{d}_{wtt}$ , and  $\{D_s : s \in \omega\}$  be a d.c.e. approximation of  $D$ . An additional addition for this approximation is: for any  $s$ ,  $|(D_{s+1} \setminus D_s) \cup (D_s \setminus D_{s+1})| = 1$ .

For the proof of Theorem 1, we will construct c.e. sets  $B$  and  $C$  satisfying the following requirements:

$\mathcal{S}$ :  $K = B \sqcup C$ ;

$\mathcal{P}_e^B$ :  $K \neq \Phi_e^{B \oplus D}$ , where the use function of  $\Phi_e^{B \oplus D}$ , i.e.  $\varphi_e$ , is bounded by  $\psi_e$ ;

$\mathcal{P}_e^C$ :  $K \neq \Phi_e^{C \oplus D}$ , where the use function of  $\Phi_e^{C \oplus D}$ , i.e.  $\varphi_e$ , is bounded by  $\psi_e$ .

Here  $\{(\Phi_e, \psi_e) : e \in \omega\}$  is a recursive list of all pairs  $(\Phi, \psi)$ ,  $\Phi$  a partial computable functional  $\Phi$  and  $\psi$  a partial computable function. As indicated at the beginning of this paper, the  $\mathcal{S}$ -requirement ensures that  $K$  and  $B \oplus C \oplus D$  are *wtt*-equivalent. All the  $\mathcal{P}_e^B$  requirements,  $e \in \omega$ , ensure that  $K$  is not *wtt*-reducible to  $B \oplus D$ , and all the  $\mathcal{P}_e^C$  requirements,  $e \in \omega$ , ensure that  $K$  is not *wtt*-reducible to  $C \oplus D$ . Thus,  $B \oplus D$  and  $C \oplus D$  are not *wtt*-reducible to each other, which implies that both are strictly above  $\mathbf{d}_{wtt}$ .

The idea of satisfying the  $\mathcal{S}$ -requirement is standard. That is, at any stage  $s$ , we find a requirement  $\mathcal{R}$  with the highest priority with  $k_s$  less than the restraint  $r(\mathcal{R}, s)$ , if exists. If  $\mathcal{R}$  is a  $\mathcal{R}_e^B$ -requirement, then enumerate  $k_s$  into  $C$ . Otherwise, enumerate  $k_s$  into  $B$ . Obviously,  $B \sqcup C = K$ .

Before we describe how to satisfy a  $\mathcal{P}$ -requirement, a  $\mathcal{P}_e^B$ -requirement, say, we first review the idea when  $D$  is c.e., and then we show the changes we need to make for the case when  $D$  is d.c.e.

The main idea for the case when  $D$  is c.e. is the Sacks preservation strategy, i.e., to find a disagreement between  $K$  and  $\Phi_e^{B \oplus D}$ , we define expansionary stages and extend a *wtt*-reduction  $\Delta_e$  at expansionary stages such that if there were infinitely many expansionary stages, then we would have  $\Delta_e^D = K$ , which is impossible. Here we define the length of the agreement between  $K$  and  $\Phi_e^{B \oplus D}$  at stage  $s$  as

$$\ell^B(e, s) = \max\{x : (\forall y < x)[\Phi_e^{B \oplus D}(y)[s] \downarrow \text{ with use } \varphi_e(y) < \psi_e(y) \text{ and } \Phi_e^{B \oplus D}(y)[s] = K_s(y)]\},$$

and say that  $s$  is an expansionary stage if for any expansionary stage  $t < s$ ,  $\ell^B(e, s) > \ell^B(e, t)$ . At an expansionary stage  $s$ , for any  $y < \ell^B(e, s)$ , if  $\Delta_e^D(y)$  has no definition at stage  $s$ , then define  $\Delta_e^D(y)[s] = K_s(y)$  with use  $\delta_e(y)[s] = \varphi_e(y)[s]$ . So,  $\delta_e$  is bounded by  $\psi_e$ , and if there were infinitely many expansionary stages, then  $\Phi_e^{B \oplus D}$  would be total and  $\Delta_e^D$  would be defined as a total function, showing that  $K \leq_{wtt} D$ , which is impossible. Thus, there are only finitely many expansionary stages, and we will have some  $y < \ell^B(e)$ , the length of agreement at the last expansionary stage, such that either  $\Phi_e^{B \oplus D}(y) \uparrow$  or  $\Phi_e^{B \oplus D}(y) \neq K(y)$ .

In this strategy, the main point is to protect computations at expansionary stages. Assume that after an expansionary stage  $s_1$  (so a restraint is imposed on the  $B$ -part to protect computations), we see that a computation  $\Phi_e^{B \oplus D}(y)$  changes, with  $y < \ell^B(e, s_1)$ , because of the changes of  $D$  between stages  $s_1$  and  $s_2$  say. We also assume that  $s_2$  is not an expansionary stage, but we see that  $\Phi_e^{B \oplus D}(y)[s_2]$  converges. The strategy says that at the next expansionary stage  $s_3$ , we will protect  $\Phi_e^{B \oplus D}(y)[s_3]$ . This computation  $\Phi_e^{B \oplus D}(y)[s_3]$  we see at stage  $s_3$  could be also different from  $\Phi_e^{B \oplus D}(y)[s_2]$ , and  $\Phi_e^{B \oplus D}(y)[s_2]$  is not protected. It is okay, for  $D$  c.e., as either there are no more expansionary stages, or the changes between stages  $s_2$  and  $s_3$  will remain forever in  $D$ , and no more computation of  $\Phi_e^{B \oplus D}(y)$  can be the same as  $\Phi_e^{B \oplus D}(y)[s_2]$ . Thus, among all the computations of  $\Phi_e^{B \oplus D}(y)$  with use  $\varphi_e(y)[s] < \psi_e(y)$ , we only protect those we see at expansionary stages, and for  $y$  above, the change of  $D$  undefines  $\Delta_e^D(y)$ , and we will redefine  $\Delta_e^D(y)$  again at the next expansionary stage. Nothing is complicated in this case.

As we are assuming that the uses of  $\Phi_e^{B \oplus D}(y)$  are bounded by  $\psi_e(y)$ , the nature of finite injury allows to protect all computations of  $\Phi_e^{B \oplus D}(y)$  we see at all stages. That is, whenever we see a new computation of  $\Phi_e^{B \oplus D}(y)$ , at stage  $s_2$  above, we can protect it and redefine  $\Delta_e^D(y)[s_2] = K_{s_2}(y)$ . Of course, if the computation  $\Phi_e^{B \oplus D}(y)$  changes between stage  $s_2$  and  $s_3$ , then the  $D$ -changes undefine  $\Delta_e^D(y)[s_2]$  again, allowing us to redefine it at stage  $s_3$ .

We adopt this idea of protecting all computations for our purpose when  $D$  is d.c.e. It can happen that a computation  $\Phi_e^{B \oplus D}(y)[s]$  changes because of some enumeration of  $z$  into  $D$ , and after many stages, when  $z$  leaves  $D$ , at stage  $t$  say, the  $D$ -part of the oracle  $B \oplus D$  recovers to the status at stage  $s$ , and if we protect  $\Phi_e^{B \oplus D}(y)[s]$  at stage  $s$ , then we will have  $\Phi_e^{B \oplus D}(y)[t] = \Phi_e^{B \oplus D}(y)[s]$ . This variation of Sacks preservation strategy allows us to deal with cases when  $D$  is any  $\Delta_2^0$  set, i.e.  $\mathbf{0}'_{wtt}$  splits above any other  $\Delta_2^0$   $wtt$ -degrees.

We are now ready to provide a full construction. We first list the requirements as follows:

$$\mathcal{S} < \mathcal{P}_0^B < \mathcal{P}_0^C < \mathcal{P}_1^B < \mathcal{P}_1^C < \dots < \mathcal{P}_e^B < \mathcal{P}_e^C < \dots$$

We say that a requirement  $\mathcal{Q}$  has priority higher than  $\mathcal{R}$  if  $\mathcal{Q} < \mathcal{R}$  in the order defined above. So  $\mathcal{S}$  has the highest priority, and at any stage  $s$ , we will enumerate  $k_s$  into one of  $B$  and  $C$ , but not both.

For a  $\mathcal{P}$ -requirement,  $\mathcal{P}_e^B$  say, we call stage  $s$  a  $\mathcal{P}_e^B$ -*identical stage* if

1.  $\ell^B(e, s) = \ell^B(e, s - 1)$ ,
2. for all  $y \leq \ell^B(e, s)$ ,  $\Phi_e^{B \oplus D}(y)[s]$  converges if and only if  $\Phi_e^{B \oplus D}(y)[s - 1]$  converges,
3. for  $\Phi_e^{B \oplus D}(y)[s]$  converges, the computation  $\Phi_e^{B \oplus D}(y)[s]$  and  $\Phi_e^{B \oplus D}(y)[s - 1]$  are the same.

We say that  $\mathcal{P}_e^B$  *requires attention at stage*  $s$  if  $s$  is not a  $\mathcal{P}_e^B$ -identical stage.

Construction at stage 0: For all the requirements, let the corresponding restraint as 0.

Construction at stage  $s > 0$ :

Step 1. Among requirements  $\mathcal{P}_0^B, \mathcal{P}_0^C, \mathcal{P}_1^B, \mathcal{P}_1^C, \dots, \mathcal{P}_s^B, \mathcal{P}_s^C$ , check which one requires attention at stage  $s$ . Let it be  $\mathcal{Q}[s]$ , and set the corresponding restraint as  $s$ . For those  $y$  less than the length of agreement of  $\mathcal{Q}[s]$ , define  $\Delta_e^D(y) = K_s(y)$  with use  $\delta_e(y)[s] = \varphi_e(y)[s]$ . Initialize all the requirements with priority lower than  $\mathcal{Q}[s]$ .

Step 2. Among all the requirements with priority not lower than  $\mathcal{Q}[s]$ , find the one with higher priority,  $\mathcal{R}[s]$  say, whose restraint is larger than  $k_s$ . If  $\mathcal{R}[s]$  is a  $\mathcal{P}^B$ -strategy, then enumerate  $k_s$  into  $C$ . Otherwise, enumerate  $k_s$  into  $B$ . Initialize all the requirements with priority lower than  $\mathcal{R}[s]$ .

This completes the construction of stage  $s$ .

*End of construction*

### 3 Verification

In this section, we verify that the constructed c.e. sets  $B$  and  $C$  satisfy all the requirements. The actions at step 2 of each stage  $s$  ensure that  $K = B \sqcup C$ , and hence

**Lemma 1.** *The requirement  $\mathcal{S}$  is satisfied.*

Now we verify that all the  $\mathcal{P}$ -requirements are satisfied. The following lemma is enough to show this.

**Lemma 2.** *For each  $e \in \omega$ ,*

1.  $\mathcal{P}_e^B$  can be initialized at most finitely many times;
2.  $\mathcal{P}_e^B$  requires attention at most finitely many times;
3.  $\mathcal{P}_e^B$  has finite restraint;
4. The same are true for requirement  $\mathcal{P}_e^C$ .

*Proof.* We prove it by induction on  $e$ . So we can assume that after a stage  $s_0$  large enough, no more  $\mathcal{P}_{e'}$ -requirements, with  $e' < e$ , requires attention, or requires further enumeration of elements of  $K$  into  $B$ . Thus, after stage  $s_0$ ,  $\mathcal{P}_e^B$  cannot be initialized anymore. (1) holds.

To prove (2), we assume that  $\mathcal{P}_e^B$  requires attention infinitely often. Then, the bounding function  $\psi_e$  is total. As  $D$  is d.c.e., we assume that after stage  $s_1$ ,  $D$  becomes fixed up to  $\psi_e(0)$ . According to steps 1 and 2 in every stage, after stage  $s_0$ , we protect all the computations of  $\Phi_e^{B\oplus D}(0)$  whenever we have a new computation, thus, for any computation of  $\Phi_e^{B\oplus D}(0)$ , if it converges before stage  $s_1$ , at stage  $s'$  say, and the  $D$ -part of the use agrees with  $D \upharpoonright \psi_e(0)$ , then this computation will converge forever. Of course, no such a computation occurs before stage  $s_1$ , then, when  $\mathcal{P}_e^B$  requires attention again, we will have a new computation of  $\Phi_e^{B\oplus D}(0)$ , which will be protected. In both cases,  $\Phi_e^{B\oplus D}(0)$  converges. The same idea can be used to prove that  $\Phi_e^{B\oplus D}$  converges at any  $n$ , by induction.

We now show that  $\Delta_e^D$  is total and computes  $K$  correctly. Again, we first show that  $\Delta_e^D(0)$  is defined, with  $\Delta_e^D(0) = K(0)$ , and the same argument can be applied to show that for any  $n$ ,  $\Delta_e^{D(n)}$  is defined and equals to  $K(n)$ .

We assume again that  $D$  has no more change below  $\psi_e(0)$  after stage  $s_1$ . Then for  $\Delta_e^D(0)$  defined at stage  $s^*$  with the  $D$ -part of the use agreeing with  $D \upharpoonright \varphi_e(0)[s^*]$ , i.e.

$$D \upharpoonright \varphi_e(0)[s^*] = D_{s_1} \upharpoonright \varphi_e(0)[s^*],$$

we have  $\Delta_e^D(0) = \Delta_e^D(0)[s^*]$ . To see this, assume that  $s^* < t_1 < t_2 < \dots < t_n < s_1$  be a list of stages with  $D_{t_i} \upharpoonright \varphi_e(0)[s^*] = D_{s^*} \upharpoonright \varphi_e(0)[s^*]$  for each  $i \in \{1, \dots, n\}$ , then  $\Delta_e^D(0)[t_i] = \Delta_e^D(0)[s^*]$  with use  $\varphi_e(0)[s^*]$ . This actually shows that for any definition of  $\Delta_e^D(0)$ , which is defined at other stages,  $D$  must have changes at some number  $z$  below  $\varphi_e(0)[s^*]$ , before stage  $s^*$  (if  $\Delta_e^D(0)$  is defined before stage  $s^*$ ), or between any two stages in this list (if  $\Delta_e^D(0)$  is defined between these two stages). Of course, if there is no such a stage  $s^*$ , then at stage  $s_1$ , we define  $\Delta_e^D(0)$  and by the choice of  $s_1$ ,  $D$  will have no change any more and hence  $\Delta_e^{D(0)}$  is defined.

Now we show that  $\Delta_e^D(0) = K(0)$ . Note that at stage  $s^*$  above, we have  $\Phi_e^{B\oplus D}(0)[s^*]$  converges, and this computation is protected since  $s^*$  onwards and hence after stage  $s_1$ , i.e. the computation  $\Phi_e^{B\oplus D}(0)$  will be the same as  $\Phi_e^{B\oplus D}(0)[s^*]$ . Thus,  $\Phi_e^{B\oplus D}(0) = \Phi_e^{B\oplus D}(0)[s^*]$ , and as we assume that there are infinitely many stages  $\mathcal{P}_e^B$  requires attention, we know that after stage  $s_1$ , the agreement of  $\mathcal{P}_e^B$  will be always larger than 0, and hence  $K(0) = \Phi_e^{B\oplus D}(0)[s^*]$  forever, and as a consequence,  $\Delta_e^D(0) = K(0)$ .

We can then apply the same idea and show that for any  $n$ ,  $\Delta_e^D(n)$  is defined and equals to  $K(n)$ . This shows that  $K \leq_{utt} D$  via  $\Delta$ . A contradiction. Thus (2) is true for  $\mathcal{P}_e^B$  requirement.

Note that (2) tells us the existence of a stage  $s_2$ , after which the  $\mathcal{P}_e^B$  requirement never requires attention again, which means that after stage  $s_2$ , all stages are  $\mathcal{P}_e^B$ -identical, and hence no more restraint is imposed. This shows that the last restraint imposed by  $\mathcal{P}_e^B$  is before stage  $s_2$ , and as a consequence, the  $\mathcal{P}_e^B$  requirement has finite restraint. (3) is true.

The same argument can show that (1), (2), (3) above are also true for  $\mathcal{P}_e^C$  requirement. (4) is true.

This completes the proof of Lemma 3.2, and hence the proof of Theorem 1.

## 4 Further Remarks

As pointed out in the introduction, we can improve Theorem 1 and show that the d.c.e. *wtt*-degrees are dense, and hence for a given Turing degree  $\mathbf{d}$ , it can either contain exactly one d.c.e. *wtt*-degree, or contain infinitely many d.c.e. *wtt*-degree. We call a d.c.e. Turing degree  $\mathbf{d}$  contiguous if it contains exactly one d.c.e. *wtt*-degree. A recent work of the authors shows the existence of properly d.c.e. contiguous degrees. We have seen that c.e. contiguous degrees have many unusual applications, like Downey's idea of using c.e. contiguous degrees to show the downwards density of c.e. degrees with strong anti-cupping property, and we are interested in problems of properly d.c.e. contiguous degrees, like the distribution of such degrees and how we can use such degrees to transfer properties of *wtt*-degrees to Turing degrees.

## References

1. Ambos-Spies, K.: Contiguous R.E. degrees. In: Börger, E., Oberschelp, W., Richter, M.M., Schinzel, B., Thomas, W. (eds.) *Computation and Proof Theory*. LNM, vol. 1104, pp. 1–37. Springer, Heidelberg (1984). doi:[10.1007/BFb0099477](https://doi.org/10.1007/BFb0099477)
2. Arslanov, M.: Structural properties of the degrees below  $0'$ . *Dokl. Nauk. SSSR* **283**, 270–273 (1985)
3. Cooper, S.B.: Degrees of unsolvability. Ph.D. thesis, Leicester University (1971)
4. Cooper, S.B.: The density of the low<sub>2</sub> *n*-r.e. degrees. *Arch. Math. Logic* **31**, 19–24 (1991)
5. Cooper, S.B.: A splitting theorem for the *n*-r.e. degrees. *Proc. Am. Math. Soc.* **115**, 461–471 (1992)
6. Cooper, S.B.: Local degree theory. In: Griffor, E.R. (ed.) *Handbook of Computability Theory*, pp. 121–153. North-Holland, Amsterdam (1999)
7. Cooper, S.B., Harrington, L., Lachlan, A.H., Lempp, S., Soare, R.I.: The d.r.e. degrees are not dense. *Ann. Pure Appl. Logic* **55**, 125–151 (1991)
8. Cooper, S.B., Li, A.: Non-uniformity and generalised Sacks splitting. *Acta Math. Sin.* **18**, 327–334 (2002)
9. Cooper, S.B., Yi, X.: Isolated d.r.e. degrees. University of Leeds, Department of Pure Mathematics, Preprint series, vol. 17 (1995)
10. Downey, R.:  $\Delta_2^0$  degrees and transfer theorems. III. *J. Math.* **31**, 419–427 (1987)
11. Downey, R.: D.r.e. degrees and the nondiamond theorem. *Bull. London Math. Soc.* **21**, 43–50 (1989)
12. Downey, R., Stob, M.: Splitting theorems in recursion theory. *Ann. Pure Appl. Logic* **65**, 1–106 (1993)
13. Ershov, Y.L.: A hierarchy of sets, Part I (Russian). *Algebra i Logika* **7**, 47–73 (1968). *Algebra and Logic* (English translation), **7**, 24–43 (1968)
14. Ershov, Y.L.: A hierarchy of sets, Part II (Russian). *Algebra i Logika* **7**, 15–47 (1968). *Algebra and Logic* (English Translation), **7**, 212–232 (1968)
15. Friedberg, R.M., Rogers, H.: Reducibility and completeness for sets of integers. *Z. Math. Logik Grundlag. Math.* **5**, 117–125 (1959)
16. Ishmukhametov, S.: D.r.e. sets, their degrees and index sets. Ph.D. thesis, Novosibirsk, Russia (1986)
17. Kaddah, D.: Infima in the d.r.e. degrees. *Ann. Pure Appl. Logic* **62**, 207–263 (1993)

18. Lachlan, A.H.: Lower bounds for pairs of recursively enumerable degrees. Proc. London Math. Soc. **16**, 537–569 (1966)
19. Ladner, R.E., Sasso, L.P.: The weak-truth-table degrees of recursively enumerable sets. Ann. Math. Logic **8**, 429–448 (1975)
20. Odifreddi, P.: Classical Recursion Theory. Studies in Logic and the Foundations of Mathematics, vol. 143. Elsevier, New York (1999)
21. Sacks, G.E.: The recursively enumerable degrees are dense. Ann. Math. **80**, 300–312 (1964)
22. Soare, R.I.: Recursively Enumerable Sets and Degrees. Springer, Heidelberg (1987)
23. Stob, M.: *Wtt*-degrees and *T*-degrees of r.e. sets. J. Symbolic Logic **48**, 921–930 (1983)
24. Wu, G.: Isolation and lattice embeddings. J. Symbolic Logic **67**, 1055–1064 (2002)
25. Wu, G., Yamaleev, M.M.: Isolation: motivations and applications. Proc. Kazan Univ. (Phys. Math. Ser.) **52**, 204–217 (2012)