

EPR study of spectra transformations of the intrinsic vanadyl-porphyrin complexes in heavy crude oils with temperature to probe the asphaltenes' aggregation



M.R. Gafurov^{a,*}, M.A. Volodin^{a,b}, A.A. Rodionov^a, A.T. Sorokina^a, M.Yu. Dolomatov^{c,d}, A.V. Petrov^e, A.V. Vakhin^a, G.V. Mamin^a, S.B. Orlinskii^a

^a Kazan Federal University, Kremlevskaya, 18, Kazan, 420008, Russia

^b Sakhalin Energy Investment Company, Ltd., Dzerzhinskogo, 35, Yuzhno-Sakhalinsk, 693020, Russia

^c Bashkir State University, Validy, 32, Ufa, 450076, Russia

^d Ufa State Petroleum Technical University, Kosmonavtov, 1, Ufa, 450000, Russia

^e Institute of Petroleum Refining and Petrochemistry, Iniciativnaya, 12, Ufa, 450065, Russia

ARTICLE INFO

Keywords:

Bitumen
Crude oil
Heavy oil
Rotational correlation time
Temperature dependence
ESR

ABSTRACT

Temperature dependencies of electron paramagnetic resonance spectra of intrinsic paramagnetic vanadyl complexes and dynamical viscosity for two heavy crude oils and asphalt samples are measured. Transitions between the different motional conditions (from the rigid to the fast motion regimes) are observed. The rotational correlation times (in the model of the isotropic diffusion) are extracted. It is shown that the characteristic temperatures for the motional regime transitions are mainly defined by the asphaltenes' content. From our analysis it follows that the thermal treatment leads to the destruction of the asphaltene complexes onto the 4–5 small pieces. The results indicate that paramagnetic vanadyl complexes are the sensitive intrinsic probes to study qualitatively and quantitatively structural transformations of asphaltenes of heavy crude oils in-situ.

1. Introduction

Asphaltene constituents of petroleum (defined as the heaviest fractions of crude oil insoluble in normal alkanes) impact all aspects of crude oil production and utilization (Trukhan et al., 2017; Speight, 2015; Ganeeva et al., 2011). The content of the high-molecular asphaltene components could reach the values of 45 wt % in native oils and up to 73 wt % in natural asphalts and bitumen. Undesirable asphaltene precipitation is a serious concern to the petroleum industry because asphaltenes can block catalyst active sites, plug up well bores and stop oil production, in addition to blocking pipelines (Ganeeva et al., 2011; Gawel et al., 2005; Yen and Chilingarian, 2000). As it is stressed anew in the recent review by Martyanov et al. (2017), a fundamental condition for the development of effective processes for heavy oil production and processing is the understanding of the key factors that determine the aggregative behaviour and stability of oil disperse systems (ODS), behaviour and chemical transformations of their components under various external conditions including elevated temperature.

Asphaltenes are known to form nanosized colloidal particles

“nanocrystallites”, “nanocolloids”, or “nanoaggregates”. It is shown that asphaltene aggregation begins already at their concentrations as high as 10 mg/L in some “good” organic solvents, as described by Evdokimov et al. (2016), Mullins et al. (2014) and by Shkalikov et al. (2010). The smallest aggregates contain 5–8 small asphaltene molecules. For example, in the well-known Yen-Mullins model (Mullins et al., 2014) the asphaltenes are defined as a central moderately-sized polycyclic aromatic hydrocarbon ring system linked with alkane chains on the exterior, with nanoaggregates up to six molecules and clusters of eight nanoaggregates on average (Lima et al., 2017).

However, after decades of intensive studies, the association and aggregation of asphaltenes are still not well characterized and are subjects of ongoing debate (Lima et al., 2017; Trukhan et al., 2017; Evdokimov et al., 2016; Ganeeva et al., 2011). Additionally, structural and molecular variability of asphaltenes do not allow to use an exact copy of one model which satisfactory describes a properties of one sort of ODS to other ODS. Therefore, the complex analytical study of each kind of ODS is in need both for the fundamental research and practical applications.

A significant factor influencing the asphaltene aggregation and

* Corresponding author.

E-mail address: Marat.Gafurov@kpfu.ru (M.R. Gafurov).

precipitation is their local environment, which determines the nature of intermolecular interactions of asphaltenes with other ODS components. It is evident that prediction of crude oil stability and solubility of asphaltenes under specified conditions requires the fundamental knowledge of the character of intermolecular interactions between asphaltenes and other heavy oil components present in the local environment. This knowledge can be gained only by means of in situ studies of local rheological properties of crude oils and molecular dynamics of asphaltenes under various conditions (Martyanov et al., 2017; Trukhan et al., 2017).

As it was already mentioned, because studying asphaltene properties in the actual (native) ODS and its fractions is difficult (especially for the optical studies), the vast majority of experimental and theoretical work has been conducted on systems wherein asphaltene obtained from the ODS was then dissolved in simple solvents. Electron paramagnetic resonance (EPR) techniques do not require the diluted samples and can sense the intrinsic paramagnetic centres (PC) also in the unfractionated species, i.e., in the native environment (see Yen and Chilingarian, 2000; Khasanova et al., 2017; Gafurov et al., 2018 for reviews on this topic). Following the changes of the EPR spectroscopic parameters of PC (for example with temperature) one can gain additional information about ODS.

Metalloporphyrins in heavy oils were first discovered by Treibs (1936) who postulated the biological origin of these compounds from chlorophyll, bacteriochlorophylls, hemes, and other tetrapyrrolic biochemical (Gourier et al., 2010). Vanadyl containing porphyrins, including paramagnetic species with VO^{2+} as a core, are the important part of many ODS and the object of EPR investigations (Reynolds, 1985; Yakubov et al., 2017; Biktagirov et al., 2017; Gracheva et al., 2016; Ramachandran et al., 2015; Gourier et al., 2010; Gilinskaya, 2008).

Since recently a role of VO^{2+} in the asphaltene and asphaltene-resin aggregation has begun to be extensively discussed (Dechaine and Gray, 2010). From the relaxation times measurements for “free” organic radical (FR) and VO^{2+} by Mamin et al. (2016), the existence of the dipole-dipole coupling manifested in the acceleration of the transverse relaxation times of FR was found for different asphaltenes. Consequently, it was proposed that VO^{2+} complexes can participate in construction of the asphaltene aggregates via the intermolecular interactions. The first evidence that the magneto-optical activity in asphaltenes can also be due to the vanadyls in asphaltenes was presented recently by Edelman et al. (2016). Trukhan et al. (2017) found that the precise simulation of the vanadyls’ EPR spectra is an effective tool for the quantitative determination of the asphaltene sizes in different oil fractions in situ and allows one to find their size distribution in crude oils. Cui et al. (2017a, 2017b) show that small changes of components of g-factor of vanadyls in asphaltenes, maltenes and resins can be used to track the mobility of VO^{2+} complexes under the different treatment.

In this work we inspect the changes of the conventional EPR spectrum of the intrinsic paramagnetic vanadyl complexes during the thermal treatment of heavy crude oil and asphalt samples with the aim to gain information about the ODS structure and dynamics as well as to show the abilities of the conventional EPR approaches for the in-situ analysis.

2. Materials and methods

2.1. Materials

There is no unique ODS classification (Briggs et al., 1988; Yen and Chilingarian, 2000; Souza et al., 2012). According to Briggs et al. (1988), heavy oils are defined as having the room temperature viscosity η_0 from 100 to 10 000 mPa s and density higher than 930 kg/m³. Heavy crude oils from Ashalchinskoe and Mordovo-Karmalskoe oilfields and of asphalt obtained by means of second stage propane deasphalting of the Western Siberian oil tar and were studied. They were investigated by SARA (Saturated and Aromatic hydrocarbons, Resins and Asphaltenes) analysis and stored sealed in dark place at room temperature. No special

procedures for the oxygen removing were conducted. Some physical properties of the samples are listed in Table 1. Analysis of the organic solutions of the samples # 1 and #2 with nuclear magnetic resonance (NMR) were presented by Rakhmatullin et al. (2017). Additional information for sample # 3 and its fractions one can find in the paper of Dolomatov et al. (2016).

2.2. Methods

The measurements of the dynamic viscosity η were performed using a rotary viscometer Brookfield DV2 with a thermostatic cell in the temperature range of (20–160) °C. The deviation of temperature during the measurements did not exceed ± 0.5 °C and with Stabinger viscometer SVM 3000/G2 with a thermostatic cell in the temperature range of (20–100) °C with the relative error of 0.5%.

EPR spectra of the samples were obtained using an ESP-300 spectrometer (Bruker) operating at 9.4–9.9 GHz (X-band) in continuous wave mode. Spectrometer’s magnetic field can be varied in the range of 20–1600 mT with an accuracy of not less than 0.01 mT, modulation frequency was fixed to 100 kHz. PC concentration was estimated at room temperature in the double cavity ER4105DR by comparing the integrated intensities of the spectra of a test sample and a reference sample (powder of DPPH and Mn^{2+} in MgO). Samples heating was done in a stream of nitrogen blown through the quartz insert in the high-temperature resonator ER 4114HT equipped with the VT thermostatic system (Oxford Instruments). For each sample the temperature dependence of the EPR spectra in the range of about (293–780) K was obtained with (8–15) °C increment, heating rate of 2 °C/min and temperature setting accuracy of not less than ± 2 °C. The chosen heating rate is slow enough to cause technical artifacts and allows to carry out the temperature experiments on a sample during a working day, Biktagirov et al. (2014) and Galukhin et al. (2016) have it recently shown. Modulation amplitude of 0.5 mT was chosen to investigate the temperature changes of vanadyl complexes. Microwave power level was equal to 25 μW in order to avoid saturation effects.

3. Results and discussion

Firstly, we present the EPR results. Fig. 1 shows the EPR spectra for sample #2 at two temperature values ($T = 297$ K and 654 K). The existence of paramagnetic centers in ODS is generally caused by the presence of *d* metals (mainly V, Ni, Fe) and stable carbon centered “free” radicals (FR). It is established that the majority of the organic FR are concentrated in the structure of a condensed polyaromatic core of asphaltene molecule while the metals locate in the polar fractions (both resin and asphaltene), with a further majority concentrated in the asphaltenes (Yen and Chilingarian, 2000). The presented spectra contain two components. The first contribution is due to the VO^{2+} complex. Because of the near planar structure of the vanadyl porphyrin complexes, the corresponding *g* and *A* tensors have an axial symmetry (Table 2). Therefore, the “powder” EPR spectrum of VO^{2+} at room and the lower temperatures consists of 16 partially overlapped components of hyperfine structure (8 lines in parallel and 8 in perpendicular orientations due to $I = 7/2$). The second (single line) can be attributed to FR. For details of the EPR spectra interpretation see Dolomatov et al., 2016; Alexandrov et al., 2014; Biktagirov et al., 2014; Yen and Chilingarian, 2000. In this work we are concentrating on the tracing of EPR of VO^{2+} complex. Temperature dependence of VO^{2+} EPR for asphalt (sample #3) is shown in Fig. 2.

From Figs. 1 and 2 it follows that the vanadyl spectra transform with temperature from the typical anisotropic “powder” spectra to the isotropic one. This transition could be characterized by two temperature values which we denote as T_{start} (EPR spectrum at $T < T_{\text{start}}$ is described by the only anisotropic “rigid” pattern) and T_{end} (at $T > T_{\text{end}}$ the EPR spectrum has only the isotropic “averaged” fast motion component). The values of T_{start} and T_{end} for the investigated samples are listed in Table 3.

As it follows from SARA analysis (Table 1), for sample #3 only the

Table 1
Physical properties of the investigated samples at near room temperature.

Sample #	Origin	Density at 20°C, kg/m ³	Viscosity at 20°C, Pa·s	Viscosity at 60°C, Pa·s	Asphaltenes, wt %	Resins, wt %	Saturates and aromatics, wt %	Concentration of VO ²⁺ complexes from EPR, 10 ¹⁸ spin/g
Sample # 1	Mordovo-Karmalskoe	948.0	0.536	0.050	3.1	25.4	71.8	8.6 ± 0.9
Sample # 2	Ashalchinskoe	968.2	2.426	0.1267	3.9	31.0	65.2	11 ± 1.2
Sample # 3	Asphalt	1044	–	158.6	17.8	30.4	65.1	5.2 ± 0.8




Fig. 1. EPR spectra for the heavy oil sample #2 measured at $T = 297$ K (curve 1) and $T = 654$ K (curve 2).

Table 2
Experimental components of g values and ⁵¹V hyperfine coupling parameters A (in MHz) of vanadyl complexes in all the investigated samples.

g_x	g_y	g_z	A_x	A_y	A_z
1.984(1)	1.984(1)	1.964(1)	157(15)	157(15)	471(20)




Fig. 2. EPR spectra for asphalt sample #3 with temperature.

concentration of asphaltenes is significantly higher than that for other species while other components are practically the same as for sample #2. From that we can speculate that the significant variations for the

Table 3
Characteristic temperatures of the vanadyls mobility transitions from isotropic to the anisotropic EPR pattern.

	T_{start} (K)	T_{end} (K)
Sample # 1	470(10)	510(10)
Sample # 2	480(10)	540(10)
Sample # 3	660(15)	740(20)

temperature values of the obtained transition (Table 3) are defined mainly by asphaltenes rather than by other constituents.

Secondly, we extract (estimate) the characteristic rotational mobility parameters. Different theoretical approaches for various rotational regimes are applied for simulation of VO²⁺ EPR spectrum (see Biktagirov et al. (2014), Trukhan et al. (2014) for details and literature references). (1) When the motional averaging is not significant a powder spectrum is observed; its lineshape is determined by inhomogeneous broadening and is not influenced by the value of τ_{rot} (so called rigid limit). (2) At some values of τ_{rot} the rotational diffusion becomes more prominent and the modulation of g and A tensors anisotropy by the rotational motion appears. Thereby, the lineshape changes significantly with τ_{rot} and its simulation requires a rigorous theoretical approach (usually, based on the density matrix theory). (3) The faster rotational diffusion leads to the effective averaging of g factor anisotropy, the linewidths can be calculated according to Wilson-Kivelson theory, Wilson and Kivelson, 1966. (4) The limiting case of rotational averaging corresponds to the effect of motional narrowing; the spectrum can be described by isotropic hyperfine constants and does not depend on the correlation time (an isotropic limit).

To analyze how the X-band EPR spectrum of VO²⁺ complex is related to its mobility, we have simulated the series of the corresponding X-band EPR spectra with respect to the value of the rotational correlation time. Spectra simulations were performed in the assumption of isotropic rotational diffusion of VO²⁺ by using EasySpin software module for Matlab (Stoll and Schweiger, 2006). The principle components of g tensor and hyperfine splitting tensors were fixed at the values obtained for room-temperature EPR spectra. The results of the simulation are presented in Fig. 3. It follows that in the X-band the EPR spectra are sensitive to the changes of τ_{rot} faster than 10⁻⁷ s, i.e., it is quite difficult to give exact values of τ_{rot} near room temperatures (at $T \ll T_{start}$, cf. Fig. 2). Therefore, the estimations of rotational mobility only in the vicinity of anisotropic-isotropic transformation are reliable.

The experimental results support the conclusion of Biktagirov et al. (2014) that the obtained temperature changes corresponds to the change of rotational mobility time from about 10⁻⁸ s up to 10⁻¹⁰ s, i.e., of two orders of magnitude. Fig. 4 graphically presents the temperature transformations of τ_{rot} for sample #3.

Fig. 5 shows the possible simulation of EPR spectra in the “mixed” regime ($T_{start} < T < T_{end}$) with two “slow” and “fast” rotational correlation times of 10 ns and 0.1 ns for sample #1. Though this simulation (as well as for other samples) looks reasonable, one cannot insist that it is the only one solution.

Interestingly, the extracted from EPR correlation times at high temperatures ($T > T_{end}$) are close to those reported by Mullins with his




Fig. 3. Simulation of EPR spectra of VO^{2+} in X-band for different values of rotational time expressed in the decimal logarithms (in seconds, with the step of 0.25 for the decimal logarithm) with the components of g and A tensors from Table 2.




Fig. 4. Extracted values of τ_{rot} as a function of temperature that accounts T_{start} and T_{end} for sample #3 (filled squares). In the mixed regime two values for τ_{rot} (filled squares and open circles) are given.

coworkers for the diluted in organic solvents asphaltenes of (0.2–1.2) ns derived from the time-resolved fluorescence depolarization (TRFD) experiments (see Mullins et al. (2014) and references therein). It is again, in our opinion, proves our suggestion that the observed transformations of vanadyls' EPR reflect mainly the mobility of asphaltenes' molecules.

Thirdly, we connect the rheological and EPR parameters. Fig. 6 presents the temperature dependence of viscosity along with the corresponding approximations (extrapolations). Viscosity is one of the key rheological parameter in petrophysics. The modelling of viscosity of asphaltene solutions is a longstanding unresolved problem. A number of empirical and semi-empirical viscosity models have been proposed. However, no consensus about the influence of asphaltenes structure and asphaltenes concentration on the viscosity is achieved and no single equation is found to adequately describe all the asphaltene systems (Daučík et al., 2008; Pal and Can, 2015; Pal, 2016). Very often Arrhenius type description is used (Daučík et al., 2008; Luo and Gu, 2006). Our results could not be described by single exponential decay. For simplicity, we have described the viscosity data with two exponents as for binary mixtures (Fig. 6).




Fig. 5. Simulation of EPR spectrum for sample #1 at $T = 517$ K as a sum of two components with $\tau_1 = 0.1$ ns (line 1) and $\tau_2 = 10$ ns (line 2). An insert demonstrates a part of the experimental spectrum in the vicinity of the magnetic fields corresponding to the nuclear spin projections of $m_l = 1/2$ and $m_l = 3/2$ as one of the most sensitive to the temperature transformation part of EPR spectrum.

Now let us to estimate the characteristic scales of the asphaltenes' aggregates. It is known that τ_{rot} at a given temperature can be determined by the local viscosity η and the characteristic radius R of the rotating particles using the Stokes–Einstein–Debye equation (Trukhan et al., 2014)

$$\tau_{\text{rot}} = 4\pi R^3 \eta(T) / 3k_B T, \quad (1)$$

where k_B is Boltzmann constant.

From Eq. (1) the characteristic radius can be expressed as

$$R = (3\tau_{\text{rot}} k_B T / 4\pi \eta(T))^{1/3}. \quad (2)$$

From Table 3, Figs. 3–5 and Eq. (2) it immediately follows that independently of the sample the characteristic radius derived from EPR in the “fast” motion regime is roughly $(100)^{1/3} \approx (4\text{--}5)$ times less than in the “slow” motion one. It is in correspondence with the Mullins calculations of aggregation of 7–8 small asphaltene molecules (see Mullins et al. (2014) and references therein).

The extraction of absolute values of R is a quite a tricky task. Obviously, as it was pointed out above, at near room temperatures (at $T \ll T_{\text{start}}$) one cannot use Eq. (2) for the valid estimations. At high temperatures one cannot reliably measure the viscosity by using the standard (approved) methods and tools. Table 4 presents the results of estimations of R in the vicinity of the isotropic-anisotropic transformation exploiting the extrapolations of $\eta(T)$ from Fig. 6.

Size distribution for the benzene diluted asphaltenes from about 0.3 nm up to 2.0 nm are reported from the small-angle X-ray scattering (SAXS) and EPR analyses (Trukhan et al., 2014). Sizes of (0.12–0.24) nm for the smallest asphaltenes molecules are given in papers of Mullins with his coworkers (see Martyanov et al. (2017), Ganeeva et al. (2011) and Mullins et al. (2014) and references therein). Our data at $T \geq T_{\text{end}}$ for the heavy oil samples #1 and #2 are in a good correspondence with those.

Obviously, the extracted values for sample #3 are far away from the reality. It means that the applied approach is apparently not valid for the high-viscous carbonaceous systems. Besides the absence of the viscosity data for the temperature range of interest, one can suppose that the simple slow and fast motion models cannot adequately describe changes of vanadyls EPR spectra in the complex systems.




Fig. 6. Temperature dependencies of dynamic viscosities (semilog plot) and their approximations (with the extrapolations) by two-exponential decays.

Table 4

Characteristic radius of asphaltenes as extracted from Eq. (2) at temperatures of the vanadyls mobility transitions from isotropic to the anisotropic regimes.

	$T = T_{\text{start}}$	$T = T_{\text{end}}$
Sample # 1	1.6(2) nm	0.36(6) nm
Sample # 2	1.0(1) nm	0.23(5) nm
Sample # 3	0.20 (5) nm	0.047(8) nm

4. Conclusions

In this work we show that the transformations of VO^{2+} EPR spectra with temperature in native (untreated) hydrocarbon systems are mainly due to the asphaltene molecules. We hope that our findings would help to improve the understanding of the asphaltene association and aggregation in native environments as well as of intermolecular interactions of asphaltenes with other ODS components. The results indicate that EPR of VO^{2+} could be used as a tool for measuring viscosity or its analogue at high temperatures. Taking into account that EPR is a non-destructive method and there is no need for sample preparation, it demonstrates that EPR can be used for the on-line monitoring or even for the EPR logging to follow the influence of thermal or catalytic treatment of heavy crude oil.

Acknowledgments

The work was funded by the subsidy of the Russian Government to support the Program of competitive Growth of Kazan Federal University.

References

- Alexandrov, A.S., Archipov, R.V., Ivanov, A.A., Gnezdilov, O.I., Gafurov, M.R., Skirda, V.D., 2014. The low-field pulsed mode dynamic nuclear polarization in the pentavalent chromium complex and crude oils. *Appl. Magn. Reson.* 45, 1275–1287.
- Biktagirov, T.B., Gafurov, M.R., Volodin, M.A., Mamin, G.V., Rodionov, A.A., Izotov, V.V., Vakhin, A.V., Isakov, D.R., Orlinskii, S.B., 2014. EPR study of rotational mobility of vanadyl-porphyrin complexes in crude oil asphaltene: probing the effect of thermal treatment of heavy oils. *Energy Fuels* 28, 6683–6687.
- Biktagirov, T., Gafurov, M., Mamin, G., Gracheva, I., Galukhin, A., Orlinskii, S., 2017. In situ identification of various structural features of vanadyl porphyrins in crude oil by high-field (3.4 T) electron–nuclear double resonance spectroscopy combined with density functional theory calculations. *Energy Fuels* 31, 1243–1249.
- Briggs, P.J., Baron, P.R., Fulleylove, R.J., 1988. Development of heavy oil reservoirs. *J. Petrol. Technol.* 40, 206–214.

- Cui, Q., Nakabayashi, K., Ma, X., Miyawaki, J., Ideta, K., Tennichi, Y., Ueda, M., Al-Mutairi, A., Marafi, A.M.J., Park, J., Yoon, S., Mochida, I., 2017a. Studying rotational mobility of V=O complexes in atmospheric residues and their resins and asphaltenes by electron spin resonance. *Energy Fuels* 31, 4748–4757. <https://doi.org/10.1021/acs.energyfuels.6b03279>.
- Cui, Q., Nakabayashi, K., Ma, X., Ideta, K., Miyawaki, J., Marafi, A.M.J., Al-Mutairi, A., Park, J., Yoon, S., Mochida, I., 2017b. Examining the molecular entanglement between V=O complexes and their matrices in atmospheric residues by ESR. *RSC Adv.* 7, 37908–37914. <https://doi.org/10.1039/C7RA06436E>.
- Daučík, P., Višňovský, J., Ambro, J., Hájeková, E., 2008. Temperature dependence of the viscosity of hydrocarbon mixtures. *Acta Chim. Slovaca* 1, 43–57.
- Dechaine, G.P., Gray, M.R., 2010. Chemistry and association of vanadium compounds in heavy oil and bitumen, and implications for their selective removal. *Energy Fuels* 24, 2795–2808.
- Dolomatov, M.U., Rodionov, A.A., Gafurov, M.R., Petrov, A.V., Biktagirov, T.B., Bakhtizin, R.Z., Makarchikov, S.O., Khairudinov, I.Z., Orlinskii, S.B., 2016. Concentration of paramagnetic centres at low-temperature thermal destruction of asphaltene of heavy petroleum distillates. *Magn. Reson. Solids* 18 (1–7), 16101. http://mrsej.kpfu.ru/contents/2016/MRSej_16101.pdf.
- Edelman, I.S., Sokolov, A.E., Zabluda, V.N., Shubin, A.A., Martyanov, O.N., 2016. Magnetic circular dichroism analysis of crude oil. *J. Structural Chemistry* 57, 382–389.
- Evdokimov, I.N., Fesan, A.A., Losev, A.P., 2016. New answers to the optical interrogation of asphaltene monomers and primary aggregates from steady state fluorescence studies. *Energy Fuels* 30, 4494–4503.
- Gafurov, M., Volodin, M., Biktagirov, T., Mamin, G., Orlinskii, S.B., 2018. High-field, pulsed, and double resonance studies of crude oils and their derivatives. In: Shukla, A.K. (Ed.), *Analytical Characterization Methods for Crude Oil and Related Products*. John Wiley & Sons Ltd, Hoboken, NJ, pp. 101–124.
- Galukhin, A.V., Khelkhal, M.A., Gerasimov, A.V., Biktagirov, T., et al., 2016. Mn-catalyzed oxidation of heavy oil in porous media: kinetics and some aspects of mechanism. *Energy Fuels* 30, 7731–7737.
- Ganeeva, Yu.M., Yusupova, T.N., Romanov, G.V., 2011. Asphaltene nano-aggregates: structure, phase transitions and effect on petroleum systems. *Russ. Chem. Rev.* 80, 993–1008.
- Gawel, I., Bociarska, D., Biskupski, P., 2005. Effect of asphaltene on hydroprocessing of heavy oils and residua. *Appl. Catal. A* 295, 89–94.
- Gilinskaya, L.G., 2008. EPR spectra of V(IV) complexes and the structure of oil porphyrins. *J. Struct. Chem.* 49, 245–254.
- Gourier, D., Delpoux, O., Bonduelle, A., Binet, L., Ciofini, I., Vezin, H., 2010. EPR, ENDOR, and HYSCORE study of the structure and the stability of vanadyl–porphyrin complexes encapsulated in silica: potential paramagnetic biomarkers for the origin of life. *J. Phys. Chem. B* 114, 3714–3725.
- Gracheva, I.N., Gafurov, M.R., Mamin, G.V., Biktagirov, T.B., Rodionov, A.A., Galukhin, A.V., Orlinskii, S.B., 2016. ENDOR study of nitrogen hyperfine and quadrupole tensors in vanadyl porphyrins of heavy crude oil. *Magn. Reson. Solids* 18 (1), 16102 (1–5). http://mrsej.kpfu.ru/contents/2016/MRSej_16102.pdf.
- Khasanova, N.M., Gabdrakhmanov, D.T., Kayukova, G.P., Mikhaylova, A.N., Morozov, V.P., 2017. EPR study of hydrocarbon generation potential of organic rich domain rocks. *Magn. Reson. Solids* 19 (1–5), 17102. http://mrsej.kpfu.ru/contents/2017/MRSej_17102.pdf.
- Lima, F.C.D.A., Alvim, R.D.S., Miranda, C.R., 2017. From single asphaltene and resins to nanoaggregates: a computational study. *Energy Fuels* 31, 11743–11754.
- Luo, P., Gu, Y., 2006. Effects of asphaltene content on the heavy oil viscosity at different temperatures. *Fuel* 86, 1069–1078.
- Mamin, G.V., Gafurov, M.R., Yusupov, R.V., Gracheva, I.N., Ganeeva, Yu.M., Yusupova, T.N., Orlinskii, S.B., 2016. Toward the asphaltene structure by electron paramagnetic resonance relaxation studies at high fields (3.4 T). *Energy Fuels* 30, 6942–6946.
- Martyanov, O.N., Larichev, Y.V., Morozov, E.V., Trukhan, S.N., Kazarian, S.G., 2017. The stability and evolution of oil systems studied via advanced methods in situ. *Russ. Chem. Rev.* 86, 999–1023.
- Mullins, O.C., Pomerantz, A.E., Zuo, J.Y., Dong, C., 2014. Downhole fluid analysis and asphaltene science for petroleum reservoir evaluation. *Annu. Rev. Chem. Biomol. Eng.* 5, 325–345.
- Pal, R., 2016. Modeling the viscosity of concentrated nanoemulsions and nanosuspensions. *Fluids* 1, 11–20. <https://doi.org/10.3390/fluids1020011>.
- Pal, R., Can, J., 2015. A new model for the viscosity of asphaltene solutions. *J. Chem. Eng.* 93, 747–755.
- Rakhmatullin, I.Z., Efimov, S.V., Margulis, B.Y., Klochkov, V.V., 2017. Qualitative and quantitative analysis of oil samples extracted from some Bashkortostan and Tatarstan oilfields based on NMR spectroscopy data. *J. Pet. Sci. Eng.* 156, 12–18.
- Ramachandran, V., van Tol, J., McKenna, A.M., Rodgers, R.P., Marshall, A.G., Dalal, N.S., 2015. High field electron paramagnetic resonance characterization of electronic and structural environments for paramagnetic metal ions and organic free radicals in deepwater horizon oil spill tar balls. *Anal. Chem.* 87, 2306–2313.
- Reynolds, J.G., 1985. Characterization of heavy residua by application of a modified D 2007 separation and electron paramagnetic resonance. *Liq. Fuel. Technol.* 3, 73–105.
- Shkalikov, N.V., Vasil'ev, S.G., Skirda, V.D., 2010. Peculiarities of asphaltene precipitation in n-alkane-oil systems. *Colloid J.* 72, 133–140.
- Souza de, T.A., Scheer, A.D.P., Khalil, M.C., Yamamoto, C.I., Luz Jr., L.F.D.L., 2012. Emulsion inversion using solid particles. *J. Pet. Sci. Eng.* 96–97, 49–57.
- Speight, J.G., 2015. *Handbook of Petroleum Product Analysis*, second ed. Wiley & Sons, Hoboken. 368 p.
- Stoll, S., Schweiger, A., 2006. EasySpin, a comprehensive software package for spectral simulation and analysis in EPR. *J. Magn. Reson.* 178, 42–55.

- Treibs, A., 1936. Chlorophyll-und Häminderivate in organischen Mineralstoffen. *Angew. Chem.* 49, 682–686.
- Trukhan, S.N., Yudanov, V.F., Gabrienko, A.A., Subramani, V., Kazarian, S.G., Martyanov, O.N., 2014. In situ electron spin resonance study of molecular dynamics of Asphaltenes at elevated temperature and pressure. *J. Energy Fuels* 28, 6315–6321.
- Trukhan, S.N., Kazarian, S.G., Martyanov, O.N., 2017. Electron spin resonance of slowly rotating vanadyls—effective tool to quantify the sizes of asphaltenes in situ. *Energy Fuels* 31, 387–394.
- Wilson, R., Kivelson, D., 1966. ESR linewidths in solution. I. Experiments on anisotropic and spin—rotational effects. *J. Chem. Phys.* 44, 154. <https://doi.org/10.1063/1.1726439>.
- Yakubov, M., Milordov, D., Yakubova, S., Abilova, G., Sinyashin, K., Tazeeva, E., Borisova, U., Mironov, N., Morozov, V., 2017. Vanadium and paramagnetic vanadyl complexes content in asphaltenes of heavy oils of various productive sediments. *Petrol. Sci. Technol.* 35, 1468–1472.
- Yen, T.F., Chilingarian, G.V., 2000. *Asphaltenes and Asphalts*, 2. Developments in Petroleum Science, 40 B. Elsevier, USA, 622 p.