


Sobolev Institute of Geology and Mineralogy SB RAS (IGM SB RAS)  
Trofimuk Institute of Petroleum Geology and Geophysics SB RAS (IPGG SB RAS)  
Novosibirsk State University (NSU)



# **X INTERNATIONAL SIBERIAN EARLY CAREER GEOSCIENTISTS CONFERENCE**

13-17 June 2022, Novosibirsk

**PROCEEDINGS OF THE CONFERENCE**



**MINERALOGY**  
2022



**Sobolev Institute of Geology and Mineralogy SB RAS (IGM SB RAS)**  
**Trofimuk Institute of Petroleum Geology and Geophysics SB RAS (IPGG SB RAS)**  
**Novosibirsk State University (NSU)**



**X INTERNATIONAL SIBERIAN EARLY CAREER  
GEOSCIENTISTS CONFERENCE**

**PROCEEDINGS OF THE CONFERENCE**



**MINERALOGY**  
2022

13-17 June 2022

Novosibirsk, Russia

УДК 55(061)  
ББК Д 26  
Т35

ORGANIZING COMMITTEE

Co-chairmen:

Dr. Nikolay N. Kruk,  
Corresponding Member of RAS, Director of IGM SB RAS  
Dr. Vyacheslav N. Glinskikh,  
Corresponding Member of RAS, Director of IPPG SB RAS  
Dr. Valery A. Vernikovsky,  
Academician of RAS, Dean of the DGG NSU

Conference Secretariat:

Dr. Maria O. Shapovalova (IGM SB RAS)  
Dr. Ekaterina A. Ovdina (IGM SB RAS)  
Elena S. Shaparenko (IGM SB RAS)

Technical Editors:

Dr. Ekaterina A. Ovdina (IGM SB RAS)  
Dr. Maria O. Shapovalova (IGM SB RAS)

X International Siberian Early Career GeoScientists Conference: Proceedings of the Conference (13-17 June 2022, Novosibirsk, Russia)/ IGM SB RAS, IPPG SB RAS, NSU. – Novosibirsk: IPC NSU, 2022. - 228 p. - ISBN 978-5-4437-1337-3.

The volume contains short papers of the X International Siberian Early Career GeoScientists Conference and offer information on multidisciplinary aspects of the Earth Science: petrology, geochemistry, geochronology, mineralogy, metallogeny, geodynamics and tectonics, sedimentology, geomorphology and neotectonics, paleontology and paleoclimatology, petroleum geosciences, economic geology, geoecology, hydrogeology, geophysics, mining engineering and geoinformation systems in the Earth Science.

The Conference was supported by IGM SB RAS, IPPG SB RAS, NSU, JSC "Siberian Anthracite", JSC "Almazy Anabara", Data East, LLC, PJSC "Gazprom Neft", Laboratory of ore-forming systems IGM SB RAS and SGA Siberian Student Chapter.

УДК 55(061)  
ББК Д 26

**ISBN 978-5-4437-1337-3**

© Sobolev Institute of Geology and Mineralogy  
SB RAS, 2022  
© Trofimuk Institute of Petroleum Geology and  
Geophysics SB RAS, 2022  
© Novosibirsk State University, 2022

# CONTENTS

## Section MINERALOGY

<b>THE COMPOSITION OF LITHOSPHERIC MANTLE BENEATH THE LOW-DIAMONDIFEROUS TSNIGRI-ARKHANGELSKAYA KIMBERLITE PIPE (ARKHANGELSK DIAMONDIFEROUS PROVINCE, NW RUSSIA)</b> <i>Agasheva E.V.</i> .....	11
<b>MODULARITY OF THE CRYSTAL STRUCTURES OF MINERALS AND SYNTHETIC COMPOUNDS WITH THE GENERAL FORMULA <math>A_2M_3(TO_4)_4</math></b> <i>Aksenov S.M., Merlino S.</i> 13	13
<b>METAMORPHISM VS METASOMATISM: STABLE ISOTOPE CONSTRAINTS (CASE STUDY OF THE KOCHUMDEK CONTACT AUREOLE, EAST SIBERIA)</b> <i>Deviatiarova A.S., Sokol E.V., Kokh S.N., Reutsky V.N., Izokh O.P., Pyryaev A.N.</i> .....	15
<b>THE DESCRIPTION OF ARAGONITES TWINNING FROM THE DEPOSITS OF SPAIN, MOROCCO AND RUSSIA</b> <i>Donskikh K.G., Gavryushkin P.N., Banaev M.V.</i> .....	17
<b>Ca-Mg-Fe OLIVINES IN PARALAVAS OF THE NYALGA COMBUSTION METAMORPHIC COMPLEX, CENTRAL MONGOLIA</b> <i>Glushkova V.E., Peretyazhko I.S., Savina E.A., Khromova E.A.</i> .....	19
<b>SPECTROSCOPIC FEATURES OF ELECTRON IRRADIATED DIAMOND CRYSTALS FROM MIR KIMBERLITE PIPE, YAKUTIA</b> <i>Komarovskikh A.Yu., Rakhmanova M.I.</i> .....	21
<b>MINERAL COMPOSITION OF SABAKTY LAKE SEDIMENTS</b> <i>Kuzina D.M., Yusupova A.R., Nourgalieva N.G., Rogov A.M.</i> .....	23
<b>THE SEQUENCE AND MECHANISMS OF PLATINUM GROUP MINERALS CRYSTALLIZATION FROM EUTECTIC GALENA-CHALCOPYRITE ORES OF THE OKTYABR'SKOE DEPOSIT, NORIL'SK REGION, RUSSIA</b> <i>Kuzmin I.A., Kalugin V.M.</i> 25	25
<b>PECULIARITIES OF THE MINERAL COMPOSITION OF THE POBEDA-2 HYDROTHERMAL FIELD (MAR)</b> <i>Lytkevich A.D.</i> .....	27
<b>VIVIANITE IN BOTTOM SEDIMENTS OF LAKE ONEGO</b> <i>Malov V.I., Strakhovenko V.D., Ovdina E.A.</i> .....	29
<b>ALLANITE IN MANTLE ECLOGITE</b> <i>Mikhailenko D.S., Stepanov A.S., Aulbach S., Gubanov N.V., Korsakov A.V., Xu Yi-G.</i> .....	31
<b>FORMATION CONDITIONS OF FE-MN CARBONATES FROM KERCH IRONSTONES: EVIDENCE FROM STABLE ISOTOPES</b> <i>Nekipelova A.V., Sokol E.V., Kokh S.N.</i> .....	33
<b>PHLOGOPITE FROM AILLIKITES OF THE ZIMINSKY COMPLEX OF THE EAST SAYAN REGION, RUSSIA</b> <i>Nugumanova Ya.N.</i> .....	35
<b>METHANE-DERIVED AUTHIGENIC CARBONATES OF THE LAPTEV SEA CONTINENTAL SLOPE</b> <i>Ruban A.S.</i> .....	37
<b>CRYSTAL CHEMISTRY OF CALCIUM DICARBONATE <math>CaC_2O_5</math>: STRUCTURAL TREND FROM DOUBLE <math>[CO_3]</math> TRIANGLES THROUGH ISOLATED <math>[CO_4]</math> TETRAHEDRA INTO FRAMEWORK AND LAYERS</b> <i>Sagatova D.N., Gavryushkin P.N.</i> 39	39
<b>FEATURES OF THE TRACE ELEMENT COMPOSITION OF MANTLE-DERIVED CHROME SPINELS</b> <i>Serebriannikov A.A., Logvinova A.M.</i> .....	41

## MINERAL COMPOSITION OF SABAKTY LAKE SEDIMENTS

*Kuzina D.M, Yusupova A.R., Nourgalieva N.G., Rogov A.M.*

Institute of Geology and Petroleum Technologies, Kazan Federal University, Kazan, Russia,  
yusupovaanast095@gmail.com

**Abstract.** In this study, we present lithology, mineralogy and scanning electron microscopy observations along a 294 cm sediment profile of the Lake Sabakty, with the objectives of understanding the geochemical processes and improving the paleohydrological and paleoecological information during the late Pleistocene and Holocene of the South Ural.

**Key words:** mineral composition, lake sediments, Holocene

Lake Sabakty (53°36'55" N; 58°39'22" E) located in the Bashkortostan republic; the length is ~2.3 km, width is ~1.0 km, average depth is 2.8 m; maximum depth is 6.0 m, the basin area is 9.0 km<sup>2</sup> [1-3]. The core No. 4 was selected for a detailed laboratory study based on the primary lithological description and the results of seismoacoustic studies [4]. The sediment was sliced at 2.0 cm thick samples. According to the radiocarbon dating, the age of the lake is ~22.5 thousand years [5].

The result of particle size analysis is presenting that the content of clay fraction in the sediment varies in the range 1.37-16.41 %, silt fraction are dominant at 33.24%-76.3%, sand - 12.02-65.39 %.

According to X-ray diffraction analysis (Fig. 1), the mineral composition is characterized by the predominance of terrigenous minerals (detrital quartz, microcline, mica, chlorite, mixed-layer clay minerals). Authigenic minerals include carbonates (calcite, dolomite), framboidal pyrite. Biogenic silica (cristobalite, tridymite) from diatom shells and stomatocysts are recorded throughout the sediment section. Scanning electron microscopy revealed the presence of framboidal pyrite, detrital and biogenic quartz in lacustrine sediments.


Fig. 1 shows, the authigenic component is inferior to the allothigenic component in content. However, in terms of variations along the section, it differs in a noticeably more differentiated behavior (especially carbonates).

The values of allothigenic component vary between 20.90%-61.14%, values of quartz ranges from 14.00% to 37.94%. The quartz content curve is similar to the amplitude of the allothigenic dominant, but with contrasting behavior at some points (since part of the crystalline quartz, together with amorphous silica (cristobalite + tridymite), is involved in the composition of diatoms and stomatocysts).

The carbonate component changes from 1.79 % to 52.12 %. The precipitation of calcite-dolomite carbonates is determined by factors: Ca/Mg-ratio in water, total carbonate alkalinity, salinity, pH value, temperature and organic productivity of the reservoir [6].

The content of pyrite changes from 1.18 % to 6.94 %. The formation of framboidal pyrite is associated with the recrystallization of amorphous Fe monosulfides, which appeared in the early diagenesis. The content of biogenic quartz increases up to section from 2.55 % to 25.70 %, indicating an increase in algae activity.

The comparison of the values variations of particle size analysis and mineralogical composition made it possible to reveal the features of climatic and other environmental changes on the studied lacustrine sediments.



**Figure 1 – Results of particle size and mineralogical analysis of Lake Sabakty sediments. Legend: 1 – clayey silt, 2 - silt, 3 – sandy silt, 4 - silty sand**

*Funding; This study was funded by the Russian Foundation for Basic Research, project 20-35-90058. Part of the study was funded with the subsidy allocated to the KFU in context of state assignment 671-2020-004.*

#### References:

1. Gareev A.M. Rivers and lakes of Bashkortostan. Ufa. Kitap. 2001. 259 p.
2. Gareev A.M. Rivers, lakes and swamp complexes of the Republic of Bashkortostan. Ufa. Gilem Publ. 2012. 248 p. (In Russian).
3. Gareev A.M. Sabakty. Bashkir Encyclopedia. Ch. ed. M.A. Ilgamov. Ufa. GAUN "Bashkir Encyclopedia". 2015-2020.
4. Krylov P.S, Nurgaliev D.K, Kuzina D.M. High-resolution seismic studies of shallow Lake Sabakty (Russia). International Multidisciplinary Scientific GeoConference Surveying Geology and Mining Ecology Management, SGEM. 2019. Vol. 19. No. 4.2. P. 229-234.
5. Yusupova A.R., Nurgalieva N.G., Batalin E.A., Gareev B.I. (2021):The first results of isotope geochemistry of bottom sediments of Lake Sabakty (Southern Ural). In Proceedings of the conference of young scientists Modern problems of geochemistry. Irkutsk: Institute of Geochemistry. A.P. Vinogradova. P. 36-38.
6. Deelman J.C. Low-temperature formation of dolomite and magnesite. Open-access e-book. 2011. 512 p.