


Sobolev Institute of Geology and Mineralogy SB RAS (IGM SB RAS)
Trofimuk Institute of Petroleum Geology and Geophysics SB RAS (IPGG SB RAS)
Novosibirsk State University (NSU)



X INTERNATIONAL SIBERIAN EARLY CAREER GEOSCIENTISTS CONFERENCE

13-17 June 2022, Novosibirsk

PROCEEDINGS OF THE CONFERENCE



MINERALOGY
2022



Sobolev Institute of Geology and Mineralogy SB RAS (IGM SB RAS)
Trofimuk Institute of Petroleum Geology and Geophysics SB RAS (IPGG SB RAS)
Novosibirsk State University (NSU)



**X INTERNATIONAL SIBERIAN EARLY CAREER
GEOSCIENTISTS CONFERENCE**

PROCEEDINGS OF THE CONFERENCE



MINERALOGY
2022

13-17 June 2022

Novosibirsk, Russia

УДК 55(061)
ББК Д 26
Т35

ORGANIZING COMMITTEE

Co-chairmen:

Dr. Nikolay N. Kruk,
Corresponding Member of RAS, Director of IGM SB RAS
Dr. Vyacheslav N. Glinskikh,
Corresponding Member of RAS, Director of IPPG SB RAS
Dr. Valery A. Vernikovsky,
Academician of RAS, Dean of the DGG NSU

Conference Secretariat:

Dr. Maria O. Shapovalova (IGM SB RAS)
Dr. Ekaterina A. Ovdina (IGM SB RAS)
Elena S. Shaparenko (IGM SB RAS)

Technical Editors:

Dr. Ekaterina A. Ovdina (IGM SB RAS)
Dr. Maria O. Shapovalova (IGM SB RAS)

X International Siberian Early Career GeoScientists Conference: Proceedings of the Conference (13-17 June 2022, Novosibirsk, Russia)/ IGM SB RAS, IPPG SB RAS, NSU. – Novosibirsk: IPC NSU, 2022. - 228 p. - ISBN 978-5-4437-1337-3.

The volume contains short papers of the X International Siberian Early Career GeoScientists Conference and offer information on multidisciplinary aspects of the Earth Science: petrology, geochemistry, geochronology, mineralogy, metallogeny, geodynamics and tectonics, sedimentology, geomorphology and neotectonics, paleontology and paleoclimatology, petroleum geosciences, economic geology, geoecology, hydrogeology, geophysics, mining engineering and geoinformation systems in the Earth Science.

The Conference was supported by IGM SB RAS, IPPG SB RAS, NSU, JSC "Siberian Anthracite", JSC "Almazy Anabara", Data East, LLC, PJSC "Gazprom Neft", Laboratory of ore-forming systems IGM SB RAS and SGA Siberian Student Chapter.

УДК 55(061)
ББК Д 26

ISBN 978-5-4437-1337-3

© Sobolev Institute of Geology and Mineralogy
SB RAS, 2022
© Trofimuk Institute of Petroleum Geology and
Geophysics SB RAS, 2022
© Novosibirsk State University, 2022

A HOLOCENE CLIMATIC RECORD INDICATED BY GEOCHEMICAL INDICATORS AND GRANULOMETRY FROM BANNOE LAKE IN THE SOUTHERN URAL <i>Yusupova A.R., Nourgalieva N.G., Kuzina D.M.</i>	128
--	-----

Section REGIONAL GEOLOGY AND TECTONICS

DEFORMATION OF THE SOUTHEASTERN PART OF THE TATARKA-ISHIMBA SUTURE ZONE (YENISEI RIDGE): STRUCTURAL AND ⁴⁰Ar/³⁹Ar GEOCHRONOLOGICAL DATA <i>Bogdanov E.A., Kadilnikov P.I., Matushkin N.Yu.</i>	130
---	-----

PETROGRAPHY, U-Pb DETRITAL ZIRCON GEOCHRONOLOGY AND PROVENANCE OF THE ILIKTA FORMATION OF THE SIBERIAN CRATON (WESTERN BAIKAL AREA) <i>Efremova U.S., Donskaya T.V.</i>	132
--	-----

COMPOSITION OF UST-BELSKY AND ALGAN TERRANES JURASSIC-CRETACEOUS TUFF-SANDSTONES (KORYAK HIGHLAND, NE RUSSIA) <i>Gushchina M.U., Moiseev A.V., Tuchkova M.I.</i>	134
---	-----

STRUCTURES AND COMPOSITIONS OF THE RESIDUAL DUNITE-HARZBURGITE COMPLEX OF THE NORTH BALKHASH OPHIOLITE ZONE (CENTRAL KAZAKHSTAN): GEOCHEMICAL AFFINITY AND TECTONIC SETTING OF FORMATION <i>Milyukova A.G., Skoblenko A.V.</i>	136
---	-----

GEODYNAMIC SETTINGS OF THE RIPHEAN MAGMATISM IN THE SOUTHEASTERN PART OF THE ANABAR SHIELD <i>Novikov V.S., Savelev A.D.</i>	138
---	-----

PALEOMAGNETISM AND GEOCHRONOLOGY OF LAMPROITES TOMTOR'S DYKES (UDZHA RIVER, SIBERIAN PLATFORM) <i>Pasenko A.M., Malyshev S.V., Pazuhina A.A.</i>	140
---	-----

KAZANIAN-STAGE CHROMSPINEL PLACERS IN THE SOUTHERN PRE-URALS, BASHKIRIA, RUSSIA <i>Rakhimov I.R.</i>	142
---	-----

HOW WIDE WAS ~1000-950 MA MAFIC MAGMATIC EVENT: DATA FROM THE SIBERIAN AND NORTH CHINA CRATONS? <i>Savelev A.D., Khudoley A.K., Malyshev S.V., Pazukhina A. A.</i>	144
---	-----

EVOLUTION OF THE AUSTRALIAN-ANTARCTIC PART OF THE SOUTH EASTERN INDIAN RIDGE IN PALEOCENE – EOCENE <i>Sergeeva V.M., Leitchenkov G.L.</i>	146
--	-----

MICROSTRUCTURAL ANALYSIS OF A TECTONIC FAULT (ON EXAMPLE OF THE PRIMORSKY FAULT OF THE BAIKAL RIFT ZONE) <i>Svecherevskiy A.D., Ustinov S.A., Ostapchuk A.A., Petrov V.A.</i>	148
--	-----

LITHOGEOCHEMICAL CHARACTERISTICS AND GEODYNAMIC OF THE UPPER PALEOZOIC YAKSHA FORMATION (WESTERN TRANSBAIKALIA) <i>Tashlykov V.S.</i>	150
--	-----

CONDITIONS AND FEATURES OF THE FORMATION OF METABASITE AND METATERRIGENOUS ROCKS OF THE TELETSK ZONE OF GORNY ALTAI <i>Zindobryi V.D., Buslov M.M.</i>	152
---	-----

A HOLOCENE CLIMATIC RECORD INDICATED BY GEOCHEMICAL INDICATORS AND GRANULOMETRY FROM BANNOE LAKE IN THE SOUTHERN URAL

Yusupova A.R., Nourgalieva N.G., Kuzina D.M.

Institute of Geology and Petroleum Technologies, Kazan Federal University, Kazan, Russia,
yusupovaanast095@gmail.com

Abstract. The work is devoted to the geochemical and granulometry study of the Bannoe lake sediments. The main aim of this investigation is to identify the events and trends of environmental and climate changes in the Late Quaternary. The present study of grain size and organic geochemistry of core sediments from Bannoe Lake provides paleoclimatological record of South Ural for the Holocene Period.

Key words: geochemical indicators, lacustrine sediments, particle size content, Holocene

The Bannoe Lake (53°35'48.13" N 58°37'47.28" E) is in the Southern Urals. The altitude of the lake is 438 m, the length is ~4.2 km, the width is ~1.9 km, and the basin area is 36.3 km² [1; 2; 3]. According to the radiocarbon dating the age of the lake is 12.5 thousand years [4]. For a detailed complex analysis, core No. 3 was selected according to the results of seismoacoustic studies [5]. The total core length is 508 cm. The sediment was sliced at 2 cm thick samples.

High resolution analyses of lake sediments retrieved from Bannoe Lake were conducted using a combination of variables including particle size content, total organic carbon (TOC), total nitrogen (TN), total inorganic carbon (TIC) and total sulfur (TS) contents, $\delta^{13}\text{C}$ value of organic matter ($\delta^{13}\text{C}_{\text{org}}$), and $\delta^{18}\text{O}$, $\delta^{13}\text{C}$ values of carbonate content ($\delta^{18}\text{O}_{\text{carb}}$ and $\delta^{13}\text{C}_{\text{carb}}$). The C/N ratio (TOC/TN) is regarded as an effective indicator of the origin of organic matter in sediments [6; 7]. Endogenous organics, rich in proteins and poor in cellulose, are characterized by C/N values from 4 to 10. Exogenous organics, poor in proteins and rich in cellulose (the remains of cell walls of higher (terrestrial) plants), are characterized by C/N values of more than 20 [8].

The result of particle size analysis is showing that the content of clay fraction in the sediment varies in the range (9.54-26.27) %, silt fraction are dominant at 56.39%-78.12%, sand - (2.76-26.71) % (Figure 1).

The TOC values changes from 1.56%-16.41%, the TN content ranged from 0.23%-2.23%. The TOC and TN indices increase up the section, which is explained by the increasing the biota activity and the organic matter accumulation. The TS values varied from 0.057% to 0.78%, which is typical for freshwater lakes [9]. The ratio C/N values vary between 5.98-17.79%; average value of C/N is 8.86 and indicates the dominance of endogenous biota. The greatest contribution of exogenous plants is noted at 408 cm (C/N=17.79) and 188 cm (C/N=13.00). The values of $\delta\text{C}_{\text{org}}$ vary from -27.65 ‰ to -24.22 ‰, which in combination with data on the C/N indicator (C/N~9), makes it possible to verify the mixed origin of organic matter in the studied lake sediments. The $\delta^{18}\text{O}_{\text{carb}}$ ranges from -22.25‰ to -5.31‰, the $\delta^{13}\text{C}_{\text{carb}}$ varies from -18.4‰ to 2.39‰, and the TIC values vary from ~0.03% to 4.86%. An increase in the values of $\delta^{18}\text{O}_{\text{carb}}$ (a rapid increase in the dryness of the climate) and $\delta^{13}\text{C}_{\text{carb}}$ (an increase in biomass) recorded at depths of 308-298 cm and 198-178 cm, indicating a decrease in the lake level.

The periods of high and low humidity in the Southern Urals identified during the study are consistent with the Blytt–Sernander classification [10; 11; 12]. The results of geochemical and particle size investigations made it possible to reveal the features of climatic and other environmental changes for the Holocene Period of South Ural.




Figure 1 – Results of particle size and isotope analysis of Lake Bannoe sediments. Legend: 1 – clayey silt, 2 - silt, 3 – sandy silt

This study was funded by the Russian Foundation for Basic Research, project #20-35-90058. Part of the study was funded by grant of the President of the Russian Federation for state support of young Russian scientists - candidates of sciences № MK-4100.2021.1.5.

References:

- Gareev A.M. Rivers and lakes of Bashkortostan. Ufa. Kitap. 2001. 259 p.
- Gareev E.Z. Geological natural monuments of the Republic of Bashkortostan. Ufa. 2004.
- Abdrakhmanov R.F. Popov V.G. Geochemistry and formation of underground waters of the Southern Urals. Ufa: Academy of Sciences of the Republic of Belarus, Gilem. Rev. ed. corresponding member RAS V.N. Puchkov. 2010. 420 p.
- Yusupova A., Kuzina D., Batalin G., Gareev B., Nourgalieva N. (2020): First Geochemical Data on Lacustrine Sediments, Lake Bannoe (Bannoe), Southern Urals. In Proceedings 4th Kazan Golovkinsky Stratigraphic Meeting 2020 Sedimentary Earth Systems: Stratigraphy, Geochronology, Petroleum Resources. Kazan. P.292-297.
- Krylov P.S., Nurgaliev D.K., Yasonov P.G., Dautov A.N., Golovtsov A.V., Sitdikov R.N., Krylova A.S. Seismoacoustic research of Lake Bannoe bottom sediments (South Ural, Russia). ARPN Journal of Engineering and Applied Sciences. 2020. Vol. 15. No. 1. P. 133–135.
- Meyers P.A. Applications of organic geochemistry to paleolimnological reconstructions: A summary of examples from the Laurentian Great Lakes. Organic Geochemistry. 2003. Vol. 34. P. 261–289.
- Routh J, Meyers P.A, Hjorth T, Baskaran M, Hallberg R. Sedimentary geochemical record of recent environmental changes around Lake Middle Marviken, Sweden. Journal Paleolimnology. 2007. Vol. 37. P. 529-545.
- Krishnamurthy R.V, Bhattacharya S.K, Kusumgar S. Palaeoclimatic changes deduced from $^{13}\text{C}/^{12}\text{C}$ and C/N ratios of Karewa lake sediments, India. Nature. 1986. Vol. 323. P. 150-152.
- Ding X., Li D., Zheng L., Bao H., Chen H., Kao S. Sulphur Geochemistry of a Lacustrine Record from Taiwan Reveals Enhanced Marine Aerosol Input during the Early Holocene. Scientific Reports. 2016. Vol. 6. Art. 38989
- Blytt A.G. Forsøg til enTheori om Indvandingen af Norges Flora Nyt Mag. Naturvid. Christiania (Oslo). Vol. 21. P. 279-362.
- Blytt A.G. Immigration of the Norwegian Flora. Cammermeyer. Oslo. Christiania. 89 p.
- Sernander R. Studier öfver den Gótländska vegetationens utvecklingshistora. Uppsala. Akademisk afhandling. 1984. 112 p.