

Discrete Mathematics, Algorithms and Applications
© World Scientific Publishing Company

**MINIMAL INSTANCES WITH NO WEAKLY STABLE MATCHING
FOR THREE-SIDED PROBLEM WITH CYCLIC
INCOMPLETE PREFERENCES**

Eduard Yu. Lerner*

*Department of Data Analysis and Operations Research
Institute of Computational Mathematics and Information Technologies
Kazan Federal University, Russia
eduard.lerner@gmail.com*

Regina E. Lerner

*Phystech School of Applied Mathematics and Informatics
Moscow Institute of Physics and Technology, Russia
ru.lerner@gmail.com*

Received 18 August 2021
Revised Day Month Year
Accepted Day Month Year
Published Day Month Year

Given n men, n women, and n dogs, each man has an incomplete preference list of women, each woman has an incomplete preference list of dogs, and each dog has an incomplete preference list of men. We understand a family as a triple consisting of one man, one woman, and one dog such that the dog belongs to the preference list of the woman, who, in turn, belongs to the preference list of the man, while the latter belongs to the preference list of the dog. We understand a matching as a collection of nonintersecting families (some agents, possibly, remain single). A matching is said to be nonstable, if one can find a man, a woman, and a dog which do not live together currently but each of them would become “happier” if they do. Otherwise, the matching is said to be stable (a weakly stable matching). We give an example of this problem for $n = 3$ where no stable matching exists. Moreover, we prove the absence of such an example for $n < 3$. Such an example was known earlier only for $n = 6$ (Biro, McDermid, 2010). The constructed examples also allows one to halve the size of the recently constructed analogous example for complete preference lists (Lam, Plaxton, 2019).

Keywords: stable matching; incomplete preferences lists; three-sided problem; cycles; weighted directed graphs.

Mathematics Subject Classification 2020: 05C22, 05C90, 91B68

*Corresponding author.

1. Introduction

Assume that there are n men and n women, and each one among them has a preference list of representatives of the opposite sex. A partition into heterogeneous families with no tuple of man and woman, who prefer each other rather than their partners (if they have ones), is called a stable matching. The initial case of complete preference lists was studied by D. Gale and L.S. Shapley: a stable matching necessarily exists, a quadratic time complexity algorithm for forming it was proposed in [4]. Note that the algorithm is also applicable for the case of incomplete preference lists, but some men and women, possibly, remain single. A certain modification of the Gale–Shapley algorithm (see, for example, [5]) allows one to find, if possible, a matching without single men and women or to prove its absence, otherwise.

In the case of random complete preference lists (when for each man and each woman the distribution of all permutations of representatives of the opposite sex is independent and uniform) the time necessary for finding a stable matching is $\Theta(n \ln(n))$ [5]. In this case, the mean value of the number of stable matchings also has the asymptote $n \ln(n)$ [9].

In [5] D. Knuth states the question whether it is possible to generalize the theory of stable matchings to the case of three genders. The most interesting variant in the k -gender case occurs when preferences are cyclic: representatives of the 1st gender rank representatives of the 2nd one, the latter rank representatives of the 3rd gender, etc., and each representative of the k th gender has a preference list of representatives of the 1st gender (see [7, Chapter 5.6] for the non-cyclic variants of the k -gender case).

A tuple containing exactly one representative of each gender is called a family, and a set of disjoint families is called a matching. A matching is said to be weakly stable, if there is no tuple outside this matching, each member of which would become “happier”, if they live together. In what follows, for brevity, we use the term “a stable matching” instead of the term “a weakly stable matching”.

Let the number of representatives of each gender equal n . Authors of the paper [2] prove that with complete preference lists a stable matching always exists, provided that $n \leq k$ (where k is the number of genders). In [3], Eriksson et al. generalize this result for the case when $k = 3$ and $n = k + 1 = 4$. Ibid, they state the conjecture that the problem of finding a stable matching in a 3-gender case with complete preference lists (problem 3-DSM-CYC or just 3DSM) has a solution for any n . Using a satisfiability problem formulation and an extensive computer-assisted search, the authors of [8] prove the validity of the conjecture stated by Eriksson et al. for $n = 5$. In [10], B. Pittel proves that with random preference lists the mean value of stable matchings in problem 3DSM grows as $\Omega(n^2 \ln^2(n))$.

The 3DSMI-problem (3-dimensional stable matching with incomplete preference lists) was studied by P. Biró and E. McDermid [1]. According to results a solution of 3DSMI does not necessarily exists in contrast to the two-dimensional case; they give an explicit example of problem 3DSMI for $n = 6$ with no stable matching.

Moreover, they prove that the problem of establishing the solvability of 3DSMI is NP-complete. Ibid, they state the problem of constructing an instance with no weakly stable matching for $n < 6$.

Finally, contrary to expectations, the conjecture stated by Eriksson et al. was recently refuted in [6]. Lam and Paxton associate problem 3DSMI with a certain problem 3DSM, where n is 15 times greater than the initial size; this problem is solvable if and only if so is the initial problem 3DSMI. Therefore, the problem of establishing the solvability of problem 3DSM is NP-complete. The example described in the paper [1] allows one to construct an instance of problem 3DSM for $n = 90 = 6 \times 15$ with no stable matching.

For this reason, the problem of finding an instance of 3DSMI with no weakly stable matching for $n < 6$ becomes more salient. The construction of such instances for the least possible values of n is the goal of this paper.

First we constructed an instance of 3DSMI problem for $n = 4$ and proved the absence of such instances for $n < 3$. But after failing to prove the absence of such instances for $n = 3$, we have proposed an algorithm for the computer search of all possible instances of 3DSMI problems for $n = 3$. Unexpectedly, the algorithm has succeeded in constructing instances of rather simple 3DSMI problems without weakly stable matching for $n = 3$.

The rest part of the paper has the following structure. In Sect. 2, we present the formal definitions of 3DSMI-CYC in terms of the graph theory. In Sect. 3, we study some properties of graphs of problem 3DSMI-CYC, prove the absence of counterexamples for $n < 3$. In Sect. 4 we describe various cases of problem 3DSMI for $n = 3$ and consider the result of their computer enumeration. We consider several instances and explicitly prove the absence of a stable matching for each of them, and describe general properties of all counterexamples. In Sect. 5, we conclude by mentioning some potential future work.

2. The statement of 3DSMI-CYC in terms of the graph theory

Let G be some directed graph. Denote the set of its edges by E ; assume that no edges are multiple. Let the vertex set $V(G) = V$ of the graph G be divided into three subsets, namely, the set of men M , women F , and dogs D . Any edge (v, v') , $v, v' \in V$ of this graph is considered to be of one of three types: either $v \in M, v' \in F$, or $v \in F, v' \in D$, or $v \in D, v' \in M$.

Evidently, the length of each cycle in the graph G is a multiple of 3. The graph G is said to be *trivial* if it has no cycles whose length exactly equals 3. In this paper, we mostly consider graphs to be untrivial.

Assume that $|M| = |F| = |D|$ (otherwise we supplement the corresponding subgraph with vertices that are not connected with the rest part of the graph). The number $n = |M| = |F| = |D|$ is called the problem *size*. Evidently, the length of all cycles in the graph G is a multiple of 3. Note also that this condition ensures the possibility to divide the vertex set of any digraph G into 3 subsets M, F, D

4 *Eduard Yu. Lerner and Regina E. Lerner*

so that all its edges are directed as indicated above. Each edge (v, v') , $v, v' \in V$, corresponds to some positive integer $r(v, v')$ which is called the rank of this edge. For fixed $v \in V$, all possible ranks $r(v, v_1), \dots, r(v, v_k)$ coincide with $\{1, \dots, k\}$, where k is the outdegree of the vertex v (if $r(v, v') = 1$, then v' is the best preference for v , and so on).

Using the language of the graph theory, for convenience, we define the notion of a matching for a graph G in terms of a special spanning subgraph H of this graph. Each vertex $v \in V$ of the subgraph H has at most one outgoing edge and the following condition is fulfilled: if a vertex v has an outgoing edge, then this edge belongs to a cycle of length 3 in the graph H . Cycles of length 3 in the subgraph H are called *families*. Evidently, each family, accurate to a cyclic shift, takes the form (m, f, d) , where $m \in M$, $f \in F$, and $d \in D$. Note that in what follows, for convenience of denotations of families, we do not fix the order of genders in a family, i. e., we treat denotations of families as triples derived from an initial one by a cyclic shift as equivalent.

In what follows, we sometimes use the notion of a family in a wider sense, namely, as any cycle of length 3 in the graph G . However, if some subgraph H is fixed, then we describe other cycles of length 3 explicitly, applying the term “a family” only to cycles that enter in H .

A *matching* μ is a collection of all families of the subgraph H . This definition is traditional; though we use it, we, for clarity, we illustrate our considerations with the help of the subgraph $H(\mu)$. For a vertex v , $v \in V$, in the matching μ , the rank $R_\mu(v)$ is defined as the rank of the edge that goes out of this vertex in the subgraph H . If some vertex v in the subgraph H has no outgoing edge, then $R_\mu(v)$ is set to $+\infty$.

A triple (v, v', v'') is said to be *blocking* for some matching μ , if it is a cycle in the graph G , and

$$r(v, v') < R_\mu(v), \quad r(v', v'') < R_\mu(v'), \quad r(v'', v) < R_\mu(v'').$$

A matching μ is said to be *stable*, if no blocking triple exists for it.

Problem *3DSMI* (3-dimensional stable matching with incomplete preference lists) consists of finding a stable matching for a given graph G . It is well known that it does not necessarily exists. Moreover, the problem of establishing its existence for a given graph G is NP-complete. It was mentioned in the Introduction, this fact was proved by Biro and McDermid. They have constructed an explicit example of the graph G of size 6, for which no stable matching exists. Moreover, the question of constructing similar examples for lesser sizes was also stated by the mentioned authors.

3. The absence of examples of problem 3DSMI with no stable matching for $n < 3$

Let G and G' be two directed graphs defined on one and the same vertex set V but, generally speaking, having distinct edge sets. Assume that rank functions r_G and

$r_{G'}$ are defined on E and E' , correspondingly. Let $L \subseteq E \cap E'$. We say that ranking orders r_G and $r_{G'}$ coincide on L , if for any two edges $(v, v'), (v, v'')$ in L ,

$$r_G(v, v') < r_G(v, v'') \iff r_{G'}(v, v') < r_{G'}(v, v'').$$

Lemma 3.1. *For any untrivial graph G of problem 3DSMI of size n there exists a graph G' of the same size such that the outgoing degree of each its vertex is nonzero and there is the following correspondence between graphs G and G' :*

- 1) *the set of all possible families of graphs G and G' coincide;*
- 2) *the ranking order of all edges that enter in these families also coincide.*

Proof. Let v be a vertex in the graph G having no outgoing edges. Then v enters in no family of the graph G . Let us delete this vertex together with all incoming edges in v . Repeating this procedure several times, we get a graph \widehat{G} such that each its vertex has at least one outgoing edge and its set of families coincides with that of the initial graph G . Let the symbol \widehat{V} stand for the vertex set of the graph \widehat{G} , denote the set of its edges by \widehat{E} . According to untriviality of graph G the set of families of graphs G and \widehat{G} is nonempty. In this case, the set \widehat{V} contains at least one vertex for each gender.

Let us now restore the initial vertices belonging to the set $V \setminus \widehat{V}$ and for each of them arbitrarily construct at least one edge directed to some vertex in \widehat{V} that corresponds to a proper gender. Since the incoming degree of restored vertices equals zero, they, as earlier, can enter in no family. Note that $\widehat{E} \subseteq E$ and, consequently, one can construct a rank function for the obtained graph G' preserving the ranking order of the graph G on \widehat{E} . The obtained graph G' with the rank function defined in the indicated way is the desired one. \square

Lemma 3.1 allows one, when studying problems 3DSMI of size n , to restrict oneself to considering the corresponding graphs G with nonzero outgoing degrees of all vertices, which we do in what follows.

Let the symbol G'' stands for a subgraph of the graph G consisting of its edges of rank 1. We call G'' the *basic subgraph* of the graph G . Since each vertex in the basic subgraph has exactly one outgoing edge, G'' represents a collection of cycles, whose lengths are multiples of 3, and trees of edges that lead to these cycles.

Theorem 3.2. *Problem 3DSMI of size $n \leq 2$ always has a stable matching.*

Proof. Note that with $n = 1$ the assertion of the lemma is trivial. In what follows, we restrict ourselves to considering the case of $n = 2$. Note also that in this case a nonstable matching can contain only one family. So let us assume the absence of two family matching for graph G .

The basic subgraph of the graph G contains cycles either of length 3 or of length 6. Let us consider both cases sequentially. In the first case, there exist vertices v_0, v_1, v_2 such that $r(v_0, v_1) = r(v_1, v_2) = r(v_2, v_0) = 1$. Therefore, the family

(v_0, v_1, v_2) is a stable matching. It remains to consider the case when the basic subgraph of the graph G is a cycle of length 6, i.e., $C = (v_0, v_1, \dots, v_5)$. Without loss of generality, we assume that the graph G , which represents a counterexample to Theorem 3.2, along with the cycle C contains the edge (v_2, v_0) of rank 2. Then the only possible blocking triple to the matching of one family (v_0, v_1, v_2) is (v_2, v_3, v_4) . Consequently, the graph G also contains the edge (v_4, v_2) . But then the only possible blocking triple for the matching consisting of one family (v_2, v_3, v_4) is (v_4, v_5, v_0) . In turn, the graph G that consists of only a basic cycle C and edges (v_0, v_4) , (v_4, v_2) , (v_2, v_0) of rank 2 has a stable matching consisting of one family (v_0, v_4, v_2) . Therefore, the graph G , along with the cycle C , contains at least 4 edges. Consequently, the graph G of size $n = 2$ has a matching of two families, and it is stable by definition. \square

4. The examples of graphs G of size $n = 3$ with no stable matching

In this section, we consider the case of $n = 3$. Let us first classify all graphs of the problem of this size; this will facilitate their computer search.

If the basic subgraph of the graph G contains cycles of length 3, then there exist vertices v_0, v_1, v_2 such that $r(v_0, v_1) = r(v_1, v_2) = r(v_2, v_0) = 1$. Therefore, if the family (v_0, v_1, v_2) enters in a matching, then these vertices can enter in no blocking triple. By deleting vertices $\{v_0, v_1, v_2\}$ from the graph G , we get the graph of 3DSMI with $n = 2$. By Theorem 3.2, this problem has a stable matching μ . Then $\mu \cup \{(v_0, v_1, v_2)\}$ is stable in G .

Therefore, the basic subgraph of the graph G represents either a cycle of length 9, or a cycle of length 6 with three edges that lead to this cycle. Altogether, accurate to the cyclic symmetry, there are 6 such subgraphs; they are shown in Fig. 1

Each of 9 vertices of these subgraphs in the graph G can have outgoing edges that lead to two remaining vertices of the corresponding gender (here we understand remaining vertices as those that differ from the vertex, to which the edge of the basic graph G'' is already directed). Generally speaking, the total number of possible cases is 5, namely,

- (1) the considered vertex has no more outgoing edges;
- (2)–(3) the considered vertex has one more outgoing edge that leads to some vertex among two ones;
- (4)–(5) the considered vertex has two edges, their ranks are equal to 2 and 3, we can associate ranks with these edges in two ways.

Therefore, it suffices to consider 6×5^9 problems 3DSMI.

Evidently, for each of these problems there exist at most 27 families (27 blocking triples). The number of possible matching μ as one can easily calculate, also is not so large. Namely, there exist at most 27 matchings consisting of one family. In addition, there exist at most 108 matchings consisting of two families, namely, there are 27 ways to form a triple consisting of representatives of genders that enter in no matching, and 4 ways to choose partners among two women and two dogs entering




Fig. 1. 6 variants of the basic subgraph of the graph G .

the matching for a fixed man that also enters this matching. Finally, the number of matchings of 3 triples is not less than 36, because there are $3! \times 3!$ ways to distribute women and dogs among three man-indexed triplets. Therefore, the total amount of matchings does not exceed $36 + 108 + 27 = 171$. For each of them we need to find the first triple among 27 potential blocking ones that really is blocking.

Therefore, the total amount of considered cases does not exceed $6 \times 5^9 \times 171 \times 27 \approx 54 \times 10^9$. For generating these cases, we have written a program in Python. See the version of this program that calculates the number of counterexamples for each of basic graphs shown in Fig. 1 at <https://github.com/reginalerner/3dsm/>.

For the first basic graph shown in Fig. 1 (a cycle formed by 9 vertices), no such graph for problem 3SDMI with no stable matching was obtained. As is mentioned in the Introduction, we even did not expect to find such instances for $n = 3$. To our surprise, the computer search has found such counterexamples for each of the rest basic graphs. One of them is shown in Fig. 2. For convenience, we enumerate vertices of the graph by numbers v , $v = 0, 1, \dots, 8$. The value $v \bmod 3$ defines the gender that corresponds to the vertex v .

Theorem 4.1. *There is no stable matching in 3SDMI for the graph shown in Fig. 2.*

Proof. Fig. 2 evidently demonstrates that each possible cycle of length 3 takes one of the following forms: $(0, 1, 5)$, $(0, 7, 8)$, $(1, 2, 3)$, $(1, 5, 3)$, $(2, 3, 4)$, $(3, 4, 5)$, and $(4, 8, 6)$. These cycles form families, while collections of disjoint families form matchings μ in the problem.

Evidently, if one can add a cycle (v, v', v'') to a matching μ (vertices v, v', v'' do not enter in μ), then μ is unstable, i.e., the triple (v, v', v'') is blocking for μ . Therefore, candidates for stable matchings should be supplemented with possible




Fig. 2. The graph of problem 3DSMI of size 3 with no stable matching consisting of 16 edges. The rank of all edges indicated by solid bold lines equals 1. The dashed lines represent the edges with the rank 2. The rank of the “dotted” edge equals 3. Vertices $\{2, 5, 7\}$ correspond to men, vertices $\{0, 3, 6\}$ correspond to women, and vertices $\{1, 4, 7\}$ correspond to dogs.

cycles. We call such matchings *incompletable* and consider only such ones.

The union of vertices of three cycles that are listed above does not coincide with the set of all vertices of the graph shown in Fig. 2. On the other hand, by using the direct search method we can prove that any set consisting of one triple is completable. Therefore, each incompletable matching consists of two families. Below we give their complete list together with blocking triples:

- (1) $\{(0, 1, 5), (2, 3, 4)\}$, the blocking triple is $(4, 8, 6)$;
- (2) $\{(0, 1, 5), (4, 8, 6)\}$, the blocking triple is $(1, 2, 3)$;
- (3) $\{(0, 7, 8), (1, 2, 3)\}$, the blocking triple is $(3, 4, 5)$;
- (4) $\{(0, 7, 8), (1, 5, 3)\}$, the blocking triple is $(2, 3, 4)$;
- (5) $\{(0, 7, 8), (2, 3, 4)\}$, the blocking triple is $(0, 1, 5)$;
- (6) $\{(0, 7, 8), (3, 4, 5)\}$, the blocking triple is $(0, 1, 5)$;
- (7) $\{(1, 2, 3), (4, 8, 6)\}$, the blocking triple is $(0, 7, 8)$ or $(3, 4, 5)$;
- (8) $\{(1, 5, 3), (4, 8, 6)\}$, the blocking triple is $(0, 7, 8)$. □

One can easily give other examples of graphs with the same set of cycles, incompletable matchings, and blocking triples. In particular, this property is characteristic for the graph that differs from that shown in Fig. 2 by the presence of the additional edge $(7, 2)$ of rank 2 or the additional edge $(6, 1)$ of rank 2, or both of these edges.

Moreover, one can find other graphs consisting of 16 edges that have no stable matching. One of them is shown in Fig. 3 (any other graph with this property differs from the indicated one only in the fact that ranks of edges $(0, 4)$ and $(0, 7)$ have interchanged). These graphs define the following families of forming matchings in problem 3DSMI: $(0, 1, 8)$, $(0, 4, 5)$, $(0, 7, 5)$, $(1, 2, 6)$, $(2, 3, 7)$, $(3, 4, 5)$, and $(3, 7, 5)$.

The list of matchings with blocking triples looks as follows:

- (1) $\{(0, 1, 8), (2, 3, 7)\}$, the blocking triple is $(3, 4, 5)$;
- (2) $\{(0, 1, 8), (3, 4, 5)\}$, the blocking triple is $(1, 2, 6)$;
- (3) $\{(0, 1, 8), (3, 7, 5)\}$, the blocking triple is $(1, 2, 6)$;
- (4) $\{(0, 4, 5), (1, 2, 6)\}$, the blocking triple is $(2, 3, 7)$;
- (5) $\{(0, 4, 5), (2, 3, 7)\}$, the blocking triple is $(0, 1, 8)$;




Fig. 3. One more graph of 3DSMI of size 3 with no stable matching which contains 16 edges. Denotations are the same as in Fig. 2.

- (6) $\{(0, 7, 5), (1, 2, 6)\}$, the blocking triple is $(2, 3, 7)$;
- (7) $\{(1, 2, 6), (3, 4, 5)\}$, the blocking triple is $(0, 7, 5)$.
- (8) $\{(1, 2, 6), (3, 7, 5)\}$, the blocking triple is $(0, 4, 5)$.

Since the counterexamples considered above are diverse, they have some common properties. We are going to describe them in a future paper.

5. Concluding remarks

In this paper, we study the problem stated by Biro and McDermid in [1], namely, we seek for instances with no weakly stable matching for 3DSM-CYC with $n < 6$. In particular, we find the minimal value of n , with which such instances exist, and describe some of them.

The idea of this study is due to the work of Lam and Paxton [6], who give an example of problem 3DSM-CYC for $n = 90$ with no stable matching. This example is based on an analogous example proposed by Biro and McDermid for problem 3DSMI-CYC with $n = 6$. Our example constructed for problem 3DSMI with $n = 3$ allows one to make the size of an example for problem 3DSM with no stable matching as low as $n = 45$. According to results obtained in Sect. 3, the further decrease of n for problem 3DSMI is impossible. However, it seems possible to find problem 3DSM with no stable matching with $n < 45$ using some other methods.^a

Actually, Lam and Paxton studied not only 3-DSM-CYC, but also its k -gender analog, k -DSM-CYC, for arbitrary $k \geq 3$. First they have represented problem 3-DSMI-CYC as a particular case of k -DSMI-CYC with n^2 representatives of each gender. Then by the reduction from k -DSMI-CYC they have proved that k -DSM-CYC is NP-complete.

Note that some development of ideas proposed in the paper [6] allows one to rather easily construct a counterexample of size $n = 5$ for k -DSMI-CYC, $k > 3$, basing on the graph shown in Fig. 2 via the subdivision of edges that go out of

^aNote in the revised paper: in arXiv:2107.10102v3 [math.CO] we reduced the size of the counterexample to $n = 20$.

vertices $\{0, 3, 6\}$ which correspond to women. Any of subdivided edges is converted to the chain with $k - 3$ vertices inside, one for each new gender (see Fig. 4). A




Fig. 4. The graph (of size 5) of 4DSMI-CYC with no stable matching. In 4DSMI-CYC, we assume that there exist 4 genders, for example, men, women, kids, and dogs, while women are partial to kids, the latter are partial to dogs, and preferences of men and dogs are similar to those described earlier in the definition for 3DSMI. The definition of the graph of 4DSMI-CYC is also completely analogous; the vertices that correspond to kids are marked with the bold dot symbol. Recall that the rank of all edges indicated by solid bold lines equals 1. The dashed lines represent the edges with the rank 2. The rank of the “dotted” edge equals 3.

k -gender family should contain the new vertices from subdivided edge, so there is a biunique correspondence between new k -gender families and old 3-gender ones. If no stable matching exists for 3-gender families, then neither one exists for the new k -gender graph.

Therefore, for any $k > 2$, we have constructed an instance of problem k -DSMI-CYC with $n = 5$ with no stable matching (where lists of preferences of two women, two men, and two dogs that are not shown in Fig. 2 can be arbitrary). The question about the existence of such counterexamples for $n = 3$ and $n = 4$ still remains open.

We hope that this work can be useful in studying other questions related to other aspects of the generalization of the theory of stable matchings to the k -dimensional case, $k > 2$. In our opinion, this study is far from completion.

Acknowledgments

The authors are grateful to Olga Kashina for her help in the translation of this paper, and to the (anonymous) reviewer for his/her careful reading of this work and for useful comments. We have taken them into account when preparing the final version of the text.

References

- [1] P. Biró and E. McDermid, Three-sided stable matchings with cyclic preferences, *Algorithmica* **58** (2010) 5–18.
- [2] E. Boros, V. Gurvich, S. Jaslar and D. Krasner, Stable matchings in three-sided systems with cyclic preferences, *Discrete Math.* **289**(13) (2004) 1–10.

- [3] K. Eriksson, J. Söstrand and P. Strimling, Three-dimensional stable matching with cyclic preferences, *Math. Soc. Sci.* **52** (2006) 77–87.
- [4] D. Gale and L.S. Shapley, College admissions and the stability of marriage, *Am. Math. Mon.* **69** (1962) 9–15.
- [5] D. E. Knuth, *Stable marriage and its relation to other combinatorial problems: an introduction to the mathematical analysis of algorithms*. Ser. CRM Proceedings and Lecture Notes (American Mathematical Society, 1996).
- [6] C.-K. Lam and C.G. Plaxton, On the Existence of Three-Dimensional Stable Matchings with Cyclic Preferences, in: *Algorithmic Game Theory*. Ser. Lecture Notes in Computer Science, Vol. 11801 (Springer, 2019), pp. 329–342.
- [7] D.F. Manlove, *Algorithmics of matching under preferences*. Ser. Theor. Comput. Sci. (World Scientific, 2013).
- [8] K. Pashkovich and L. Poirrier, Three-dimensional stable matching with cyclic preferences, *Optimization Letters*, **2020**(3) (2020) 1–9.
- [9] B. Pittel, The average number of stable matchings, *SIAM J. Discrete Math.* **2** (1989) 530–549.
- [10] B. Pittel, On random stable matchings: Cyclic ones with strict preferences and two-sided ones with partially ordered preferences, *Advances in Appl. Math.* **120** (2020) 1–27.