



Contents lists available at ScienceDirect

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laa

Locally algebraic linear operators and their centralizers



Adel Abyzov*, Alexander Maklakov

ARTICLE INFO

Article history:

Received 7 November 2022

Accepted 27 December 2022

Available online 2 January 2023

Submitted by P. Semrl

MSC:

15A04

16S50

16D70

16W80

Keywords:

Locally algebraic linear operators

Finite topologies

Principal ideal domain

Discrete valuation ring

Torsion module

ABSTRACT

In this paper, using finite topologies defined on the algebra of linear operators, we investigate centralizers and double centralizers of locally algebraic linear operators. In particular, for an arbitrary locally algebraic operator A , we establish the equality $CC(A) = \overline{F[A]}$ and the conditions under which the equality $CC(A) = C(A)$ is fulfilled. Besides, in the case of an algebraically closed field, we describe minimal locally algebraic linear operators.

© 2022 Elsevier Inc. All rights reserved.

1. Introduction

Let M be the right R -module. For any finite subset X of the module M , let us denote by U_X the set $\{f \in \text{End}(M) \mid f(X) = 0\}$. Then a *finite topology* on $\text{End}(M)$ is defined as a topology whose base consists of all sets of the form $f + U_X$, where $f \in \text{End}(M)$, and X is a finite subset of M . The finite topology on $\text{End}(M) \subseteq M^M$ is a topology

* Corresponding author.

E-mail address: aabyzov@kpfu.ru (A. Abyzov).