

HYPONORMAL MEASURABLE OPERATORS AFFILIATED WITH A SEMIFINITE VON NEUMANN ALGEBRA. IV

A. M. Bikchentaev

UDC 517.983+517.986

Abstract—Let τ be a faithful normal semifinite trace on a von Neumann algebra $\mathcal{M}$ of operators. For a normal operator A in $\mathcal{M}$, a condition on a τ -integrable operator B is found under which the operator $A + B$ is normal. For an operator whose square is τ -integrable, equivalent conditions for its normality are established in terms of trace inequalities. For an operator in $\mathcal{M}$, a criterion for hyponormality is found in terms of trace inequalities. It is shown that, given an arbitrary natural n , the power $(PQ)^n$ of the product of projections P and Q in $\mathcal{M}$ is hyponormal if and only if $PQ = QP$. Operator inequalities are obtained for powers of hyponormal contractions. It is shown that every natural power of a hyponormal partial isometry is a hyponormal partial isometry with the same initial space.

DOI: 10.1134/S0037446626010027

Keywords: Hilbert space, von Neumann algebra, normal trace, measurable operator, hyponormal operator

1. Introduction

Hyponormal bounded operators on a Hilbert space have been studied by many authors (see, for instance, [1–10] and the references therein). In the context of semifinite von Neumann algebras, the author published the papers [11–19] devoted to properties of (unbounded) τ -measurable hyponormal operators (see also [20]). Let $\mathcal{M}$ be a von Neumann algebra of operators acting in a Hilbert space $\mathcal{H}$, let $\mathcal{M}^{\text{pr}}$ be the lattice of projections in $\mathcal{M}$, and let τ be a faithful normal semifinite trace on $\mathcal{M}$. We list the main results of the paper; some of them are new even in the case $\mathcal{M} = \mathcal{B}(\mathcal{H})$ with $\tau = \text{tr}$.

- Let $A \in \mathcal{M}$ be normal and let $B \in L_1(\mathcal{M}, \tau) \cap L_2(\mathcal{M}, \tau)$. If the operator $A + B$ is hyponormal, then $A + B$ is normal (Theorem 1).
- For an operator $A \in L_2(\mathcal{M}, \tau)$, the following conditions are equivalent (see Theorem 2):
 - (i) A is normal;
 - (ii) $\tau(PA^*AP) \geq \tau(PAA^*P)$ for all $P \in \mathcal{M}^{\text{pr}}$;
 - (iii) $\tau(PA^*AP) \leq \tau(PAA^*P)$ for all $P \in \mathcal{M}^{\text{pr}}$.
- $A \in \mathcal{M}$ is hyponormal if and only if $\tau(PA^*AP) \geq \tau(PAA^*P)$ for all $P \in \mathcal{M}^{\text{pr}}$ with $\tau(P) < +\infty$.
- If $A \in \mathcal{B}(\mathcal{H})_1$ is hyponormal, then $(A^{*n}A^n)^{1/4} \geq (A^*)^{n-2}A^{n-2}$ for all $n \in \mathbb{N}$ with $3 \leq n$ (Theorem 4).
- Given $P, Q \in \mathcal{B}(\mathcal{H})^{\text{pr}}$ and $n \in \mathbb{N}$, the operator $(PQ)^n$ is hyponormal if and only if $PQ = QP$ (Theorem 5).
- If a partial isometry $U \in \mathcal{B}(\mathcal{H})$ is hyponormal, then U^n is also a hyponormal partial isometry and $U^{*n}U^n = U^*U$ for each $n \in \mathbb{N}$ (Corollary 5).

Apparently, some of our assertions extend to locally measurable operators from [21].

2. Definitions and Notation

Let $\mathcal{M}$ be a von Neumann algebra of operators on a Hilbert space $\mathcal{H}$, let $\mathcal{M}^{\text{pr}}$ be the lattice of projections ($P = P^2 = P^*$) in $\mathcal{M}$, let I be the unit of $\mathcal{M}$, and let $P^\perp = I - P$ for $P \in \mathcal{M}^{\text{pr}}$. Let $\mathcal{M}^+$ be the cone of positive elements in $\mathcal{M}$, let $\|\cdot\|$ be the C^* -norm on $\mathcal{M}$, and let $\mathcal{M}_1 = \{X \in \mathcal{M} : \|X\| \leq 1\}$ be the unit ball of $\mathcal{M}$.

Given $P, Q \in \mathcal{M}^{\text{pr}}$, we write $P \sim Q$ (Murray–von Neumann equivalence) if $P = U^*U$ and $Q = UU^*$ for some $U \in \mathcal{M}$.

For $(P_n)_{n=1}^\infty \subset \mathcal{M}^{\text{pr}}$, the infimum $\bigwedge_{n=1}^\infty P_n \in \mathcal{M}^{\text{pr}}$ is defined by $(\bigwedge_{n=1}^\infty P_n)\mathcal{H} = \bigcap_{n=1}^\infty P_n\mathcal{H}$.

An operator $V \in \mathcal{M}$ is an *isometry* if $V^*V = I$. An operator $V \in \mathcal{M}$ is a *coisometry* if V^* is an isometry.

A mapping $\varphi : \mathcal{M}^+ \rightarrow [0, +\infty]$ is called a *trace* if $\varphi(X + Y) = \varphi(X) + \varphi(Y)$, $\varphi(\lambda X) = \lambda\varphi(X)$ for all $X, Y \in \mathcal{M}^+$ and $\lambda \geq 0$ (with $0 \cdot (+\infty) \equiv 0$), and $\varphi(Z^*Z) = \varphi(ZZ^*)$ for all $Z \in \mathcal{M}$. A trace φ is called

- *faithful* if $\varphi(X) > 0$ for all $X \in \mathcal{M}^+$, $X \neq 0$;
- *normal* if $X_i \nearrow X$ ($X_i, X \in \mathcal{M}^+$) implies $\varphi(X) = \sup \varphi(X_i)$;
- *semifinite* if $\varphi(X) = \sup\{\varphi(Y) : Y \in \mathcal{M}^+, Y \leq X, \varphi(Y) < +\infty\}$ for all $X \in \mathcal{M}^+$ (see [22, Chapter V, Section 2; 23, Chapter 1, Section 1.15]).

An operator on $\mathcal{H}$ (not necessarily bounded or densely defined) is *affiliated with the von Neumann algebra* $\mathcal{M}$ if it commutes with every unitary operator from the commutant $\mathcal{M}'$ of $\mathcal{M}$.

Further on, τ denotes a faithful normal semifinite trace on $\mathcal{M}$, and

$$\mathcal{M}_\tau^{\text{pr}} = \{P \in \mathcal{M}^{\text{pr}} : \tau(P) < +\infty\}.$$

A closed operator X affiliated with $\mathcal{M}$ whose domain $\mathcal{D}(X)$ is dense in $\mathcal{H}$ is called τ -*measurable* if for every $\varepsilon > 0$ there exists $P \in \mathcal{M}^{\text{pr}}$ such that $P\mathcal{H} \subset \mathcal{D}(X)$ and $\tau(P^\perp) < \varepsilon$. The set $S(\mathcal{M}, \tau)$ of all τ -measurable operators is a $*$ -algebra with respect to taking the adjoint, scalar multiplication, and the operations of strong addition and multiplication obtained by taking closures of the usual operations (see [24, Chapter IX; 23, Chapter 2, Section 2.3]). For a family $\mathcal{L} \subset S(\mathcal{M}, \tau)$, we denote by $\mathcal{L}^+$ and $\mathcal{L}^h$ the positive and Hermitian parts of $\mathcal{L}$, respectively. The partial order on $S(\mathcal{M}, \tau)^h$ induced by the proper cone $S(\mathcal{M}, \tau)^+$ is denoted by $\leq$. If $X \in S(\mathcal{M}, \tau)$ and $X = U|X|$ is the polar decomposition of X , then $U \in \mathcal{M}$ and $|X| = \sqrt{X^*X} \in S(\mathcal{M}, \tau)^+$.

An operator $A \in S(\mathcal{M}, \tau)$ is called *hyponormal* if $A^*A \geq AA^*$. It is called *cohyponormal* if the operator A^* is hyponormal.

By $\mu(t; X)$ we denote the *singular value function* of an operator $X \in S(\mathcal{M}, \tau)$, i.e., the nonincreasing right-continuous function $\mu(\cdot; X) : (0, +\infty) \rightarrow [0, +\infty)$ defined by

$$\mu(t; X) = \inf\{\|XP\| : P \in \mathcal{M}^{\text{pr}}, \tau(P^\perp) \leq t\}, \quad t > 0.$$

Lemma 1 [25]. *Let $X, Y \in S(\mathcal{M}, \tau)$ and let $A, B \in \mathcal{M}$. Then*

- (i) $\mu(t; X) = \mu(t; |X|) = \mu(t; X^*)$ for all $t > 0$;
- (ii) if $|X| \leq |Y|$, then $\mu(t; X) \leq \mu(t; Y)$ for all $t > 0$;
- (iii) $\mu(t; AXB) \leq \|A\| \|B\| \mu(t; X)$ for all $t > 0$;
- (iv) $\mu(t; f(|X|)) = f(\mu(t; X))$ for all continuous functions $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $f(0) = 0$ and all $t > 0$.

The set

$$S_0(\mathcal{M}, \tau) = \{X \in S(\mathcal{M}, \tau) : \mu(\infty; X) := \lim_{t \rightarrow \infty} \mu(t; X) = 0\}$$

of τ -compact operators is an ideal in $S(\mathcal{M}, \tau)$.

Let m be the Lebesgue measure on $\mathbb{R}$. The noncommutative Lebesgue L_p -space ($1 \leq p < +\infty$) associated with $(\mathcal{M}, \tau)$ can be defined by

$$L_p(\mathcal{M}, \tau) = \{X \in S(\mathcal{M}, \tau) : \mu(\cdot; X) \in L_p(\mathbb{R}^+, m)\}$$

with the norm $\|X\|_p = \|\mu(\cdot; X)\|_p$, $X \in L_p(\mathcal{M}, \tau)$. We denote by the same symbol τ the extension of the trace τ to the entire Banach space $L_1(\mathcal{M}, \tau)$.

If $\mathcal{M} = \mathcal{B}(\mathcal{H})$ is the $*$ -algebra of all bounded linear operators on $\mathcal{H}$ and $\tau = \text{tr}$ is the canonical trace, then $S(\mathcal{M}, \tau)$ coincides with $\mathcal{B}(\mathcal{H})$, and $S_0(\mathcal{M}, \tau)$ coincides with the ideal of compact operators $\mathcal{S}(\mathcal{H})$. The following holds:

$$\mu(t; X) = \sum_{n=1}^{\infty} s_n(X) \chi_{[n-1, n)}(t), \quad t > 0,$$

where $\{s_n(X)\}_{n=1}^{\infty}$ is the sequence of s -numbers of a compact operator X , and χ_A denotes the indicator function of a set $A \subset \mathbb{R}$ (see [26, Chapter II]). Then the space $L_p(\mathcal{M}, \tau)$ is the Schatten-von Neumann ideal $\mathcal{S}_p(\mathcal{H})$, $1 \leq p < +\infty$.

3. Main Results

Lemma 2 [24, Chapter IX, Theorem 2.13]. *If $X \in \mathcal{M}$ and $Y \in L_1(\mathcal{M}, \tau)$, then $XY, YX \in L_1(\mathcal{M}, \tau)$.*

Lemma 3 [27, Theorem 17]. *If $X, Y \in S(\mathcal{M}, \tau)^+$ and $XY, YX \in L_1(\mathcal{M}, \tau)$, then $\tau(XY) = \tau(YX)$.*

Theorem 1. *Let an operator $A \in \mathcal{M}$ be normal and let $B \in L_1(\mathcal{M}, \tau) \cap L_2(\mathcal{M}, \tau)$. If the operator $T := A + B$ is hyponormal, then T is normal.*

PROOF. Since $T^*T \geq TT^*$ and $A^*A = AA^*$, we have

$$A^*B + B^*A + B^*B \geq AB^* + BA^* + BB^*.$$

Therefore,

$$D := A^*B + B^*A + B^*B - AB^* - BA^* - BB^* \geq 0.$$

The terms A^*B , B^*A , AB^* , and BA^* belong to $L_1(\mathcal{M}, \tau)$ by Lemma 2, and $B^*B, BB^* \in L_1(\mathcal{M}, \tau)$ by the definition of the space $L_2(\mathcal{M}, \tau)$. Hence $D \in L_1(\mathcal{M}, \tau)^+$. By Lemma 3 we have

$$\tau(A^*B) = \tau(BA^*), \quad \tau(B^*A) = \tau(AB^*), \quad \tau(B^*B) = \tau(BB^*),$$

and therefore, by linearity of the extension of τ to $L_1(\mathcal{M}, \tau)$, we get $\tau(D) = 0$. Since this extension is faithful on the cone $L_1(\mathcal{M}, \tau)^+$, we conclude that $D = 0$. Thus $T^*T = TT^*$, and the operator T is normal. $\square$

Passing to adjoint operators, we obtain the following.

Corollary 1. *Let an operator $A \in \mathcal{M}$ be normal and let $B \in L_1(\mathcal{M}, \tau) \cap L_2(\mathcal{M}, \tau)$. If the operator $T := A + B$ is cohyponormal, then T is normal.*

Corollary 2 [28]. *Let an operator $A \in \mathcal{B}(\mathcal{H})$ be normal and let $B \in \mathcal{S}_2(\mathcal{H})$. If the operator $T := A + B$ is hyponormal (or cohyponormal), then T is normal.*

PROOF. The assertion follows from the inclusion $\mathcal{S}_2(\mathcal{H}) \subset \mathcal{S}_1(\mathcal{H})$.

Theorem 2. *For an operator $A \in L_2(\mathcal{M}, \tau)$, the following conditions are equivalent:*

- (i) A is normal;
- (ii) $\tau(PA^*AP) \geq \tau(PAA^*P)$ for all $P \in \mathcal{M}^{\text{pr}}$;
- (iii) $\tau(PA^*AP) \leq \tau(PAA^*P)$ for all $P \in \mathcal{M}^{\text{pr}}$.

PROOF. (i) $\Rightarrow$ (ii): Every normal operator $X \in S(\mathcal{M}, \tau)$ is hyponormal. Hence, from the inequality $A^*A \geq AA^*$ we get

$$PA^*AP \geq PAA^*P \quad \text{for all } P \in \mathcal{M}^{\text{pr}}.$$

Therefore, by monotonicity on the cone $L_1(\mathcal{M}, \tau)^+$ of the extension of τ to $L_1(\mathcal{M}, \tau)$, we obtain (ii).

(ii) $\Rightarrow$ (i): Assume that (ii) holds, but A is not normal. Since $L_2(\mathcal{M}, \tau) \subset S_0(\mathcal{M}, \tau)$, the operator A is not hyponormal by [11, Theorem 2.2]. Therefore, in the Jordan decomposition

$$X := A^*A - AA^* = X_+ - X_-, \tag{1}$$

with $X_+, X_- \in L_1(\mathcal{M}, \tau)^+$ and $X_+X_- = 0$, we have $X_- \neq 0$. Multiplying both sides of equality (1) on the left and on the right by the projection $P = \text{supp}(X_-)$, we get

$$PA^*AP - PAA^*P = -X_-. \tag{2}$$

Consequently, $\tau(PA^*AP) - \tau(PAA^*P) = -\tau(X_-) < 0$ since the extension of τ to $L_1(\mathcal{M}, \tau)$ is faithful on the cone $L_1(\mathcal{M}, \tau)^+$. This contradicts (ii).

Since an operator $X \in S(\mathcal{M}, \tau)$ is normal if and only if X^* is normal, we obtain (i) $\Leftrightarrow$ (iii). $\square$

Since the trace τ on the algebra $\mathcal{M}$ is semifinite, there exists a nonzero subprojection Q of the projection P such that $Q \in \mathcal{M}_\tau^{\text{pr}}$ and $QX_\perp Q \neq 0$ (see (2)). Therefore, each of the conditions

- $\tau(PA^*AP) \geq \tau(PAA^*P)$ for all $P \in \mathcal{M}_\tau^{\text{pr}}$;
- $\tau(PA^*AP) \leq \tau(PAA^*P)$ for all $P \in \mathcal{M}_\tau^{\text{pr}}$

implies that the operator $A \in L_2(\mathcal{M}, \tau)$ is normal.

An operator $T \in S(\mathcal{M}, \tau)$ is hyponormal (cohyponormal) if and only if $\mu(t; TP) \geq \mu(t; T^*P)$ (respectively, $\mu(t; T^*P) \geq \mu(t; TP)$) for all $t > 0$ and all $P \in \mathcal{M}_\tau^{\text{pr}}$ (see [17, Theorem 6]). Hence, by Lemma 1(i), (iv), we conclude the following:

*An operator $T \in S(\mathcal{M}, \tau)$ is hyponormal (cohyponormal) if and only if $\mu(t; PT^*TP) \geq \mu(t; PTT^*P)$ (respectively, $\mu(t; PTT^*P) \geq \mu(t; PT^*TP)$) for all $t > 0$ and all $P \in \mathcal{M}_\tau^{\text{pr}}$.*

Arguing in the same way as in the proof of Theorem 2, we verify the following statement:

For an operator $A \in \mathcal{M}$, the following conditions are equivalent:

- (i) *A is hyponormal;*
- (ii) *$\tau(PA^*AP) \geq \tau(PAA^*P)$ for all $P \in \mathcal{M}_\tau^{\text{pr}}$.*

Theorem 3. *Let an operator $A \in \mathcal{M}$ be an isometry, let $B \in S_0(\mathcal{M}, \tau)$, and let $T := A + B$. If $TT^* \geq I$ and T is hyponormal, then T is normal.*

PROOF. By assumption, $P := AA^* \in \mathcal{M}^{\text{pr}}$. Since $T^*T \geq TT^* \geq I$, we have

$$A^*B + B^*A + B^*B \geq AB^* - BA^* - BB^* - P^\perp \geq 0. \quad (3)$$

In particular, $P^\perp \in S_0(\mathcal{M}, \tau)$. If $X, Y \in S(\mathcal{M}, \tau)^+$, $Y \neq 0$, and $X \geq \mu(\infty; X)I$, then there exists $s > 0$ such that $\mu(s; X) < \mu(s; X + Y)$ (see [29, Proposition 2.2]). The operator

$$X := AB^* - BA^* - BB^* - P^\perp$$

belongs to $S_0(\mathcal{M}, \tau)^+$ and satisfies $X \geq \mu(\infty; X)I = 0 \cdot I = 0$. Put

$$X + Y := A^*B + B^*A + B^*B$$

(see the left-hand side of inequality (3)). Assume that T is not normal, i.e., $Y \neq 0$. For an arbitrary $Z \in S(\mathcal{M}, \tau)^+$, we have

$$\mu(t; I + Z) = 1 + \mu(t; Z) \quad \text{for all } t > 0$$

(see the proof of [30, Theorem 7]). By Lemma 1(i), (iv), we obtain

$$\mu(t; T^*T) = \mu(t; |T|^2) = \mu(t; |T|)^2 = \mu(t; T)^2 = \mu(t; T^*)^2 = \mu(t; |T^*|)^2 = \mu(t; |T^*|^2) = \mu(t; TT^*) \quad (4)$$

for all $t > 0$. Now, for $s > 0$ we have

$$\mu(s; T^*T) = \mu(s; I + X + Y) = 1 + \mu(s; X + Y) > 1 + \mu(s; X) = \mu(s; I + X) = \mu(s; TT^*).$$

This contradicts (4). Therefore, $Y = 0$, and the operator T is normal. $\square$

Passing to adjoint operators, we obtain the following.

Corollary 3. *Let an operator $A \in \mathcal{M}$ be a coisometry, let $B \in S_0(\mathcal{M}, \tau)$, and let $T := A + B$. If $TT^* \geq I$ and T is cohyponormal, then T is normal.*

Lemma 4. *The function $f(t) = t^p$ is operator monotone on the half-line $[0, +\infty)$ for $0 < p \leq 1$.*

Lemma 5 [31]. *If $A \in \mathcal{B}(\mathcal{H})^+$ and $B \in \mathcal{B}(\mathcal{H})_1$, then $Bf(A)B^* \leq f(BAB^*)$ for every function f that is operator monotone on $[0, +\infty)$ and satisfies $f(0) \leq 0$.*

Theorem 4. *Let an operator $A \in \mathcal{B}(\mathcal{H})_1$ be hyponormal. Then $(A^{*n}A^n)^{1/4} \geq (A^*)^{n-2}A^{n-2}$ for all $3 \leq n \in \mathbb{N}$.*

PROOF. Multiplying both sides of the inequality $A^*A \geq AA^*$ on the left by A^* and on the right by A , we obtain

$$A^{*2}A^2 \geq (A^*A)^2. \quad (5)$$

Then $(A^{*2}A^2)^{1/2} \geq A^*A$ by Lemma 4. Multiplying both sides of the latter inequality on the left by A^* and on the right by A , by Lemmas 4, 5, and inequality (5) we get

$$(A^{*3}A^3)^{1/2} \geq A^*(A^{*2}A^2)^{1/2}A \geq A^{*2}A^2 \geq (A^*A)^2,$$

and hence $(A^{*3}A^3)^{1/4} \geq A^*A$. Multiplying both sides of the latter inequality on the left by A^* and on the right by A , by Lemmas 4 and 5 we obtain $(A^{*4}A^4)^{1/4} \geq A^*(A^{*3}A^3)^{1/4}A \geq A^{*2}A^2$; hence, in the same way we derive

$$(A^{*5}A^5)^{1/4} \geq A^*(A^{*4}A^4)^{1/4}A \geq A^{*3}A^3.$$

Continuing this process, we get $(A^{*n}A^n)^{1/4} \geq (A^*)^{n-2}A^{n-2}$, i.e., $|A^n|^{1/2} \geq |A^{n-2}|^2$ for all $3 \leq n \in \mathbb{N}$. $\square$

Corollary 4. *Let an operator $A \in \mathcal{M}_1$ be hyponormal and let $A^n \in S_0(\mathcal{M}, \tau)$ for some $2 \leq n \in \mathbb{N}$. Then A belongs to $S_0(\mathcal{M}, \tau)$ and is normal.*

PROOF. Without loss of generality, we may assume that n is odd (if n is even, consider the operator $A^{n+1} = A \cdot A^n \in S_0(\mathcal{M}, \tau)$). For every $t > 0$, by Lemma 1(i), (ii), (iv), we have

$$\mu(t; A^n)^{1/2} = \mu(t; A^{*n}A^n)^{1/4} = \mu(t; (A^{*n}A^n)^{1/4}) \geq \mu(t; (A^*)^{n-2}A^{n-2}) = \mu(t; |A^{n-2}|^2) = \mu(t; A^{n-2})^2.$$

On the other hand, for every $t > 0$, by Lemma 1(iii),

$$\mu(t; A^n) = \mu(t; A \cdot A^{n-2} \cdot A) \leq \|A\|^2 \mu(t; A^{n-2}) \leq \mu(t; A^{n-2}),$$

and hence $\mu(t; A^{n-2}) \geq \mu(t; A^n) \geq \mu(t; A^{n-2})^4$, $t > 0$. Thus $A^{n-2} \in S_0(\mathcal{M}, \tau)$. Continuing the process of reducing the exponent of A , we obtain $A \in S_0(\mathcal{M}, \tau)$. By [11, Theorem 2.2], the operator A is normal. In particular, $\mu(t; A^k) = \mu(t; A)^k$ for all $t > 0$ and all $k \in \mathbb{N}$ by [11, Theorem 3.1(ii)]. $\square$

Recall that in [16, Theorem 3.3(i)] it was established that if an operator $A \in S(\mathcal{M}, \tau)$ is paranormal and $A^n \in S_0(\mathcal{M}, \tau)$ for some $n \in \mathbb{N}$, then $A \in S_0(\mathcal{M}, \tau)$.

Theorem 5. *Let $P, Q \in \mathcal{B}(\mathcal{H})^{\text{pr}}$, let $n \in \mathbb{N}$, and let $A := (PQ)^n$. Then the following are equivalent:*

- (i) A is hyponormal;
- (ii) $PQ = QP$ (hence $A = P \wedge Q \in \mathcal{B}(\mathcal{H})^{\text{pr}}$).

PROOF. (i) $\Rightarrow$ (ii): We have $(QP)^n(PQ)^n \geq (PQ)^n(QP)^n$, that is,

$$(QPQ)^{2n-1} \geq (PQP)^{2n-1}.$$

Applying Lemma 4 with $p = (2n-1)^{-1} \in (0, 1]$, we obtain $QPQ \geq PQP$. Multiplying both sides of this inequality on the left and on the right by the projection P , we get $PQPQP = (PQP)^2 \geq PQP$. Since $PQP \in \mathcal{B}(\mathcal{H})^+ \cap \mathcal{B}(\mathcal{H})_1$, we obtain $PQP \geq (PQP)^2$. Consequently,

$$PQP = (PQP)^2 \in \mathcal{B}(\mathcal{H})^{\text{pr}}.$$

By von Neumann's theorem [32, Problem 122] (a new proof is given in [33, Theorem 2.2]), the sequence $(PQP)^k$ is nonincreasing and converges in the strong operator topology of $\mathcal{B}(\mathcal{H})$ to the projection $P \wedge Q$. Thus $PQP = P \wedge Q$, and the operator $V := QP$ is a partial isometry. Then $V^* = (QP)^* = PQ$ is also a partial isometry [32, Corollary 2 to Problem 127], and $VV^* = PQP \in \mathcal{B}(\mathcal{H})^{\text{pr}}$. Applying von Neumann's theorem once again, we obtain $QPQ = P \wedge Q$. Hence $PQP = QPQ = P \wedge Q$. Since $\|PQ - P \wedge Q\| = \sqrt{\|PQP - P \wedge Q\|} = 0$ (see [33, p. 6]), we get $PQ = P \wedge Q$ and $A = (PQ)^n = P \wedge Q$. $\square$

Lemma 6. Let $A \in \mathcal{B}(\mathcal{H})^+$, let $P \in \mathcal{B}(\mathcal{H})^{\text{pr}}$, and let $I \geq A \geq P$. Then

- (i) $AP = PA$;
- (ii) if $AP = A$, then $A = P$.

PROOF. (i): Since $P^\perp \geq I - A \geq 0$; according to [34, Chapter 2, 2.17] the operators $P^\perp \in \mathcal{B}(\mathcal{H})^{\text{pr}}$ and $I - A$ commute. Hence $AP = PA$.

(ii): We have $B := A - P \geq 0$ and $A = P + B$. Multiplying both sides of the latter equality on the left by the projection $P^\perp$, we get $BP^\perp = AP^\perp = 0$. Multiplying both sides of the inequality $I \geq P + B$ on the left and on the right by the projection P , we obtain $P \geq P + PBP$. Since $PBP \geq 0$, we have $PBP = |B^{1/2}P|^2 = 0$. Consequently, $B^{1/2}P = 0$ and $BP = B^{1/2} \cdot B^{1/2}P = 0$. Thus $B = BP + BP^\perp = 0 + 0 = 0$, and hence $A = P$. $\square$

Theorem 6. Let $P, Q \in \mathcal{M}^{\text{pr}}$. Then

- (i) $\mu(t; PQ^\perp)^2 = \mu(t; P - PQP) \leq \mu(t; P - Q)$ for all $t > 0$;
- (ii) if $P \sim Q$ and $V \in \mathcal{M}$ with $V^*V = P$ and $VV^* = Q$, then

$$\mu(t; P - V) = \mu(t; Q - V) \leq \mu(t; I - V) \quad \text{for all } t > 0;$$

- (iii) if a partial isometry $U \in \mathcal{B}(\mathcal{H})$ is hyponormal, then

$$\dots \geq U^{*n}U^n \geq \dots \geq U^{*2}U^2 \geq U^*U \geq UU^* \geq U^2U^{*2} \geq \dots \geq U^nU^{*n} \geq \dots$$

PROOF. (i): For every $t > 0$, by Lemma 1(i), (iii), (iv), we have

$$\begin{aligned} \mu(t; PQ^\perp)^2 &= \mu(t; Q^\perp P)^2 = \mu(t; PQ^\perp P) = \mu(t; P - PQP) = \mu(t; P(P - Q)P) \\ &\leq \|P\|^2 \mu(t; P - Q) = \mu(t; P - Q). \end{aligned}$$

(ii): If $P \sim Q$ with $V \in \mathcal{M}$, then V is a partial isometry and $V = VV^*V$ (see [32, Corollary 3 to Problem 127]). Therefore, by Lemma 1(i), (iii), for every $t > 0$ we have

$$\begin{aligned} \mu(t; P - V) &= \mu(t; V^*V - VV^*V) = \mu(t; (V^* - VV^*)V) \\ &\leq \|V\| \mu(t; V^* - VV^*) = \mu(t; V^* - Q) = \mu(t; Q - V). \end{aligned}$$

In the same way, we obtain $\mu(t; Q - V) \leq \mu(t; P - V)$ for every $t > 0$. By Lemma 1(i), (iii), for every $t > 0$ we have

$$\mu(t; Q - V) = \mu(t; VV^* - V) = \mu(t; V(V^* - I)) \leq \|V\| \mu(t; V^* - I) = \mu(t; I - V).$$

(iii): Multiplying both sides of the inequality $U^*U \geq UU^*$ on the left by U and on the right by U^* , and taking into account the equality $U = UU^*U$, we obtain $UU^* \geq U^2U^{*2}$. Multiplying all parts of the chain $U^*U \geq UU^* \geq U^2U^{*2}$ on the left by U and on the right by U^* , and again using $U = UU^*U$, we get $UU^* \geq U^2U^{*2} \geq U^3U^{*3}$. Continuing this process, we obtain

$$U^*U \geq UU^* \geq U^2U^{*2} \geq \dots \geq U^nU^{*n} \geq U^{n+1}U^{*(n+1)} \geq \dots$$

Multiplying both sides of the inequality $U^*U \geq UU^*$ on the left by U^* and on the right by U , and taking into account the equality $U = UU^*U$, we obtain $U^{*2}U^2 \geq U^*U$. Multiplying all parts of the chain $U^{*2}U^2 \geq U^*U \geq UU^*$ on the left by U^* and on the right by U , and again using $U = UU^*U$, we get $U^{*3}U^3 \geq U^{*2}U^2 \geq U^*U$. Continuing this process, we obtain

$$\dots \geq U^{*(n+1)}U^{n+1} \geq U^{*n}U^n \geq \dots \geq U^{*2}U^2 \geq U^*U \geq UU^*. \quad \square$$

Corollary 5. *If a partial isometry $U \in \mathcal{B}(\mathcal{H})$ is hyponormal, then U^n is also a hyponormal partial isometry and $U^{*n}U^n = U^*U$ for every $n \in \mathbb{N}$.*

PROOF. Since $I \geq U^{*n}U^n \geq U^*U \geq UU^* \geq U^nU^{*n} \geq 0$, the operator U^n is hyponormal for every $n \in \mathbb{N}$. Fix $n \in \mathbb{N}$ and put $A := U^{*n}U^n$, $P := U^*U$. Then $I \geq A \geq P$ and

$$AP = U^{*n}U^n \cdot U^*U = U^{*n}U^{n-1} \cdot UU^*U = U^{*n}U^n = A$$

by the equality $UU^*U = U$. Now, by Lemma 6(ii), we obtain $U^{*n}U^n = U^*U = P$ for every $n \in \mathbb{N}$. Consequently, U^n is also a partial isometry; hence, U^{*n} is a partial isometry for every $n \in \mathbb{N}$.

The sequence of projections $(Q_n)_{n=1}^\infty = (U^nU^{*n})_{n=1}^\infty$ is decreasing and, according to Vigier's theorem (see [35, Theorem 4.1.1] or [36, Chapter 1, Theorem 4.5]), it converges in the strong operator topology of $\mathcal{B}(\mathcal{H})$ to some nonnegative operator in $\mathcal{B}(\mathcal{H})_1$. Since the lattice $\mathcal{B}(\mathcal{H})^{\text{pf}}$ is monotone complete, the limit operator is the projection $\bigwedge_{n=1}^\infty Q_n$. $\square$

FUNDING

The research was carried out in the framework of the Development Program of the Scientific Educational Mathematical Center of the Volga Federal District (Agreement 075-02-2025-1725/1).

CONFLICT OF INTEREST

The author of this work declares that he has no conflicts of interest.

References

1. Akhmadiev M., Alhasan H., Bikchentaev A., and Ivanshin P., "Commutators and hyponormal operators on a Hilbert space," J. Iran. Math. Soc., vol. 4, no. 1, 67–78 (2023).
2. Bogdanović K., "A class of norm inequalities for operator monotone functions and hyponormal operators," Complex Anal. Oper. Theory, vol. 18, no. 2, 32, 12 pp. (2024).
3. Fu Y., Cui P., Zu Ch., and Lu Y., "The hyponormal block dual Toeplitz operators," J. Math. Res. Appl., vol. 45, no. 3, 377–394 (2025).
4. Uchiyama A., "Decomposition of hyponormal operator," Nihonkai Math. J., vol. 34, no. 2, 91–102 (2023).
5. Ramesh G. and Sequeira S.S., "Representation and normality of hyponormal operators in the closure of AN -operators," Acta Math. Hungar., vol. 174, no. 2, 341–359 (2024).
6. Bala N. and Ramesh G., "A representation of hyponormal absolutely norm attaining operators," Bull. Sci. Math., vol. 171, 103020, 15 pp. (2021).
7. Gu C., Hendricks J., and Rutherford D., "Hyponormality of block Toeplitz operators," Pacific J. Math., vol. 223, no. 1, 95–111 (2006).
8. Chō M. and Itoh M., "Putnam's inequality for p -hyponormal operators," Proc. Amer. Math. Soc., vol. 123, no. 8, 2435–2440 (1995).
9. Curto R.E., Hwang I.S., and Lee W.Y., "Hyponormality and subnormality of block Toeplitz operators," Adv. Math., vol. 230, no. 4–6, 2094–2151 (2012).
10. Duggal B.P., "On p -hyponormal contractions," Proc. Amer. Math. Soc., vol. 123, no. 1, 81–86 (1995).
11. Bikchentaev A.M., "On normal τ -measurable operators affiliated with semifinite von Neumann algebras," Math. Notes, vol. 96, no. 3, 332–341 (2014).
12. Bikchentaev A.M., "Paranormal measurable operators affiliated with a semifinite von Neumann algebra," Lobachevskii J. Math., vol. 39, no. 6, 731–741 (2018).
13. Bikchentaev A., "Paranormal measurable operators affiliated with a semifinite von Neumann algebra. II," Positivity, vol. 24, no. 5, 1487–1501 (2020).
14. Bikchentaev A.M., "Concerning the theory of τ -measurable operators affiliated to a semifinite von Neumann algebra. II," Matematika i Teor. Komp'yutern. Nauki, vol. 1, no. 2, 3–11 (2023).
15. Bikchentaev A.M., "Concerning the theory of τ -measurable operators affiliated to a semifinite von Neumann algebra. II," Lobachevskii J. Math., vol. 44, no. 10, 4507–4511 (2023).
16. Bikchentaev A., "Hyponormal measurable operators, affiliated to a semifinite von Neumann algebra," Adv. Oper. Theory, vol. 9, no. 4, 83, 17 pp. (2024).
17. Bikchentaev A.M., "Hyponormal measurable operators affiliated to a semifinite von Neumann algebra," Sib. Math. J., vol. 66, no. 3, 656–663 (2025).
18. Bikchentaev A., "Hyponormal measurable operators affiliated to a semifinite von Neumann algebra. III," in: Methusalem Seminars at Ghent Analysis and PDE Center. Extended Abstracts 2023/2024, Basel, Birkhäuser, 2026 (in press).

19. Bikhentaev A.M., “Trace inequalities for measurable operators affiliated to a von Neumann algebra,” *Russian Math. (Iz. VUZ)*, vol. 69, no. 1, 90–95 (2025).
20. Dehimi S. and Mortad M.H., “Unbounded operators having self-adjoint, subnormal, or hyponormal powers,” *Math. Nachr.*, vol. 296, no. 9, 3915–3928 (2023).
21. Muratov M.A. and Chilin V.I., “Topological algebras of measurable and locally measurable operators,” *J. Math. Sci. (N.Y.)*, vol. 239, no. 5, 654–705 (2019).
22. Takesaki M., *Theory of Operator Algebras. I. Operator Algebras and Non-Commutative Geometry V*, Springer, Berlin (2002) (Encyclopaedia of Mathematical Sciences, 124).
23. Dodds P.G., de Pagter B., and Sukochev F.A., *Noncommutative Integration and Operator Theory*, Birkhäuser, Cham (2023) (Progr. Math., 349).
24. Takesaki M., *Theory of Operator Algebras. II. Operator Algebras and Non-Commutative Geometry VI*, Springer, Berlin (2003) (Encyclopaedia of Mathematical Sciences, 125).
25. Fack T. and Kosaki H., “Generalized s -numbers of τ -measurable operators,” *Pacific J. Math.*, vol. 123, no. 2, 269–300 (1986).
26. Gokhberg I.Ts. and Krein M.G., *An Introduction to the Theory of Linear Nonselfadjoint Operators in Hilbert Space*, Amer. Math. Soc., Providence (1969) (Transl. Math. Monogr., 18).
27. Brown L.G. and Kosaki H., “Jensen’s inequality in semifinite von Neumann algebra,” *J. Operator Theory*, vol. 23, no. 1, 3–19 (1990).
28. Kittaneh F., “A note on hyponormal operators,” *Math. Rep. Toyama Univ.*, vol. 9, 105–107 (1986).
29. Chilin V.I., Krygin A.V., and Sukochev Ph.A., “Extreme points of convex fully symmetric sets of measurable operators,” *Integral Equations Operator Theory*, vol. 15, no. 2, 186–226 (1992).
30. Bikhentaev A.M., “The trace and integrable commutators of the measurable operators affiliated to a semifinite von Neumann algebra,” *Sib. Math. J.*, vol. 65, no. 3, 522–533 (2024).
31. Hansen F., “An operator inequality,” *Math. Ann.*, vol. 246, no. 3, 249–250 (1979, 1980).
32. Halmos P.R., *A Hilbert Space Problem Book*. Second edition, Springer, New York and Berlin (1982) (Grad. Texts in Math., 19).
33. Bikhentaev A.M. and Moslehian M.S., “On pairs of projections,” *Positivity*, vol. 29, no. 4, 47, 13 pp. (2025).
34. Strătilă S.V. and Zsidó L., *Lectures on von Neumann Algebras*. Second edition, Cambridge University, Delhi (2019) (Cambridge–IISc Series).
35. Murphy G., *C^* -Algebras and Operator Theory*, Academic, Boston (1990).
36. Sherstnev A.N., *Methods of Bilinear Forms in Noncommutative Measure and Integral Theory*, Fizmatlit, Moscow (2008) [Russian].

Publisher’s Note

Pleiades Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. AI tools may have been used in the translation or editing of this article.

A. M. BIKCHENTAEV
 Kazan (Volga Region) Federal University, Kazan, Russia
<https://orcid.org/0000-0001-5992-3641>
Airat.Bikhentaev@kpfu.ru