

Temperature dynamics of organic matter of the bazhenov suite from electron paramagnetic resonance

Sitdikova L., Khasanova N., Izotov V., Vakhin A., Sidorova E.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2017. All Rights Reserved. Revealing the conditions and mechanisms of evolution of oil deposit based on optical and electron microscopy, electron paramagnetic resonance (EPR) is a necessary step to create improved of recovery methods. Application EPR for study temperature, mechanical and radiation dynamics of organic matters were presented in this communication. The subject of the investigation was the core material from Bazhenov suite (East Siberian oil and gas province). In order to reveal the features of the material composition and the generation potential of the organic matter of rocks, the EPR spectra of the Bazhenov suite samples were investigated. Methods of chloroform extraction, isothermal treatment at a temperature of 350°C and 600°C in hydrogen atmosphere for samples from core section were used. The main hydrocarbon generation zone was established during the degradation of kerogen at 350°C. The inhomogeneity of distribution of carbonate matter is revealed. The main phase of oil generation in the area of 350°C is accompanied by the destruction of kerogen and the formation of a liquid phase of hydrocarbons and leaching of iron. Potential hydrocarbon generation opportunities are associated with destruction of kerogen at 350°C.

<http://dx.doi.org/10.5593/sgem2017/11/S01.071>

Keywords

Bazhenov suite, Electron paramagnetic resonance, Oil deposit, Organic matter

References

- [1] Predtechenskaya E.A., Zlobin O.N. Bazhenov suite as "intermediate" collector in tectonic disorders areas, All-Russian Conference on the Deep Genesis of Oil and Gas "4th Kudryavtsev Readings", Russia, 2015.
- [2] Dorofeeva T.V., Krasnov S.G. The oil collectors of the Bazhenov suite of Western Siberia, Russia, 131 p, 1983.
- [3] Levorsen A.I. The geology of oil and gas, Russia, 693 p, 1970.
- [4] Filina S.I., Korzh M.V., Zonn M.S. Paleogeography and oil content of the novskaya suite of Western Siberia, Russia, 34 p, 1984.
- [5] Sitdikova L.M., Khasanova N.M., Talipov I.P., Izotov V.G., Nizamutdinov N.M., Khasanov R.A., Sidorova E.U., Salimov R.I. Residual organic matter the Jurassic deposits West Siberia, Georesources, Russia, vol. 1(12), 2012, pp 4-7.
- [6] Akramkhodzhaev A.M. Oil and gas - products of the transformation of organic matter, Russia, 281 p, 1982.
- [7] Löhr S.C., Kennedy M.J., Brugger J., Etschmann B. The role of mineral surfaces in the preservation and transformation of organic carbon: meso- to nanoscale spatial complexity of organic matter forms in organic-rich shales, Australian Regolith and Clays Conference, 2012, pp 113-116.

- [8] Bulka G.R., Nizamutdinov N.M., Mukhutdinova N.G., Khasanova N.M., Galeev A.A., Vinokurov V.M. EPR probes in sedimentary rocks: the feature of Mn²⁺ and free radicals distribution in the Permian formation in Tatarstan, *Applied Magnetic Resonance*, vol. 2, 1991, pp 107-115.