

Systemic and local cytokine profile following spinal cord injury in rats: A multiplex analysis

Mukhamedshina Y., Akhmetzyanova E., Martynova E., Khaiboullina S., Galieva L., Rizvanov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Mukhamedshina, Akhmetzyanova, Martynova, Khaiboullina, Galieva and Rizvanov. Our study of the changes in cytokine profile in blood serum and in the spinal cord after traumatic spinal cord injury (SCI) has shown that an inflammatory reaction and immunological response are not limited to the CNS, but widespread. This fact was confirmed by changes detected in a cytokine profile in blood serum samples [MIP-1 α , interleukin 1 (IL-1) α , IL-2, IL-5, IL-1 β , MCP-1, RANTES]. There were also changes in the levels of MIP-1 α , IL-1 α , IL-2, IL-5, IL-18, GM-colon-stimulating factor, IL-17 α , IFN- γ , IL-10, IL-13, MCP-1, and GRO KC CINC-1 in samples of the rat injured spinal cord. The results underscore the complex cytokine network imbalance exhibited after SCI and show significant changes in the concentrations of 14 cytokines/chemokines with different inflammatory and immunological activities.

<http://dx.doi.org/10.3389/fneur.2017.00581>

Keywords

Acute and subacute periods, Cytokine profile, Multiplex analysis, Rats, Spinal cord injury

References

- [1] Laliberte AM, Fehlings MG. The immunological response to spinal cord injury: helpful or harmful? *Exp Neurol* (2013) 247:282-5. doi:10.1016/j.expneurol.2013.01.008
- [2] Scheff SW, Rabchevsky AG, Fugaccia I, Main JA, Lump JJ Jr. Experimental modeling of spinal cord injury: characterization of a force-defined injury device. *J Neurotrauma* (2003) 20(2):179-93. doi:10.1089/08977150360547099
- [3] Ren M, Guo Q, Guo L, Lenz M, Qian F, Koenen RR, et al. Polymerization of MIP-1 chemokine (CCL3 and CCL4) and clearance of MIP-1 by insulin-degrading enzyme. *EMBO J* (2010) 29(23):3952-66. doi:10.1038/emboj.2010.256
- [4] Craig G, Rollins BJ. Chemokines and disease. *Nat Immunol* (2001) 2(2):108. doi:10.1038/84209
- [5] Zhang JM, An J. Cytokines, inflammation and pain. *Int Anesthesiol Clin* (2007) 45(2):27. doi:10.1097/AIA.0b013e318034194e
- [6] Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. Helper T cells and lymphocyte activation. (2002).
- [7] Sprague AH, Khalil RA. Inflammatory cytokines in vascular dysfunction and vascular disease. *Biochem Pharmacol* (2009) 78(6):539-52. doi:10.1016/j.bcp.2009.04.029
- [8] Deo SS, Mistry KJ, Kakade AM, Niphadkar PV. Role played by Th2 type cytokines in IgE mediated allergy and asthma. *Lung India* (2010) 27(2):66-71. doi:10.4103/0970-2113.63609
- [9] Takahashi JL, Giuliani F, Power C, Imai Y, Yong VW. Interleukin-1 β promotes oligodendrocyte death through glutamate excitotoxicity. *Ann Neurol* (2003) 53(5):588-95. doi:10.1002/ana.10519

- [10] John GR, Lee SC, Song X, Riviuccio M, Brosnan CF. IL-1-regulated responses in astrocytes: relevance to injury and recovery. *Glia* (2005) 49(2):161-76. doi:10.1002/glia.20109
- [11] Bose S, Cho J. Role of chemokine CCL2 and its receptor CCR2 in neurodegenerative diseases. *Arch Pharm Res* (2013) 36(9):1039-50. doi:10.1007/s12272-013-0161-z
- [12] Cushing SD, Berliner JA, Valente AJ, Territo MC, Navab M, Parhami F, et al. Minimally modified low density lipoprotein induces monocyte chemotactic protein 1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci U S A* (1990) 87(13):5134-8. doi:10.1073/pnas.87.13.5134
- [13] Standiford TJ, Kunkel SL, Phan SH, Rollins BJ, Strieter RM. Alveolar macrophage-derived cytokines induce monocyte chemoattractant protein-1 expression from human pulmonary type II-like epithelial cells. *J Biol Chem* (1991) 266(15):9912-8.
- [14] Brown Z, Strieter RM, Neild GH, Thompson RC, Kunkel SL, Westwick J. IL-1 receptor antagonist inhibits monocyte chemotactic peptide 1 generation by human mesangial cells. *Kidney Int* (1992) 42(1):95-101. doi:10.1038/ki.1992.266
- [15] Barna BP, Pettay J, Barnett GH, Zhou P, Iwasaki K, Estes ML. Regulation of monocyte chemoattractant protein-1 expression in adult human non-neoplastic astrocytes is sensitive to tumor necrosis factor (TNF) or antibody to the 55-kDa TNF receptor. *J Neuroimmunol* (1994) 50(1):101-7. doi:10.1016/0165-5728(94)90220-8
- [16] Ghirnikar RS, Lee YL, He TR, Eng LF. Chemokine expression in rat stab wound brain injury. *J Neurosci Res* (1996) 46(6):727-33. doi:10.1002/(SICI)1097-4547(19961215)46:6<727::AID-JNR9>3.0.CO;2-H
- [17] Yamagami S, Tamura M, Hayashi M, Endo N, Tanabe H, Katsuura Y, et al. Differential production of MCP-1 and cytokine-induced neutrophil chemoattractant in the ischemic brain after transient focal ischemia in rats. *J Leukoc Biol* (1999) 65(6):744-9.
- [18] Wang X, Yue TL, Barone FC, Feuerstein GZ. Monocyte chemoattractant protein-1 messenger RNA expression in rat ischemic cortex. *Stroke* (1995) 26(4):661-6. doi:10.1161/01.STR.26.4.661
- [19] Arakelyan A, Petrakova J, Hermanova Z, Boyajyan A, Lukl J, Petrek M. Serum levels of the MCP-1 chemokine in patients with ischemic stroke and myocardial infarction. *Mediators Inflamm* (2005) 2005(3):175-9. doi:10.1155/MI.2005.175
- [20] Reynolds MA, Kirchick HJ, Dahlen JR, Anderberg JM, McPherson PH, Nakamura KK, et al. Early biomarkers of stroke. *Clin Chem* (2003) 49(10):1733-9. doi:10.1373/49.10.1733
- [21] Wu X, Wittwer AJ, Carr LS, Crippes BA, DeLarco JE, Lefkowitz JB. Cytokine-induced neutrophil chemoattractant mediates neutrophil influx in immune complex glomerulonephritis in rat. *J Clin Invest* (1994) 94(1):337. doi:10.1172/JCI117326
- [22] Zhang ZJ, Cao DL, Zhang X, Ji RR, Gao YJ. Chemokine contribution to neuropathic pain: respective induction of CXCL1 and CXCR2 in spinal cord astrocytes and neurons. *Pain* (2013) 154(10):2185-97. doi:10.1016/j.pain.2013.07.002
- [23] Vexler ZS, Tang XN, Yenari MA. Inflammation in adult and neonatal stroke. *Clin Neurosci Res* (2006) 6(5):293-313. doi:10.1016/j.cnr.2006.09.008
- [24] Ghirnikar RS, Lee YL, Eng LF. Inflammation in traumatic brain injury: role of cytokines and chemokines. *Neurochem Res* (1998) 23(3):329-40. doi:10.1023/A:1022453332560
- [25] Lumpkins K, Bochicchio GV, Zagol B, Ulloa K, Simard JM, Schaub S, et al. Plasma levels of the beta chemokine regulated upon activation, normal T cell expressed, and secreted (RANTES) correlate with severe brain injury. *J Trauma Acute Care Surg* (2008) 64(2):358-61. doi:10.1097/TA.0b013e318160df9b
- [26] Duque GA, Descoteaux A. Macrophage cytokines: involvement in immunity and infectious diseases. *Front Immunol* (2014) 5:491. doi:10.3389/fimmu.2014.00491
- [27] Yatsiv I, Morganti-Kossmann MC, Perez D, Dinarello CA, Novick D, Rubinstein M, et al. Elevated intracranial IL-18 in humans and mice after traumatic brain injury and evidence of neuroprotective effects of IL-18-binding protein after experimental closed head injury. *J Cereb Blood Flow Metab* (2002) 22(8):971-8. doi:10.1097/00004647-200208000-00008
- [28] Hedtjörn M, Leverin AL, Eriksson K, Blomgren K, Mallard C, Hagberg H. Interleukin-18 involvement in hypoxic-ischemic brain injury. *J Neurosci* (2002) 22(14):5910-9.
- [29] Dickensheets HL, Donnelly RP. IFN-gamma and IL-10 inhibit induction of IL-1 receptor type I and type II gene expression by IL-4 and IL-13 in human monocytes. *J Immunol* (1997) 159(12):6226-33.
- [30] Ueta M, Hamuro J, Ueda E, Katoh N, Yamamoto M, Takeda K, et al. Stat6-independent tissue inflammation occurs selectively on the ocular surface and perioral skin of I β B β -/-mice. *Invest Ophthalmol Vis Sci* (2008) 49(8):3387-94. doi:10.1167/iovs.08-1691
- [31] Mayadas TN, Cullere X, Lowell CA. The multifaceted functions of neutrophils. *Annu Rev Pathol* (2014) 9:181-218. doi:10.1146/annurev-pathol-020712-164023
- [32] Zhu S, Qian Y. IL-17/IL-17 receptor system in autoimmune disease: mechanisms and therapeutic potential. *Clin Sci* (2012) 122(11):487-511. doi:10.1042/CS20110496

- [33] Komiyama Y, Nakae S, Matsuki T, Nambu A, Ishigame H, Kakuta S, et al. IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J Immunol* (2006) 177(1):566-73. doi:10.4049/jimmunol.177.1.566
- [34] Romagnani S. Cytokines and chemoattractants in allergic inflammation. *Mol Immunol* (2002) 38(12):881-5. doi:10.1016/S0161-5890(02)00013-5
- [35] Zhou X, Zöller T, Krieglstein K, Spittau B. TGF β 1 inhibits IFN γ -mediated microglia activation and protects mDA neurons from IFN γ -driven neurotoxicity. *J Neurochem* (2015) 134(1):125-34. doi:10.1111/jnc.13111
- [36] Vojdani A, Lambert J. The role of Th17 in neuroimmune disorders: target for CAM therapy. Part I. *Evid Based Complementary Altern Med* (2011) 2011:984965. doi:10.1093/ecam/nep063
- [37] Lobo-Silva D, Carriche GM, Castro AG, Roque S, Saraiva M. Balancing the immune response in the brain: IL-10 and its regulation. *J Neuroinflammation* (2016) 13(1):297. doi:10.1186/s12974-016-0763-8
- [38] Schmitz T, Chew LJ. Cytokines and myelination in the central nervous system. *Sci World J* (2008) 8:1119-47. doi:10.1100/tsw.2008.140
- [39] Othieno C, Hirsch CS, Hamilton BD, Wilkinson K, Ellner JJ, Toossi Z. Interaction of Mycobacterium tuberculosis-induced transforming growth factor β 1 and interleukin-10. *Infect Immun* (1999) 67(11):5730-5.
- [40] Imitola J, Chitnis T, Khoury SJ. Cytokines in multiple sclerosis: from bench to bedside. *Pharmacol Ther* (2005) 106(2):163-77. doi:10.1016/j.pharmthera.2004.11.007
- [41] Konyuchenko EA, Yu UV, Puchinyan DM, Norkin IA, Drozdova GA. Comparative analysis of neurospecific proteins and cytokines in the acute and early period of traumatic disease in the spinal cord. *Fundam Res* (2014) 7:974-9.
- [42] Odinak MM, Tsygan NV. Mechanisms of damage and regeneration of neurons in spinal cord injury (state of the problem). *Bull Russian Mil Med Acad* (2003) 2:97-103.
- [43] Kothur K, Wienholt L, Brilot F, Dale RC. CSF cytokines/chemokines as biomarkers in neuroinflammatory CNS disorders: a systematic review. *Cytokine* (2016) 77:227-37. doi:10.1016/j.cyto.2015.10.001