

Identification of large rivers of siberia (Ob, Yenisei, Lena) by using GIS technology based on remote sensing of Earth from Cosmos

Shynbergenov Y., Sihanova N.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Decryption of high-resolution space pictures in the territory of an Asian part of Russia is executed. This research is conducted for the purpose of a territory exception within which there is no washout of soils. Methodical questions of application of remote sensing for assessment of inundated sections of the large rivers of Siberia are considered. One of criteria of creation of an electronic vectorial card of river basins is separation of inundated sections. Visual decryption of satellite images for the purposes of detection of inundated sections of the large rivers, can be used in hardly accessible, severe conditions of Siberia, in particular behind a polar circle. As a result of the research conducted on materials of space shooting inundated sections pp are revealed. Ob, Yenisei, Lena. The largest number of inundated sections is created by the river Ob, at the same time the decoded territories are found from a river source in case of confluence of Biya and Katun, to an estuarial section. Rather the most smaller number of floodplains is identified on the Yenisei River. The mid position on the decoded inundated sections is held by the river Lena. Results of a research are planned to be used further when forming the specialized geoinformation database on small river basins of the Arctic basins of an Asian part of Russia. As the model river basin for calculation of potential washout of soils it is planned to use the Lena River catchment.

Keywords

Databases, GIS-analysis, Inundated sections, Remote sensing, Rivers of Siberia, Visual decryption, Washout of the soil

References

- [1] T. V. Vereshchaka, A. T. Zverev S. A. Sladkopevtsev, S. S. Sudakova, Visual decryption methods, Moscow, Nedra, 1990, p. 341.
- [2] I. V. Kozlova, Decryption of aerial photographs in mapping landscapes, Tomsk, UMP, Tomsk, 2006, p. 21- 28.
- [3] V. V. Konakh, A. A. Topaz, F. E., Shalkevich, "Decryption of soils in floodplain landscapes by digital processing of aerial photographs, " in Materials of the II International Conference "Information systems and technologies (IST'2004)", 1st part, Moscow, 2004, p. 180-186.
- [4] V. V. Konakh, A. A. Topaz, F. E. Shalkevich, "Decryption of soils in floodplain landscapes by digital processing of aerial photographs, " in Materials of the II International Conference "Information systems and technologies (IST'2004)", 1st part, Moscow, 2004, p. 180-186

- [5] I. A. Labutina, E. A. Baldina, Workshop on the course "Aerospace Photographs Decryption": Study guide, Moscow, Geographical Department of Moscow State University, 2013, p. 168
- [6] V. F. Usachev, "Monitoring of Russian Rivers, " in Gim International. The Worldwide magazine for Geomatics, XVII (2003), p. 34-37.
- [7] V. N. Sudarikov, O. N. Kalinina, Geological decryption of aerial photographs: methodological guidelines, Orenburg, IPK GOU OSU, 2008, p. 22
- [8] E. A. Shynbergenov, Identification of the floodplains of the major rivers of Siberia (Ob, Yenisei, Lena) according to the satellite-based Earth remote sensing data (ERSD), Treshnikovskii, Readings, 2017
- [9] N. A. Iliina et all. (eds.), Modern Geographic World View and Technology of Geographical Education: Materials of the Scientific-Practical Conference, Ulyanovsk, 2017, p. 304-305
- [10] L. I. Usova, Practical manual on landscape interpretation of aerial photographs of various marshals in Western Siberia, St. Petersburg, Nestor-Istoriia, 2009, p. 80.
- [11] F. V., Recommendations on the use of aerospace information in the study of the channel process, Leningrad, Gidrometeoizdat, 1985, p. 88.