

# Light curve modeling for time-dependent accretion disks in X-ray novae with general relativity effects taken into account

Suleimanov V., Lipunova G., Shakura N.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

We present a method of modeling X-ray and optical light curves of time-dependent accretion  $\alpha$ -disks in X-ray novae. The model is based on an analytic solution for  $\alpha$ -disk evolution after the outburst maximum. The method involves relativistic effects near a Kerr black hole and self-irradiation of an accretion disk. The method is applied to the outbursts of X-ray Nova Monocerotis 1975 (A 0620-00) and X-ray Nova Muscae 1991 (GRS 1124-68). Recently, narrow limits were obtained for the masses and the distances of these binaries ( $11 \pm 2M_{\odot}$ , 1.1 kpc for A 0620-00 and  $7 \pm 0.6M_{\odot}$ , 5.1 kpc for GRS 1124-68, Gelino et al. (2001a,b)). This allows us to limit other model parameters: Kerr parameter (0.5-0.7 and  $\sim 0.95$  respectively for A 0620-00 and GRS 1124-68) and  $\alpha$ -parameter (0.6-0.8 and 0.4-0.5). The inner radius of the accretion disk in GRS 1124-68 has to be few times larger than that of the last marginally stable orbit. Modeled disks in both systems should be thicker than in the standard theory and should thermalize a significant part of the incident X-ray flux ( $\geq 20\%$ ). Possible reasons for these results are discussed.

---

## Keywords

Accretion disks, Binaries, Novae: Individual (Nova Mon 1975; Nova Mus 1991)