

Specific features of reaction of 2-- -benzo[e][1,3,2]dioxaphosphinin-4-ones with perfluorodiacetyl. Synthesis and steric structure of 4',5'- bis(trifluoromethyl)-4-oxo-2-(2,2- 3,3-tetrafluoropropoxy)-2λ 5- spiro[benzo[e][1,3,2]dioxaphosphinine-2,2'-[1,3,2] dioxaphosphole]

Konovalova I., Mironov V., Ivkova G., Zagidullina E., Gubaidullin A., Litvinov I., Kurykin M.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

2-R-benzo[e][1,3,2]dioxaphosphinin-4-ones react with perfluorodiacetyl under mild conditions to form relatively labile spirophosphoranes containing a 1,3,2-dioxaphosphole ring. These compounds gradually convert to more stable 2-R-4,5-bis(trifluoromethyl)-1,3-2λ5-dioxaphosphole 2-oxides and diastereometric 2-R-4-(trifluoroacetyl)-4-(trifluoromethyl)benzo[f][1,3, 2λ5]dioxaphosphepine 2-oxides, whose structure was confirmed by means of NMR and IR spectroscopy. The structure of 4',5'-bis(trifluoromethyl)-4-oxo-2-(2,2,3,3-tetrafluoropropoxy)-2λ 5-spiro[benzo[e][1,3,2]dioxaphosphinine-2,2'-[1,3,2]dioxaphosphole] was confirmed by X-ray diffraction analysis. ©2005 Pleiades Publishing, Inc.

<http://dx.doi.org/10.1007/s11176-005-0270-6>
