



ELSEVIER

Thermochimica Acta 264 (1995) 211–218

thermochimica  
acta

## Thermochemistry of heteroatomic compounds. Part 6<sup>☆</sup>. Enthalpy of solvation and complex formation of some halides of three-coordinated phosphorus compounds in pyridine



V.V. Ovchinnikov<sup>a,\*</sup>, L.I. Lapteva<sup>a</sup>, V.Yu. Kudryavtzev<sup>a</sup>, A.A. Sobanov<sup>b</sup>

<sup>a</sup> Kazan Construction Engineers Institute, Zeligonaya str. 1, Kazan 420043, Tatarstan, Russia

<sup>b</sup> Kazan State University, Lenin str. 18, Kazan 420008, Tatarstan, Russia

Received 8 August 1994; accepted 28 February 1995

### Abstract

Using previously published values of the enthalpies of solvation of some derivatives of three-coordinated phosphorus compounds, the contributions of the non-specific solvation enthalpies in the general solvation effect in benzene, *p*-xylene, methanol and pyridine have been calculated. Taking into account the known possibility of intramolecular complex formation between halogenated derivatives of P<sup>III</sup> and pyridine, the magnitudes of specific interaction enthalpies for  $\text{PCl}_3$ ,  $\text{PBr}_3$ ,  $(\text{EtO})_2\text{PCl}$ ,   $\text{PCl}$  and   $\text{PBr}$  were determined.

Thermodynamic parameters of complex formation between cyclic chlorophosphites and 1-dimethylamino-3,3-dimethylpropan-2-one have been obtained and are discussed.

**Keywords:** Complexation, Enthalpy of solvation; Halide; Phosphorous; Pyridine

### 1. Introduction

As has been shown previously [2], the enthalpies of specific interaction between three-coordinated phosphorus derivatives and some hydrogen-donor solvents, such as chloroform, can serve to some extent as special indicators of the electron density distributions in molecules and the intramolecular interactions of the bonds. This is

\* Corresponding author.

<sup>☆</sup> For Part 5, see Ref. [1].