



Locally most powerful sequential tests of a simple hypothesis vs. one-sided alternatives

Andrey Novikov^{a,*}, Petr Novikov^b

^a Department of Mathematics, Autonomous Metropolitan University - Iztapalapa, San Rafael Atlixco 186, col. Vicentina, C.P. 09340 Mexico City, Mexico

^b Department of Mathematical Statistics, Kazan State University, Kremlevskaya 18, Kazan, Russia

ARTICLE INFO

Article history:

Received 8 May 2009

Received in revised form

30 August 2009

Accepted 7 September 2009

Available online 15 September 2009

MSC:

62L10

62L15

60G40

62C10

Keywords:

Dependent observations

Discrete-time stochastic process

Optimal sequential test

Sequential analysis

Sequential hypothesis testing

Locally most powerful test

ABSTRACT

Let X_1, X_2, \dots be a discrete-time stochastic process with a distribution P_θ , $\theta \in \Theta$, where Θ is an open subset of the real line. We consider the problem of testing a simple hypothesis $H_0: \theta = \theta_0$ vs. a composite alternative $H_1: \theta > \theta_0$, where $\theta_0 \in \Theta$ is some fixed point. The main goal of this article is to characterize the structure of locally most powerful sequential tests in this problem.

For any sequential test (ψ, ϕ) with a (randomized) stopping rule ψ and a (randomized) decision rule ϕ let $\alpha(\psi, \phi)$ be the type I error probability, $\beta_0(\psi, \phi)$ the derivative, at $\theta = \theta_0$, of the power function, and $\mathcal{N}(\psi)$ an average sample number of the test (ψ, ϕ) . Then we are concerned with the problem of maximizing $\beta_0(\psi, \phi)$ in the class of all sequential tests such that

$$\alpha(\psi, \phi) \leq \alpha \quad \text{and} \quad \mathcal{N}(\psi) \leq \mathcal{N},$$

where $\alpha \in [0, 1]$ and $\mathcal{N} \geq 1$ are some restrictions. It is supposed that $\mathcal{N}(\psi)$ is calculated under some fixed (not necessarily coinciding with one of P_θ) distribution of the process X_1, X_2, \dots

The structure of optimal sequential tests is characterized.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

Let $X_1, X_2, \dots, X_n, \dots$ be a discrete-time stochastic process with a distribution P_θ , $\theta \in \Theta$, where Θ is an open subset of the real line. We consider the problem of testing a simple hypothesis $H_0: \theta = \theta_0$ vs. a composite alternative $H_1: \theta > \theta_0$, where $\theta_0 \in \Theta$ is some fixed point. The main goal of this article is to characterize the structure of locally most powerful, in the sense of Berk (1975), sequential tests in this problem.

We follow Novikov (2009b) in the definitions and notation related to sequential hypothesis tests, as well as their interpretation and characteristics (see also Wald, 1950; Ferguson, 1967; DeGroot, 1970; Schmitz, 1993; Ghosh et al., 1997, among many others).

In particular, we say that a pair (ψ, ϕ) is a sequential hypothesis test if

$$\psi = (\psi_1, \psi_2, \dots, \psi_n, \dots) \quad \text{and} \quad \phi = (\phi_1, \phi_2, \dots, \phi_n, \dots),$$

* Corresponding author.

E-mail addresses: an@xanum.uam.mx (A. Novikov), pnovi@mail.ru (P. Novikov).

URL: <http://mat.izt.uam.mx/profs/anovikov/en> (A. Novikov).