

Electrochemical properties and reactivity of organonickel sigma-complex [NiBr(Mes)(bpy)] (Mes = 2,4,6-trimethylphenyl, bpy = 2,2'-bipyridine)

Sakhapov I., Gafurov Z., Babaev V., Kurmaz V., Mukhametbareev R., Rizvanov I., Sinyashin O., Yakhvarov D.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2015, Pleiades Publishing, Ltd. Electrochemical properties and reactivity of electrochemically activated forms of organonickel sigma-complex [NiBr(Mes)(bpy)] (where Mes = 2,4,6-trimethylphenyl, bpy = 2,2'-bipyridine) were studied. The activation of the organonickel sigma-complex was found to proceed under both electrochemical reduction and oxidation conditions to give coordinatively unsaturated forms of the complex: radical [Ni(Mes)(bpy)]• and cationic complex [Ni(Mes)(bpy)]+, respectively. It was shown experimentally that the active forms of organonickel complex [NiBr(Mes)(bpy)] can react with organic substrates (cyclohexene, octene-1, tetrahydrofuran) and convert nitriles (acetonitrile, acetonitrile-d₃, chloroacetonitrile) into corresponding imines containing 2,4,6-trimethylphenyl fragment.

<http://dx.doi.org/10.1134/S1023193515110142>

Keywords

2,2'-bipyridine, cyclic voltammetry, mass spectrometry, organonickel sigma-complexes, oxidation, reduction