

Kinetics of CO₂-hydrate formation from ice powders: Data summary and modeling extended to low temperatures

Falenty A., Salamatin A., Kuhs W.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

The shrinking-core model of the formation of gas hydrates from ice spheres with a well-defined geometry gives experimental access to the gas permeation in bulk hydrates. Here we report on results obtained for CO₂ clathration experiments in the temperature range from 185 to 272 K, extending earlier work to much lower temperature conditions. The activation energy deduced from the permeation coefficients changes its value from ~46 kJ/mol at higher temperatures to ~19 kJ/mol below 225 K. We compare our results with published molecular dynamics simulation as well as nuclear magnetic resonance studies and provide arguments that the rate limiting process at lower temperatures is the cage-to-cage jumping of CO₂ molecules via a "hole-in-the-cage" mechanism involving extrinsic water vacancies in cage walls. The rate-limiting process at higher temperatures can be explained by the temperature-dependent creation of intrinsic water-vacancy-interstitial pairs. The results obtained for CO₂-hydrate are compared to earlier results for CH₄-hydrate formation. The permeation of CO₂ molecules through bulk hydrate is found to be about three times faster when compared to the CH₄ case. This explains the faster clathration reaction of CO₂-hydrate in comparison to CH₄-hydrate. © 2013 American Chemical Society.

<http://dx.doi.org/10.1021/jp310972b>
