

# Nuclear magnetic relaxation induced by the relaxation of electron spins

Borich M., Bunkov Y., Kurkin M., Tankeyev A.  
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017, Pleiades Publishing, Inc. A physical mechanism responsible for the relaxation of nuclear spins coupled by the hyperfine interaction to relaxed electron spins in materials with spin ordering is proposed. The rate of such induced nuclear spin relaxation is proportional to the dynamic shift of the nuclear magnetic resonance (NMR) frequency. Therefore, its maximum effect on the NMR signal should be expected in the case of nuclear spin waves existing in the system. Our estimates demonstrate that the induced relaxation can be much more efficient than that occurring due to the Bloch mechanism. Moreover, there is a qualitative difference between the induced and Bloch relaxations. The dynamics of nuclear spin sublattices under conditions of the induced relaxation is reduced to the rotation of  $m_1$  and  $m_2$  vectors without any changes in their lengths ( $m_{12}(t) = m_{22}(t) = m_{02}(t) = \text{const}$ ). This means that the excitation of NMR signals by the resonant magnetic field does not change the temperature  $T_n$  of the nuclear spin system. This is a manifestation of the qualitative difference between the induced and Bloch relaxations. Indeed, for the latter, the increase in  $T_n$  accompanying the saturation of NMR signals is the dominant effect.

<http://dx.doi.org/10.1134/S002136401701009X>

---

## References

- [1] A. S. Borovik-Romanov, Yu. M. Bunkov, V. V. Dmitriev, and Yu. M. Mukharskiy, JETP Lett. 40, 1033 (1984).
- [2] Yu. M. Bunkov, S. N. Fisher, A. M. Guenault, and G. R. Pickett, Phys. Rev. Lett. 69, 3092 (1992).
- [3] Yu. M. Bunkov and G. E. Volovik, Phys. Rev. Lett. 98, 265302 (2007).
- [4] S. Autti, Yu. M. Bunkov, V. B. Eltsov, P. J. Heikkinen, J. J. Hosio, P. Hunger, M. Krusius, and G. E. Volovik, Phys. Rev. Lett. 108, 145303 (2012).
- [5] A. S. Borovik-Romanov, Yu. M. Bun'kov, A. de Vaard, V. V. Dmitriev, V. Makrotsieva, Yu. M. Mukharskii, and D. A. Sergatskov, JETP Lett. 47, 478 (1988).
- [6] G. E. Volovik, J. Low Temp. Phys. 153, 266 (2008).
- [7] Y. M. Bunkov and G. Volovik, J. Phys.: Condens. Matter 22, 164210 (2010).
- [8] A. S. Borovik-Romanov, Yu. M. Bunkov, V. V. Dmitriev, and Yu. M. Mukharskiy, JETP Lett. 39, 469 (1984).
- [9] T. Sato, T. Kunimatsu, K. Izumina, A. Matsubara, M. Kubota, T. Mizusaki, and Yu. M. Bunkov, Phys. Rev. Lett. 101, 055301 (2008).
- [10] P. Hunger, Yu. M. Bunkov, E. Collin, and H. Godfrin, J. Low Temp. Phys. 158, 129 (2010).
- [11] Yu. M. Bunkov and G. E. Volovik, JETP Lett. 89, 306 (2009).
- [12] Yu. M. Bunkov and G. E. Volovik, in Novel Superfluids, Ed. by K. H. Bennemann and J. B. Ketterson (Oxford Univ. Press, Oxford, 2013), Chap.4.
- [13] Yu. M. Bunkov, E. M. Alakshin, R. R. Gazizulin, A. V. Klochkov, V. V. Kuzmin, V. S. L'vov, and M. S. Tagirov, Phys. Rev. Lett. 108, 177002 (2012).

- [14] E. A. Turov and M. P. Petrov, Nuclear Magnetic Resonance in Ferro-and Antiferromagnets (Nauka, Moscow, 1969; Wiley, Chichester, 1972).
- [15] M. I. Kurkin and E. A. Turov, Nuclear Magnetic Resonance in Magnetically Ordered Materials and Its Application (Nauka, Moscow, 1990) [in Russian].
- [16] A. S. Borovik-Romanov, Yu. M. Bun'kov, B. S. Dumesh, M. I. Kurkin, M. P. Petrov, and V. P. Chekmarev, Sov. Phys. Usp. 27, 235 (1984).
- [17] A. S. Borovik-Romanov, in Achievements of Science and Engineering, Vol. 4: Antiferromagnetism and Ferrites (Akad. Nauk SSSR, Moscow, 1962), p. 7 [in Russian].
- [18] E. A. Turov, M. I. Kurkin, and V. V. Nikolaev, Sov. Phys. JETP 37, 147 (1973).
- [19] L. D. Landau and E. M. Lifshits, Phys. Zs. Sow. 8, 153 (1935).
- [20] T. L. Gilbert, Phys. Rev. 100, 1243 (1955).