

# Advanced modeling of a moderate-resolution holographic spectrograph

Muslimov E., Valyavin G., Fabrika S., Pavlycheva N.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2017 Optical Society of America. In the present article we consider an accurate modeling of a spectrograph with a cascade of volume-phase holographic gratings. The proposed optical scheme allows us to detect spectra in an extended wavelength range without gaps, providing relatively high spectral resolution and high throughput. However, modeling and minimization of possible cross-talk between gratings and stray light in such a scheme represents a separate task. We use analytical equations of the coupled-wave theory together with rigorous coupled-wave analysis to optimize the gratings parameters and further apply the latter together with a non-sequential ray-tracing algorithm to model propagation of beams through the spectrograph. The results show relatively high throughput up to 53% and the absence of any significant cross-talk or ghost images, even for ordinary holograms recorded on dichromated gelatin.

<http://dx.doi.org/10.1364/AO.56.004284>

---

## References

- [1] H. J. Caulfield, *Handbook of Optical Holography* (Academic, 1979).
- [2] S. C. Barden, J. A. Arns, W. S. Colburn, and J. B. Williams, "Volume-phase holographic gratings and the efficiency of three simple volume-phase holographic gratings," *Publ. Astron. Soc. Pac.* 112, 809-820 (2000).
- [3] P.-A. Blanche, P. Gailly, S. L. Habraken, P. Lemaire, and C. Jamar, "Volume phase holographic gratings: large size and high diffraction efficiency," *Opt. Eng.* 43, 2603-2612 (2004).
- [4] E. Oliva, S. Todd, M. Cirasuolo, H. Schnetler, D. Lunney, P. Rees, A. Bianco, E. Diolaiti, D. Ferruzzi, M. Fisher, I. Guinouard, M. Iuzzolino, I. Parry, X. Sun, A. Tozzi, and F. Vitali, "Updated optical design and trade-off study for MOONS, the multi-object optical and near infrared spectrometer for the VLT," *Proc. SPIE* 9147, 91472C (2014).
- [5] H. Sugai, N. Tamura, H. Karoji, A. Shimono, N. Takato, M. Kimura, Y. Ohyama, A. Ueda, H. Aghazarian, M. V. de Arruda, R. H. Barkhouser, C. L. Bennett, S. Bickerton, A. Bozier, D. F. Braun, K. Bui, C. M. Capocasale, M. A. Carr, B. Castilho, Y.-C. Chang, H.-Y. Chen, R. C. Y. Chou, O. R. Dawson, R. G. Dekany, E. M. Ek, R. S. Ellis, R. J. English, D. Ferrand, D. Ferreira, C. D. Fisher, M. Golebiowski, J. E. Gunn, M. Hart, T. M. Heckman, P. T. P. Ho, S. Hope, L. E. Hovland, S.-F. Hsu, Y.-S. Hu, P. J. Huang, M. Jaquet, J. E. Karr, J. G. Kempenaar, M. E. King, O. Le Fèvre, D. Le Mignant, H.-H. Ling, C. Loomis, R. H. Lupton, F. Madec, P. Mao, L. S. Marrara, B. Ménard, C. Morantz, H. Murayama, G. J. Murray, A. C. de Oliveira, C. M. de Oliveira, L. S. de Oliveira, J. D. Orndorff, R. de Paiva Vilaça, E. J. Partos, S. Pascal, T. Pegot-Ogier, D. J. Reiley, R. Riddle, L. Santos, J. B. dos Santos, M. A. Schwochert, M. D. Seiffert, S. A. Smee, R. M. Smith, R. E. Steinkraus, L. Sodr e, Jr., D. N. Spergel, C. Surace, L. Tresse, C. Vidal, S. Vives, S.-Y. Wang, C.-Y. Wen, A. C. Wu, R. Wyse, and C.-H. Yan, "Prime focus spectrograph for the Subaru telescope: massively multiplexed optical and near-infrared fiber spectrograph," *J. Astron. Telesc. Instrum. Syst.* 1, 035001 (2015).
- [6] E. Muslimov, N. K. Pavlycheva, G. G. Valyavin, and S. N. Fabrika, "Moderate-resolution holographic spectrograph," *Astrophys. Bull.* 71, 357-365 (2016).
- [7] E. Muslimov, G. G. Valyavin, S. N. Fabrika, and N. K. Pavlycheva, "Design and modeling of a moderate-resolution astronomic spectrograph with volume-phase holographic gratings," *Proc. SPIE* 9908, 990842 (2016).

- [8] H. Kogelnik, "Coupled wave theory for thick hologram gratings," *Bell Syst. Tech. J.* 48, 2909-2947 (1969).
- [9] N. Chateau and J.-P. Hugonin, "Algorithm for the rigorous coupled-wave analysis of grating diffraction," *J. Opt. Soc. Am. A* 11, 1321-1331 (1994).
- [10] P. Lalanne and J.-P. Hugonin, "High-order effective-medium theory of subwavelength gratings in classical mounting: application to volume holograms," *J. Opt. Soc. Am. A* 15, 1843-1851 (1998).
- [11] E. Muslimov, "Transmission holographic grating with improved diffraction efficiency for a flat-field spectrograph," *Proc. SPIE* 8787, 87870B (2013).
- [12] T. Loukina, S. Massenot, R. Chevallier, K. Heggarty, N. M. Shigapova, and A. F. Skochilov, "Volume diffraction gratings for optical telecommunications applications: design study for a spectral equalizer," *Opt. Eng.* 43, 2658-2665 (2004).
- [13] S. K. Case and R. Alferness, "Index modulation and spatial harmonic generation in dichromated gelatin films," *Appl. Phys.* 10, 41-51 (1976).
- [14] R. Pawluczyk, "Modified Brewster angle technique for the measurement of the refractive index of a DCG layer," *Appl. Opt.* 29, 589-592 (1990).
- [15] M. J. Weber, *Handbook of Optical Materials* (CRC Press, 2002).
- [16] M. Mayor, F. Pepe, D. Queloz, F. Bouchy, G. Rupprecht, G. Lo Curto, G. Avila, W. Benz, J.-L. Bertaux, X. Bonfils, Th. Dall, H. Dekker, B. Delabre, W. Eckert, M. Fleury, A. Gilliotte, D. Gojak, J. C. Guzman, D. Kohler, J.-L. Lizon, A. Longinotti, C. Lovis, D. Megevand, L. Pasquini, J. Reyes, J.-P. Sivan, D. Sosnowska, R. Soto, S. Udry, A. van Kesteren, L. Weber, and U. Weilenmann, "Setting R. New Standards with HARPS," *Messenger* 114, 20-24 (2003).
- [17] K. G. Strassmeier, I. Ilyin, A. Järvinen, M. Weber, M. Woche, S. I. Barnes, S.-M. Bauer, E. Beckert, W. Bittner, R. Bredthauer, T. A. Carroll, C. Denker, F. Dionies, I. DiVarano, D. Döscher, T. Fechner, D. Feuerstein, T. Granzer, T. Hahn, G. Harnisch, A. Hofmann, M. Lesser, J. Paschke, S. Pankratow, V. Plank, D. Plüschke, E. Popow, D. Sablowski, and J. Storm, "PEPSI: the high-resolution echelle spectrograph and polarimeter for the large binocular telescope," *Astron. Nachr.* 336, 324-361 (2015).
- [18] N. Kashikawa, K. Aoki, R. Asai, N. Ebizuka, M. Inata, M. Iye, K. S. Kawabata, G. Kosugi, Y. Ohyama, K. Okita, T. Ozawa, Y. Saito, T. Sasaki, K. Sekiguchi, Y. Shimizu, H. Taguchi, T. Takata, Y. Yadoumaru, and M. Yoshida, "FOCAS: the faint object camera and spectrograph for the Subaru telescope," *Publ. Astron. Soc. Jpn.* 54, 819-832 (2002).
- [19] O. Sholukhova, D. Bizyaev, S. Fabrika, A. Sarkisyan, V. Malanushenko, and A. Valeev, "New luminous blue variables in the Andromeda galaxy," *Mon. Not. R. Astron. Soc.* 447, 2459-2467 (2015).
- [20] S. Fabrika, Y. Ueda, A. Vinokurov, O. Sholukhova, and M. Shidatsu, "Supercritical accretion disks in ultraluminous X-ray sources and SS 433," *Nat. Phys.* 11, 551-553 (2015).
- [21] J. Poutanen, S. Fabrika, A. F. Valeev, O. Sholukhova, and J. Greiner, "On the association of the ultraluminous X-ray sources in the Antennae galaxies with young stellar clusters," *Mon. Not. R. Astron. Soc.* 432, 506-519 (2013).
- [22] G. G. Valyavin, A. O. Grauzhanina, G. A. Galazutdinov, D. R. Gadelshin, R. Y. Zhuchkov, V. G. Orlov, T. E. Burlakova, A. F. Valeev, A. F. Kholtygin, A. K. Rzaev, and D. E. Mkrtichian, "Search for signatures of reflected light from the exoplanet HD 189733b by the method of residual dynamical spectra," *Astrophys. Bull.* 70, 466-473 (2015).
- [23] G. Valyavin, D. Shulyak, G. A. Wade, K. Antonyuk, S. V. Zharikov, G. A. Galazutdinov, S. Plachinda, S. Bagnulo, L. Fox Machado, M. Alvarez, D. M. Clark, J. M. Lopez, D. Hiriart, I. Han, Y.-B. Jeon, C. Zurita, R. Mujica, T. Burlakova, T. Szeifert, and A. Burenkov, "Suppression of cooling by strong magnetic fields in white dwarf stars," *Nature* 515, 88-91 (2014).
- [24] J. D. Landstreet, S. Bagnulo, A. Martin, and G. Valyavin, "Discovery of an extremely weak magnetic field in the white dwarf LTT 16093 = WD 2047+372," *Astron. Astrophys.* 591, A80 (2016).
- [25] G. Valyavin, G. A. Wade, S. Bagnulo, T. Szeifert, J. D. Landstreet, I. Han, and A. Burenkov, "The peculiar magnetic field morphology of the white dwarf WD 1953-011: evidence for a large-scale magnetic flux tube?" *Astrophys. J.* 683, 466-478 (2008).