

The role of ATP-dependent potassium channels and nitric oxide system in the neuroprotective effect of preconditioning

Deryagin O., Gavrilova S., Buravkov S., Andrianov V., Yafarova G., Gainutdinov K., Koshelev V.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Media Sphera. All rights reserved. Objective. To study a role of ATP-dependent potassium channels (K⁺ATP) in the neuroprotective effect of ischemic (IP) and pharmacological (PP) preconditioning and evaluate the dynamics of blood nitric oxide (NO) metabolites in cerebral ischemia. Material and methods. A model of ischemic stroke induced by the electrocoagulation of a middle cerebral artery (MCA) branch was used in male rats (n=86). Glibenclamide, a selective inhibitor of ATP-sensitive K⁺ channels, and diazoxide, a potassium channel activator, were used. IP and PP were performed 24 h before MCA occlusion. Blood concentrations of NO, NO₃- and NO₂- were measured 5, 24 and 72 h after occlusion. Results. IP decreased a lesion area by 37% (p<0/05) and the preliminary introduction of glibenclamide levelled the effect of IP. A protective effect of PP was similar to that of IP. A decrease in oxygenated R-conformers of Hb-NO and a reverse increase in non-oxygenated T-conformers as well as NO₃- and NO₂- were noted 5h after MCA occlusion. In the first 24 h after MCA occlusion, contents of NO₃- and NO₂- returned to normal values. There were changes in the concentrations of Hb-NO complexes as well, with the predominance of R-conformers and minimal contents of T-conformers. Moreover, the correlations between K⁺ATP channel blockade and the decrease in serum NO₃- and NO₂ were found (p<0/03). Conclusion. The neuroprotective effect of preconditioning is caused by the activation of K⁺ATP channels. An analysis of NO metabolite concentrations in the blood of rats with IP suggests that Hb-NO complexes belonging to R-conformers deposit and carry NO in tissues releasing NO accumulated via R→T transfer in conditions of ischemia.

<http://dx.doi.org/10.17116/jnevro20161168217-23>

Keywords

ATP-dependent potassium channels, Cerebral ischemia, Electron paramagnetic resonance, Hemoglobin, Ischemic tolerance, Neuroprotection, Nitric oxide, Phenomenon of preconditioning, Spectrophotometry