

Complexes of 1,8-bis(hydrazidomethylsulfonyl)octane and 1,10-bis(hydrazidomethylsulfonyl)decane with copper(II)

Neklyudov V.V., Boos G.A., Shulaeva M.M., Lodochnikova O.A., Chmutowa G.A., Bulavina E.P., Amirov R.R.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2020 Kazan Federal University. All rights reserved. 1,8-bis(hydrazidomethylsulfonyl)octane and 1,10-bis(hydrazidomethylsulfonyl)decane differ by hydrocarbon spacer length, solubility in water, and polymerization tendencies. They and their amides are known to be new antitubercular agents. The substances act as polydentate ligands; in the pH range of 4.5–5.0 and with copper(II), they form cationic mono- and binuclear complexes. Preparatively isolated copper complexes with each of the two compounds have different compositions, which also differ from the composition of complexes in the solution.

<http://dx.doi.org/10.26907/2542-064X.2020.1.33-51>

Keywords

1,10-bis(hydrazidomethylsulfonyl)decane, 1,8-bis(hydrazidomethylsulfonyl)octane, Acid-base properties, Complexation, Copper(II), Elemental analysis, Ligand denticity, Three-dimensional structure, X-ray crystallography

References

- [1] Fattakhov S.G., Shulaeva M.M., Kravchenko M.A., Mingaleev D.N., Skornyakov S.N., Sinyashin O.G. α,ω -Bis(amide- and hydrazide methyl sulfinyl- and sulphonyl)alkanes having antituberculotic activity and α,ω -Bis(methoxy carbonyl methyl sulfinyl- or sulphonyl)alkanes for their production. Patent RF no. 2015101008/04, 2016. (In Russian)
- [2] Neklyudov V.V., Boos G.A., Shulaeva M.M., Chmutova G.A., Bagina Y.I., Salnikov Yu.I., Amirov R.R. Solution state and complexing ability of 1,4-bis(amidomethylsulfinyl)butane toward iron(III), copper(II), cobalt(II), nickel(II), and manganese(II). Russ. Chem. Bull., 2017, vol. 66, no. 4, pp. 628–635. doi: 10.1007/s11172-01-1783-y.
- [3] Neklyudov V.V., Boos G.A., Chmutova G.A., Shulaeva M.M., Bagina Y.I., Amirov R.R. Complex formation of 1,5-bis(amidomethylsulfonyl)pentane with copper(II) and iron(III). Russ. J. Gen. Chem., 2018, vol. 88, no. 8, pp. 1672–1680. doi: 10.1134/S1070363218080194.
- [4] Schönfeldt N. Grenzflächenaktive Äthylenoxid-Addukte. Stuttgart, Wiss. Verlagsges. mbH, 1976. 40 Abb., 137 Tab., XXIV, 1288 S. (In German)
- [5] Gordon A.J., Ford R.A. The Chemist's Companion. A Handbook of Practical Data, Techniques, and References. New York, Wiley, 1973. 560 p.
- [6] Aleksandrov V.V. Kislotnost' nevodnykh rastvorov [Acidity of Nonaqueous Solutions]. Kharkov, Vish. Shk., 1981. 152 p. (In Russian)

- [7] Salnikov Yu.I., Glebov A.N., Devyatov F.V. Poliyadernye kompleksy v rastvorakh [Polynuclear Complexes in Solutions]. Kazan, Izd. Kazan. Univ., 1989. 287 p. (In Russian)
- [8] Vasil'ev V.P. Termodinamicheskie svoistva rastvorov elektrolitov [Thermodynamic Properties of Electrolyte Solutions]. Moscow, Vyssh. Shkola, 1982. 320 p. (In Russian)
- [9] Shcherbakova E.S., Gol'dshtein I.P., Gur'yanova E.N., Kocheshkov K.A. Method of treatment of the results of a physicochemical investigation of complex compounds in solutions on an electronic computer. Bull. Acad. Sci. USSR, Div. Chem. Sci., 1975, vol. 24, no. 6, pp. 1165–1172. (In Russian)
- [10] Hartley F.R., Burgess C., Alcock R.M. Solution in Equilibria. New York, Ellis Horwood, 1980. 361 p.
- [11] Afanas'eva G.V. Complex formation and chemical exchange in water and water-organic solutions of copper(II), nickel(II), and cobalt(II) with hydrazides of some acids. Diss. Cand. Chem. Sci. Kazan, 2008. 240 p. (In Russian)
- [12] Přibil R. Komplexony v chemické analyse. Praha, Nakladatelství Československé akad., 1957. 473 s. (In Czech)
- [13] Sheldrick G.M. SHELXT – integrated space-group and crystal-structure determination. Acta Crystallogr., Sect. A: Found. Adv., 2015, vol. 71, pt. 5, pp. 3–8. doi: 10.1107/S2053273314026370.
- [14] Sheldrick G.M. Crystal structure refinement with SHELXL. Acta Crystallogr., Sect. C: Struct. Chem., 2015, vol. 71, pt. 1, pp. 3–8. doi: 10.1107/S2053229614024218.
- [15] Farrugia L.J. WinGX suite for small-molecule single-crystal crystallography. J. Appl. Crystallogr., 1999, vol. 32, pt. 4, pp. 837–838. doi: 10.1107/S0021889899006020.
- [16] APEX2 (Version 2.1), SAINTPlus. Data Reduction and Correction Program (Version 7.31A, Bruker Advanced X-ray Solutions. Madison, Wis., BrukerAXS Inc., 2006.
- [17] Spektroskopicheskie metody v khimii kompleksnykh soedinenii [Spectral Methods in the Chemistry of Complex Compounds]. Vdovenko V.M. (Ed.). Moscow, Khimiya, 1964. 268 p. (In Russian)
- [18] Neklyudov V.V., Voronina Yu.K., Boos G.A., Shulaeva M.M., Fattakhov S.G., Salnikov Yu.I. Pirafeneband melfene drugs and their components in the reaction with copper(II) chloride in a chloride medium. Uchenye Zapiski Kazanskogo Universiteta. Seriya Estestvennye Nauki, 2012, vol. 154, no. 4, pp. 105–111. (In Russian)
- [19] Lever A.B.P. Inorganic Electronic Spectroscopy (Studies in Physical and Theoretical Chemistry). Elsevier Sci., 1986. 864 p. (In Russian)
- [20] Salnikov Yu.I., Boos G.A., Gibadullina Ch.V. Solvation of copper(II) and ethylenediamine tetraacetic acid in some water-organic solvent media. Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol., 1991, vol. 17, no. 5, pp. 20–24. (In Russian)
- [21] Rossotti F.J.C., Rossotti H. The Determination of Stability Constants and Other Equilibrium Constants in Solution. New York, Toronto, London, McGraw-Hill Book Co., 1961. 425 p.
- [22] Umland F., Janssen A., Thierig D., Wünsch G. Theorie und praktische Anwendung von Komplexbildnern. Frankfurt am Main, Frankfurt Akad. Verlagsges., 1971. 759 S. (In German)
- [23] Tsitsishvili A.D., Tsutsunava T.I., Gogorishvili P.V., Kharitonov Yu.Ya. Complexes of cobalt(II), nickel(II), and copper(II) with dihydrazide of aselaic acid. Issled. Obl. Khim. Kompleksn. Prostykh Soedin. Nek. Perekhodnykh Redk. Met., 1978, no. 3, pp. 21–23. (In Russian)
- [24] Kiselev Yu.M. Stabilization of oxidation states in transition metals. Russ. J. Inorg. Chem., 2007, vol. 52, no. 11, pp. 1717–1725. doi: 10.1134/S0036023607110149.