

# Males and ephippial females of oriental chydoridae (Cladocera: Anomopoda) from Cat Tien National Park, South Vietnam

Sinev A.Y., Semenyuk I.I.  
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

Cladocera (Crustacea: Branchiopoda) of South-East Asia have been intensively investigated over the last decades, but males and ephippial females of many taxa remain unknown or poorly studied. Males and ephippial females of ten species of Chydoridae were collected by monthly sampling of eight water bodies during autumn-winter dry season in Cat Tien National Park, South Vietnam. Morphology of studied specimens is described and taxonomic status of studied species is discussed. Gamogenetic specimens were sparse in all studied taxa, contributing less than 8 % of the populations in samples. Gamogenetic specimens were found mostly during December-January, what suggests that daylight period is the main factor inducing a sexual reproduction in most studied species.

<http://dx.doi.org/10.11646/zootaxa.4941.3.4>

---

## Keywords

Chydoridae, Cladocera, Ehipial female, Gamogenesis, Male, Morphology, Taxonomy

## References

- [1] Alonso, M. (1987) Morphological differentiation of two new *Ephemeropterus* species (Cladocera, Chydoridae) belonging to the barroisi complex: *E. margalefi* and *E. epiphantooii*, in Spain. *Hydrobiologia*, 145, 131-146. [https://doi.org/10.1007/978-94-009-4039-0\\_15](https://doi.org/10.1007/978-94-009-4039-0_15)
- [2] Alonso, M. (1996) Crustacea Branchiopoda. In: Ramos, M. (Ed.), *Fauna Ibérica*. Vol. 7. Museo Nacional de Ciencias Naturales, CSIC, Madrid, pp. 1-486.
- [3] Alonso M. & Pretus J.L. (1989) *Alona iberica*, new species: first evidence of noncosmopolitanism within the *A. karua* complex (Cladocera: Chydoridae). *Journal of Crustacean Biology*, 9, 453-476. <https://doi.org/10.1163/193724089X00449>
- [4] Belyaeva, M. & Taylor, D.J. (2009) Cryptic species within the *Chydorus sphaericus* species complex (Crustacea: Cladocera) revealed by molecular markers and sexual stage morphology. *Molecular Phylogenetics and Evolution*, 50, 534-546. <https://doi.org/10.1016/j.ympev.2008.11.007>
- [5] Brooks, J.L. (1957) The systematics of North American *Daphnia*. *Memories of Connecticut Academy of Arts and Science*, 13, 1-180.
- [6] Chiang, S.C. & Du, N.S. (1979) *Fauna Sinica: Crustacea: Freshwater Cladocera*. Science Press, Beijing, 297 pp. [in Chinese]
- [7] Dumont, H.J. & Negrea, S.V. (2002) Introduction to the class Branchiopoda. In: Dumont H.J. (Ed.), *Guides to the identification of the microinvertebrates of the continental waters of the world*. Vol. 19. Backhuys Publishers, Leiden, 398 pp.

- [8] Dumont, H.J. & Van de Velde, I. (1977) Cladocères et Conchostracés récoltes par le professeur Th. Monod dans la moyenne vallée du Niger en décembre 1972 et janvier 1973. Bulletin de l'Institut Français d'Afrique Noire, Serie A, Sciences Naturelles, 39, 75–93.
- [9] Frey, D.G. (1982) The reticulated species of Chydorus (Cladocera, Chydoridae): two new species with suggestions of convergence. *Hydrobiologia*, 93, 255–279. <https://doi.org/10.1007/BF00012335>
- [10] Frey, D.G. (1991) A new genus of alonine chydorid cladocerans from athalassic saline waters of New South Wales, Australia. *Hydrobiologia*, 224, 11–48. <https://doi.org/10.1007/BF00006361>
- [11] Kim, K., Kotov, A.A. & Taylor, D.J. (2006) Hormonal induction of undescribed males resolves cryptic species of cladocerans. *Proceedings of the Royal Society of London B*, 273 (1583), 141–147. <https://doi.org/10.1098/rspb.2005.3299>
- [12] Klimovsky, A.I. & Kotov, A.A. (2015) Cladocera (Crustacea, Branchiopoda) of Central Yakutia 3. Taxa from the Chydorus sphaericus s. l. species group (Anomopoda, Chydoridae). *Zoologicheskiy Zhurnal*, 94, 1257–1267. <https://doi.org/10.7868/S0044513415110057>
- [13] Korovchinsky, N.M. (2004) Cladocerans of the order Ctenopoda of the world fauna (morphology, systematics, ecology, biogeography). KMK Press, Moscow, 410 pp. [in Russian]
- [14] Korovchinsky, N.M. (2011) Description of Sarsitalona cf. fernandoi (Rane, 1983) (Crustacea: Cladocera: Sididae) found in the south of Vietnam. *Zootaxa*, 3129 (1), 29–38. <https://doi.org/10.11646/zootaxa.3129.1.2>
- [15] Kotov, A. A. (2013). Morphology and phylogeny of the Anomopoda (Crustacea: Cladocera). KMK, Moscow, 638 pp. [in Russian with English abstract]
- [16] Kotov, A.A., Van Damme, K., Bekker, E.I., Siboualipha, S., Silva-Briano, M., Ortiz, A.A., de la Rosa, R.G. & Sanoamuang, L. (2013) Cladocera (Crustacea: Branchiopoda) of Vientiane province and municipality, Laos. *Journal of Limnology*, 72 (S2), 81–180. <https://doi.org/10.4081/jlimnol.2013.s2.e6>
- [17] Kotov, A.A., Van Damme, K. & Elías-Gutiérrez, M. (2003) Differentiation between African Leydigia ciliata Gauthier, 1939 and Neotropical L. cf. striata Birabén, 1939 (Chydoridae, Anomopoda, Cladocera). *Hydrobiologia*, 505, 179–197. <https://doi.org/10.1023/B:HYDR.0000007246.04478.6a>
- [18] Michael, R.G. & Hann, B.J. (1979) On the resurrection of the cladoceran species Chydorus reticulatus Daday, 1898 (Chydoridae, Cladocera) and its relationship to Chydorus ventricosus Daday, 1898. *Hydrobiologia*, 65, 225–232. <https://doi.org/10.1007/BF00038863>
- [19] Neretina, A.N. & Sinev, A.Y. (2016) A revision of the genus Leberis Smirnov, 1989 (Cladocera: Chydoridae) in the Old World and Australia. *Zootaxa*, 4079 (5), 501–533. <https://doi.org/10.11646/zootaxa.4079.5.1>
- [20] Olmstead, A.W. & LeBlanc, G.A. (2001) Temporal and quantitative changes in sexual reproductive cycling of the cladoceran *Daphnia magna* by a juvenile hormone analog. *Journal of Experimental Zoology*, 290 (2), 148–155. <https://doi.org/10.1002/jez.1044>
- [21] Rajapaksa, R. & Fernando, C.H. (1982) The first description of the male and ephippial female of *Dadaya macrops* (Daday, 1898) (Cladocera, Chydoridae), with additional notes on this common tropical species. *Canadian Journal of Zoology*, 60 (8), 1841–1850. <https://doi.org/10.1139/z82-239>
- [22] Rajapaksa, R. & Fernando, C.H. (1985) Reallocation of *Indialona macronyx* (Daday, 1898) (Cladocera, Chydoridae) to the genus *Alona*. *Canadian Journal of Zoology*, 63 (4), 970–976. <https://doi.org/10.1139/z85-143>
- [23] Rajapaksa, R. & Fernando, C.H. (1986) Tropical species of *Kurzia* (Crustacea, Cladocera), with a description of *Kurzia brevilabris* sp. nov. *Canadian Journal of Zoology*, 64, 2590–2602. <https://doi.org/10.1139/z86-379>
- [24] Rajapaksa, R. & Fernando, C.H. (1987a) A redescription of *Euryalona orientalis* (Daday, 1898), with a consideration of the other species in the genus *Euryalona* (Cladocera: Chydoridae). *Hydrobiologia*, 150, 75–90. <https://doi.org/10.1007/BF00006611>
- [25] Rajapaksa, R. & Fernando, C.H. (1987b) Redescription and assignment of *Alona globulosa* Daday, 1898 to a new genus *Notoalona* and a description of *Notoalona freyi* sp. nov. *Hydrobiologia*, 144, 131–153. <https://doi.org/10.1007/BF00014527>
- [26] Sars, G.O. (1888) Additional notes on Australian Cladocera raised from dried mud. *Forhandlinger i Videnskabs-Selskabet i Christiania*, 1888, 1–74.
- [27] Sars, G.O. (1896) On fresh-water Entomostraca from the neighbourhood of Sydney, partly raised from dried mud. *Archiv for Matematik og Naturvidenskab Christiania*, 18, 1–81.
- [28] Sars, G.O. (1901) Contributions to the knowledge of the freshwater Entomostraca of South America, as shown by artificial hatching from dried material. 1. Cladocera. *Archiv for Matematik og Naturvidenskab Christiania*, 23, 1–102.
- [29] Sinev, A.Y. (2011) Redescription of the rheophilous cladocera *Camptocercus vietnamensis* Than, 1980 (Cladocera: Anomopoda: Chydoridae). *Zootaxa*, 2934 (1), 53–60. <https://doi.org/10.11646/zootaxa.2934.1.5>
- [30] Sinev, A.Y. (2012) *Alona kotovi* sp. nov., a new species of Aloninae (Cladocera: Anomopoda: Chydoridae) from South Vietnam. *Zootaxa*, 3475 (1), 45–54. <https://doi.org/10.11646/zootaxa.3475.1.4>
- [31] Sinev, A.Y. (2014) A new and a rare species of Chydorus Leach, 1816 (Branchiopoda: Cladocera: Anomopoda) from Cat Tien National Park, Vietnam. *Zootaxa*, 3861 (2), 127–144. <https://doi.org/10.11646/zootaxa.3861.2.2>

- [32] Sinev, A.Y. (2017) Morphology, systematics and zoogeography of the cladocerans of subfamily Aloninae (Cladocera: Anomopoda: Chydoridae). Dr.Sci. Thesis, Biological Faculty of M.V. Lomonosov Moscow State University, Moscow, 428 pp.
- [33] Sinev, A.Y., Gu, Y. & Han, B. (2015) Cladocera of Hainan Island, China. Zootaxa, 4006 (3), 569-585. <https://doi.org/10.11646/zootaxa.4006.3.9>
- [34] Sinev, A.Y. & Korovchinsky, N.M. (2013) Cladocera (Crustacea: Branchiopoda) of Cat Tien National Park, South Vietnam. Journal of Limnology, 70 (2), 125-141. <https://doi.org/10.4081/jlimnol.2013.s2.e8>
- [35] Sinev, A.Y. & Kotov, A.A. (2012) New and rare Aloninae (Cladocera: Anomopoda: Chydoridae) from Indochina. Zootaxa, 3334 (1), 1-28. <https://doi.org/10.11646/zootaxa.3334.1.1>
- [36] Sinev, A.Y. & Sanoamuang, L. (2011) Hormonal induction of males as a method for studying tropical cladocerans: description of males of four chydorid species (Cladocera: Anomopoda: Chydoridae). Zootaxa, 2826 (1), 45-56. <https://doi.org/10.11646/zootaxa.2826.1.2>
- [37] Sinev, A.Y., Van Damme, K. & Kotov, A.A. (2005) Redescription of tropical-temperate cladocerans *Alona diaphana* King, 1853 and *Alona davidi* Richard, 1895 and their translocation to Leberis Smirnov, 1989 (Branchiopoda: Anomopoda: Chydoridae). Arthropoda Selecta, 14 (3), 183-205.
- [38] Sinev, A.Y. & Yusoff, F.M. (2015) Cladocera (Crustacea: Branchiopoda) of Sabah state in Borneo Island, Malaysia. Zootaxa, 4000 (5), 581-591. <https://doi.org/10.11646/zootaxa.4000.5.7>
- [39] Sinev, A.Y. & Yusoff, F.M. (2016) A new species of genus *Ephemeroporus* Frey, 1982 (Cladocera: Anomopoda: Chydoridae) from Peninsular Malaysia. Zootaxa, 4117 (3), 399-410. <https://doi.org/10.11646/zootaxa.4117.3.7>
- [40] Sinev, A.Y. & Yusoff, F. M. (2018) New data on Cladocera (Crustacea: Branchiopoda) of Sabah State, Borneo Island, Malaysia. Zootaxa, 4438 (2), 362-372. <https://doi.org/10.11646/zootaxa.4438.2.10>
- [41] Tiang-nga, S., Sinev, A.Y. & Sanoamuang, L. (2020) High diversity of Cladocera (Crustacea: Branchiopoda) in a Ramsar site lake Kud-Thing, Northeast Thailand. Zootaxa, 4780 (2), 275-290. <https://doi.org/10.11646/zootaxa.4780.2.3>
- [42] Van Damme, K., Sinev, A.Y. & Dumont, H.J. (2011) Separation of *Anthalona* gen.n. from *Alona* Baird, 1843 (Branchiopoda: Cladocera: Anomopoda): morphology and evolution of scraping stenothermic alonines. Zootaxa, 2875 (2), 1-64. <https://doi.org/10.11646/zootaxa.4438.2.10>
- [43] Van Damme, K. & Maiphae, S. (2013) *Salinalona* gen. nov., an euryhaline chydorid lineage (Crustacea: Branchiopoda: Cladocera: Anomopoda) from the Oriental region. Journal of Limnology, 72 (S2), 142-173. <https://doi.org/10.4081/jlimnol.2013.s2.e9>
- [44] Van Damme, K., Maiphae, S. & Sa-ardrit, P. (2013) Inland swamps in South East Asia harbour hidden cladoceran diversities: species richness and the description of new paludal Chydoridae (Crustacea: Branchiopoda: Cladocera) from Southern Thailand. Journal of Limnology, 72 (S2), 174-208. <https://doi.org/10.4081/jlimnol.2013.s2.e10>
- [45] Van Damme, K. & Dumont, H.J. (2010) Cladocera of the Lençóis Maranhenses (NE-Brazil): faunal composition and a reappraisal of Sars' Method. Brazilian Journal of Biology, 70 (3), 755-779. <https://doi.org/10.1590/S1519-69842010000400008>
- [46] Van Damme, K. & Sinev, A.Y. (2013) Tropical Amphi-Pacific disjunctions in the Cladocera (Crustacea: Branchiopoda). Journal of Limnology, 70 (S2), 209-244. <https://doi.org/10.4081/jlimnol.2013.s2.e11>
- [47] Venkataraman, K. (1995) Cladoceran males from the Indian region. Journal of Bombay Natural History Society, 92, 378-385.
- [48] Yalim, F.B. & Chiplak, B. (2010) Redescription of *Ephemeroporus barroisi* (Richard, 1894) (Cladocera, Chydoridae) on the basis of material from Mediterranean Anatolia (Turkey). Turkish Journal of Fisheries and Aquatic Sciences, 10, 551-558. <https://doi.org/10.4194/trjfas.2010.0416>