



**X Международная конференция молодых
учёных по химии
«МЕНДЕЛЕЕВ-2017»**

**II школа-конференция «Направленный
дизайн веществ и материалов с заданными свойствами»**



Mendeleev2017

**Сборник тезисов
докладов**

**4-7 апреля 2017 г.
Санкт-Петербург**

THE REACTIONS OF ALKYLATION OF CARBOXYLATE PHOSPHABETAINES


Aksunova A.F., Bakhtiyarov D.I., Morozov M.V., Minnullin R.R., Galkina I.V., Galkin V.I., Bakhtiyarova Yu.V.

Kazan Federal University
Kazan, Russia
Graduate student
alfiya-nizamieva@mail.ru



Alkylation of monocarboxylate betaine leads to the formation of phosphonium salt **1**. The structure of obtained compound has been confirmed by spectral methods and X-ray diffraction analysis (pic.1).



In the case of reaction proceeding in the media of acetonitrile under heating, decarboxylation occurs with the formation of phosphonium salt **2**.



Similar reaction of dicarboxylate phosphobetaine proceeds slightly different. The final product of the reaction of alkylation depends on conditions. In the case of using of ethanol as a solvent the alkylation reaction of dicarboxylate betaine with methyl iodide leads to the formation of phosphonium salt which contains two ester groups. The structure of phosphonium salt **3** has been confirmed by a single crystal X-ray diffraction studies.



Picture 1. Molecular structure of (+-ethoxy-4-oxobutyl)triphenylphosphonium iodide **1** in the crystal



Picture 2. Molecular structure of bis(3-ethoxy-3-oxopropyl)diphenylphosphonium triiodide **3** in the crystal

The work is performed according to the Russian Government Program of Competitive Growth of Kazan Federal University.