

Collagen XIII secures pre- and postsynaptic integrity of the neuromuscular synapse

Härönen H., Zainul Z., Tu H., Naumenko N., Sormunen R., Miinalainen I., Shakirzyanova A., Oikarainen T., Abdullin A., Martin P., Santoleri S., Koistinaho J., Silman I., Giniatullin R., Fox M., Heikkinen A., Pihlajaniemi T.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© The Author 2017. Published by Oxford University Press. All rights reserved. Both transmembrane and extracellular cues, one of which is collagen XIII, regulate the formation and function of the neuromuscular synapse, and their absence results in myasthenia. We show that the phenotypical changes in collagen XIII knock-out mice are milder than symptoms in human patients, but the Col13a1 $-/-$ mice recapitulate major muscle findings of congenital myasthenic syndrome type 19 and serve as a disease model. In the lack of collagen XIII neuromuscular synapses do not reach full size, alignment, complexity and function resulting in reduced muscle strength. Collagen XIII is particularly important for the preterminal integrity, and when absent, destabilization of the motor nerves results in muscle regeneration and in atrophy especially in the case of slow muscle fibers. Collagen XIII was found to affect synaptic integrity through binding the ColQ tail of acetylcholine esterase. Although collagen XIII is a muscle-bound transmembrane molecule, it also undergoes ectodomain shedding to become a synaptic basal lamina component. We investigated the two forms' roles by novel Col13a1 tm/tm mice in which ectodomain shedding is impaired. While postsynaptic maturation, terminal branching and neurotransmission was exaggerated in the Col13a1 tm/tm mice, the transmembrane form's presence sufficed to prevent defects in transsynaptic adhesion, Schwann cell invagination/retraction, vesicle accumulation and acetylcholine receptor clustering and acetylcholinesterase dispersion seen in the Col13a1 $-/-$ mice, pointing to the transmembrane form as the major conductor of collagen XIII effects. Altogether, collagen XIII secures postsynaptic, synaptic and presynaptic integrity, and it is required for gaining and maintaining normal size, complexity and functional capacity of the neuromuscular synapse.

<http://dx.doi.org/10.1093/hmg/ddx101>

References

- [1] Engel, A.G., Shen, X.M., Selcen, D. and Sine, S.M. (2015) Congenital myasthenic syndromes: Pathogenesis, diagnosis, and treatment. *Lancet Neurol.*, 14, 420-434
- [2] Rodriguez Cruz, P.M., Palace, J. and Beeson, D. (2014) Congenital myasthenic syndromes and the neuromuscular junction. *Curr. Opin. Neurol.*, 27, 566-575
- [3] Latvanlehto, A., Fox, M.A., Sormunen, R., Tu, H., Oikarainen, T., Koski, A., Naumenko, N., Shakirzyanova, A., Kallio, M., Ilves, M., et al. (2010) Muscle-derived collagen XIII regulates maturation of the skeletal neuromuscular junction. *J. Neurosci.*, 30, 12230-12241

- [4] Hägg, P., Rehn, M., Huhtala, P., Väisänen, T., Tamminen, M. and Pihlajaniemi, T. (1998) Type XIII collagen is identified as a plasma membrane protein. *J. Biol. Chem.*, 273, 15590-15597
- [5] Väisänen, M.R., Väisänen, T. and Pihlajaniemi, T. (2004) The shed ectodomain of type XIII collagen affects cell behaviour in a matrix-dependent manner. *Biochem. J.*, 380, 685-693
- [6] Logan, C.V., Cossins, J., Rodriguez Cruz, P.M., Parry, D.A., Maxwell, S., Martinez-Martinez, P., Riepsaame, J., 2088 | *Human Molecular Genetics*, 2017, Vol. 26, No. 11 Abdelhamed, Z.A., Lake, A.V., Moran, M., et al. (2015) Congenital myasthenic syndrome type 19 is caused by mutations in COL13A1, encoding the atypical non-fibrillar collagen type XIII alpha1 chain. *Am. J. Hum. Genet.*, 97, 878-885
- [7] Ohno, K., Brengman, J., Tsujino, A. and Engel, A.G. (1998) Human endplate acetylcholinesterase deficiency caused by mutations in the collagen-like tail subunit (ColQ) of the asymmetric enzyme. *Proc. Natl. Acad. Sci. U. S. A.*, 95, 9654-9659
- [8] Donger, C., Krejci, E., Serradell, A.P., Eymard, B., Bon, S., Nicole, S., Chateau, D., Gary, F., Fardeau, M., Massoulié, J., et al. (1998) Mutation in the human acetylcholinesterase-associated collagen gene, COLQ, is responsible for congenital myasthenic syndrome with end-plate acetylcholinesterase deficiency (Type Ic). *Am. J. Hum. Genet.*, 63, 967-975
- [9] Feng, G., Krejci, E., Molgo, J., Cunningham, J.M., Massoulié, J. and Sanes, J.R. (1999) Genetic analysis of collagen Q: Roles in acetylcholinesterase and butyrylcholinesterase assembly and in synaptic structure and function. *J. Cell Biol.*, 144, 1349-1360
- [10] Arikawa-Hirasawa, E., Rossi, S.G., Rotundo, R.L. and Yamada, Y. (2002) Absence of acetylcholinesterase at the neuromuscular junctions of perlecan-null mice. *Nat. Neurosci.*, 5, 119-123
- [11] Cartaud, A., Strohlic, L., Guerra, M., Blanchard, B., Lambergeon, M., Krejci, E., Cartaud, J. and Legay, C. (2004) MuSK is required for anchoring acetylcholinesterase at the neuromuscular junction. *J. Cell Biol.*, 165, 505-515
- [12] Karmouch, J., Dobbertin, A., Sigoillot, S. and Legay, C. (2013) Developmental consequences of the ColQ/MuSK interactions. *Chem. Biol. Interact.*, 203, 287-291
- [13] Sigoillot, S.M., Bourgeois, F., Lambergeon, M., Strohlic, L. and Legay, C. (2010) ColQ controls postsynaptic differentiation at the neuromuscular junction. *J. Neurosci.*, 30, 13-23
- [14] Nishimune, H., Sanes, J.R. and Carlson, S.S. (2004) A synaptic laminin-calcium channel interaction organizes active zones in motor nerve terminals. *Nature*, 432, 580-587
- [15] Noakes, P.G., Gautam, M., Mudd, J., Sanes, J.R. and Merlie, J.P. (1995) Aberrant differentiation of neuromuscular junctions in mice lacking s-laminin/laminin beta 2. *Nature*, 374, 258-262
- [16] Lin, W., Burgess, R.W., Dominguez, B., Pfaff, S.L., Sanes, J.R. and Lee, K.F. (2001) Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse. *Nature*, 410, 1057-1064
- [17] Huze, C., Bauche, S., Richard, P., Chevessier, F., Goillot, E., Gaudon, K., Ben Ammar, A., Chaboud, A., Grosjean, I., Lecuyer, H.A., et al. (2009) Identification of an agrin mutation that causes congenital myasthenia and affects synapse function. *Am. J. Hum. Genet.*, 85, 155-167
- [18] Fox, M.A., Sanes, J.R., Borza, D.B., Eswarakumar, V.P., Fassler, R., Hudson, B.G., John, S.W., Ninomiya, Y., Pedchenko, V., Pfaff, S.L., et al. (2007) Distinct target-derived signals organize formation, maturation, and maintenance of motor nerve terminals. *Cell*, 129, 179-193
- [19] Fox, M.A., Ho, M.S., Smyth, N. and Sanes, J.R. (2008) A synaptic nidogen: Developmental regulation and role of nidogen-2 at the neuromuscular junction. *Neural Develop.*, 3, 24
- [20] Schiaffino, S., Rossi, A.C., Smerdu, V., Leinwand, L.A. and Reggiani, C. (2015) Developmental myosins: Expression patterns and functional significance. *Skelet. Muscle*, 5, 22-015-0046-6. eCollection 2015
- [21] Thomsen, R.H. and Wilson, D.F. (1983) Effects of 4-aminopyridine and 3, 4-diaminopyridine on transmitter release at the neuromuscular junction. *J. Pharmacol. Exp. Ther.*, 227, 260-265
- [22] Snellman, A., Tu, H., Väisänen, T., Kvist, A.P., Huhtala, P. and Pihlajaniemi, T. (2000) A short sequence in the N-terminal region is required for the trimerization of type XIII collagen and is conserved in other collagenous transmembrane proteins. *embo J.*, 19, 5051-5059
- [23] Kvist, A.P., Latvanlehto, A., Sund, M., Eklund, L., Väisänen, T., Hägg, P., Sormunen, R., Komulainen, J., Fässler, R. and Pihlajaniemi, T. (2001) Lack of cytosolic and transmembrane domains of type XIII collagen results in progressive myopathy. *Am. J. Pathol.*, 159, 1581-1592
- [24] Hägg, P., Väisänen, T., Tuomisto, A., Rehn, M., Tu, H., Huhtala, P., Eskelinen, S. and Pihlajaniemi, T. (2001) Type XIII collagen: A novel cell adhesion component present in a range of cell-matrix adhesions and in the intercalated discs between cardiac muscle cells. *Matrix Biol.*, 19, 727-742
- [25] Ellman, G.L., Courtney, K.D., Andres, V., Jr. and Feather-Stone, R.M. (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.*, 7, 88-95
- [26] Yang, Y., Coleman, M., Zhang, L., Zheng, X. and Yue, Z. (2013) Autophagy in axonal and dendritic degeneration. *Trends Neurosci.*, 36, 418-428

- [27] Tu, H., Sasaki, T., Snellman, A., Gohring, W., Pirilä, P., Timpl, R. and Pihlajaniemi, T. (2002) The type XIII collagen ectodomain is a 150-nm rod and capable of binding to fibronectin, nidogen-2, perlecan, and heparin. *J. Biol. Chem.*, 277, 23092-23099
- [28] Tanaka, T., Wakabayashi, T., Oizumi, H., Nishio, S., Sato, T., Harada, A., Fujii, D., Matsuo, Y., Hashimoto, T. and Iwatsubo, T. (2014) CLAC-P/collagen type XXV is required for the intramuscular innervation of motoneurons during neuromuscular development. *J. Neurosci.*, 34, 1370-1379
- [29] Su, J., Stenbjorn, R.S., Gorse, K., Su, K., Hauser, K.F., Ricard-Blum, S., Pihlajaniemi, T. and Fox, M.A. (2012) Target-derived matricryptins organize cerebellar synapse formation through alpha3beta1 integrins. *Cell. Rep.*, 2, 223-230
- [30] Nykvist, P., Tu, H., Ivaska, J., Kapyla, J., Pihlajaniemi, T. and Heino, J. (2000) Distinct recognition of collagen subtypes by alpha(1)beta(1) and alpha(2)beta(1) integrins. alpha(1)beta(1) mediates cell adhesion to type XIII collagen. *J. Biol. Chem.*, 275, 8255-8261
- [31] Martin, P.T., Kaufman, S.J., Kramer, R.H. and Sanes, J.R. (1996) Synaptic integrins in developing, adult, and mutant muscle: Selective association of alpha1, alpha7A, and alpha7B integrins with the neuromuscular junction. *Dev. Biol.*, 174, 125-139
- [32] Murray, L.M., Talbot, K. and Gillingwater, T.H. (2010) Review: Neuromuscular synaptic vulnerability in motor neurone disease: Amyotrophic lateral sclerosis and spinal muscular atrophy. *Neuropathol. Appl. Neurobiol.*, 36, 133-156
- [33] Ylö nen, R., Kyrö nlahti, T., Sund, M., Ilves, M., Lehenkari, P., Tuukkanen, J. and Pihlajaniemi, T. (2005) Type XIII collagen strongly affects bone formation in transgenic mice. *J. Bone Miner. Res.*, 20, 1381-1393
- [34] Sakai, K. and Miyazaki, J. (1997) A transgenic mouse line that retains cre recombinase activity in mature oocytes irrespective of the cre transgene transmission. *Biochem. Biophys. Res. Commun.*, 237, 318-324
- [35] Shi, Y., Ivannikov, M.V., Walsh, M.E., Liu, Y., Zhang, Y., Jaramillo, C.A., Macleod, G.T. and Van Remmen, H. (2014) The lack of CuZnSOD leads to impaired neurotransmitter release, neuromuscular junction destabilization and reduced muscle strength in mice. *PLoS One*, 9, e100834
- [36] Dudai, Y., Silman, I., Kalderon, N. and Blumberg, S. (1972) Purification by affinity chromatography of acetylcholinesterase from electric organ tissue of the electric eel subsequent to tryptic treatment. *Biochim. Biophys. Acta*, 268, 138-157
- [37] Dudai, Y., Herzberg, M. and Silman, I. (1973) Molecular structures of acetylcholinesterase from electric organ tissue of the electric eel. *Proc. Natl. Acad. Sci. U. S. A.*, 70, 2473-2476
- [38] Lwebuga-Mukasa, J.S., Lappi, S. and Taylor, P. (1976) Molecular forms of acetylcholinesterase from *torpedo californica*: Their relationship to synaptic membranes. *Biochemistry*, 15, 1425-1434
- [39] Sussman, J.L., Harel, M., Frolow, F., Varon, L., Toker, L., Futerman, A.H. and Silman, I. (1988) Purification and crystallization of a dimeric form of acetylcholinesterase from *torpedo californica* subsequent to solubilization with phosphatidylinositol-specific phospholipase C. *J. Mol. Biol.*, 203, 821-823