ISSN 1070-3632, Russian Journal of General Chemistry, 2017, Vol. 87, No. 6, pp. 1143–1147. © Pleiades Publishing, Ltd., 2017. Original Russian Text © Ya.V. Veremeichik, O.A. Tevs, D.B. Krivolapov, O.A. Lodochnikova, V.V. Plemenkov, 2017, published in Zhurnal Obshchei Khimii, 2017, Vol. 87, No. 6, pp. 921–926.

Synthesis and Structure of Novel Substituted N-Sulfinylanilines

Ya. V. Veremeichik^a, O. A. Tevs^a, D. B. Krivolapov^b,
O. A. Lodochnikova^{b,c}*, and V. V. Plemenkov^{a,c}

^a Chemical Biological Institute, Kant Baltic Federal University, Kalinigrad, Russia ^bArbuzov Institute of Organic and Physical Chemistry, Kazan Research Center, Russian Academy of Sciences, ul. Akademika Arbuzova 8, Kazan, Tatarstan, 420088 Russia *e-mail: lod_olga@mail.ru ^c Butlerov Chemical Institute, Kazan Federal University, Kazan, Tatarstan, Russia

Received February 15, 2017

Abstract—*N*-Sulfinylanilines derived from 4-bromoaniline, 3-nitroaniline, and 4,4'-di(ethane-1,2-diyl)dianiline were synthesized. X-ray diffraction analysis of 4-bromo-*N*-sulfinylaniline, 3-nitro-*N*-sulfinylaniline, and 4,4'- (ethane-1,2-diyl)di-*N*-sulfinylaniline was performed. The sulfinyl function in the planar conformation of the Ar-NSO fragment was found to have *Z* configuration. The nature of intra- and intermolecular structure-forming interactions was established.

Keywords: N-sulfinylanilines, X-ray diffraction analysis, intra- and intermolecular interactions

DOI: 10.1134/S107036321706007X

The aromatic sulfinylamine function (Ar–N=S=O) is quite reactive, which is manifested in that it is highly prone to addition across the N=S bond. Especially characteristic of aromatic sulfinylamines are Diels–Alder reactions, where they can act as a dienophile (route *a*) [1, 2], diene (route *b*) [3], or 1,3-dipolarophile (in 1,3-dipolar cycloaddition, route *c*) [4, 5] (Scheme 1). In view of the fact that such reactions

provide a facile synthetic approach to cyclic sulfonamides, in the present work we decided to study in more detail the structure of the starting aromatic sulfinylanilines.

N-Sulfinylanilines **1–3** were prepared in yields of 84–90% by the reactions of the corresponding anilines with thionyl chloride in benzene under reflux (Scheme 2).

Scheme 1.

