

Towards water-free biobanks: long-term dry-preservation at room temperature of desiccation-sensitive enzyme luciferase in air-dried insect cells

Kikuta S., Watanabe S., Sato R., Gusev O., Nesmelov A., Sogame Y., Cornette R., Kikawada T.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 The Author(s). Desiccation-tolerant cultured cells Pv11 derived from the anhydrobiotic midge embryo endure complete desiccation in an ametabolic state and resume their metabolism after rehydration. These features led us to develop a novel dry preservation technology for enzymes as it was still unclear whether Pv11 cells could preserve an exogenous enzyme in the dry state. This study shows that Pv11 cells protect an exogenous desiccation-sensitive enzyme, luciferase (Luc), preserving the enzymatic activity even after dry storage for 372 days at room temperature. A process including preincubation with trehalose, dehydration, storage, and rehydration allowed Pv11 (Pv11-Luc) cells stably expressing luciferase to survive desiccation and still emit luminescence caused by luciferase after rehydration. Luminescence produced by luciferase in Pv11-Luc cells after rehydration did not significantly decrease in presence of a translation inhibitor, showing that the activity did not derive from de novo enzyme synthesis following the resumption of cell metabolism. These findings indicate that the surviving Pv11 cells almost completely protect luciferase during desiccation. Lacking of the preincubation step resulted in the loss of luciferase activity after rehydration. We showed that preincubation with trehalose associated to induction of desiccation tolerance-related genes in Pv11 cells allowed effective in vivo preservation of enzymes in the dry state.

<http://dx.doi.org/10.1038/s41598-017-06945-y>

References

- [1] Cooper, G. M. The Central Role of Enzymes as Biological Catalysts in The Cell: A Molecular Approach. 2nd edition. Sunderland (MA): (Sinauer Associates, 2000) NCBI Bookshelf ID: NBK9921 <https://www.ncbi.nlm.nih.gov/books/NBK9921/>
- [2] Berg, J. M., Tymoczko, J. L. & Stryer, L. Biochemistry. 5th edition. (NY) (Freeman W. H., 2002) NCBI Bookshelf ID: NBK21154 <https://www.ncbi.nlm.nih.gov/books/NBK21154/>
- [3] Anjem, A. & Imlay, J. A. Mononuclear iron enzymes are primary targets of hydrogenperoxide stress. *J Biol Chem* 287, 15544-15556 (2012)
- [4] Lodish, H. et al. Protein Structure and Function in Molecular Cell Biology 4th edition. (NY) (Freeman W. H., 2000) 963 NCBI Bookshelf ID: NBK21475 <https://www.ncbi.nlm.nih.gov/books/NBK21475/>
- [5] Bradbury, S. L. & Jakoby, W. B. Glycerol as an enzyme-stabilizing agent: effects on aldehyde dehydrogenase. *Proc Natl Acad Sci USA* 69, 2373-2376 (1972)
- [6] Carpenter, J. F., Martin, B., Loomis, S. H. & Crowe, J. H. Long-term preservation of dried phosphofructokinase by sugars and sugar/zinc mixtures. *Cryobiol* 25, 372-376 (1988)
- [7] Argüelles, J. C. Physiological roles of trehalose in bacteria and yeasts: a comparative analysis. *Arch Microbiol* 174, 217-224 (2000)

- [8] Crowe, J. H., Hoekstra, F. A. & Crowe, L. M. Anhydrobiosis. *Annu Rev Physiol* 54, 579-599 (1992)
- [9] Sakurai, M. et al. Vitrification is essential for anhydrobiosis in an African chironomid, *Polypedilum vanderplanki*. *Proc Natl Acad Sci USA* 105, 5093-5098 (2008)
- [10] Golovina, E. A., Golovin, A. V., Hoekstra, F. A. & Faller, R. Water replacement hypothesis in atomic detail-factors determining the structure of dehydrated bilayer stacks. *Biophys J* 97, 490-429 (2009)
- [11] Crowe, J. H., Crowe, L. M., Carpenter, J. F. & Aurell Wistrom, C. Stabilization of dry phospholipid bilayers and proteins by sugars. *Biochem J* 242, 1-10 (1987)
- [12] Green, J. L. & Angell, C. A. Phase relations and vitrification in saccharide-water solutions and the trehalose anomaly. *J Phys Chem* 93, 2880-2882 (1989)
- [13] Belton, P. S. & Gil, A. M. IR and Raman spectroscopic studies of the interaction of trehalose with hen egg white lysozyme. *Biopolymers* 34, 957-61 (1994)
- [14] Grasmeyer, N., Stankovic, M., de Waard, H., Frijlink, H. W. & Hinrichs, W. L. Unraveling protein stabilization mechanisms: vitrification and water replacement in a glass transition temperature controlled system. *Biochim Biophys Acta* 1834, 763-769 (2013)
- [15] Kreilgaard, L., Frokjaer, S., Flink, J. M., Randolph, T. W. & Carpenter, J. F. Effects of additives on the stability of *Humicola lanuginosa* lipase during freeze-drying and storage in the dried solid. *J Pharm Sci* 88, 281-290 (1999)
- [16] O'Fágáin, C. Storage and lyophilisation of pure proteins. *Methods Mol Biol* 681, 179-202 (2011)
- [17] Patel, S. M. & Pikal, M. J. Emerging Freeze-Drying Process Development and Scale-up Issues. *AAPS PharmSciTech* 12, 372-378 (2011)
- [18] Sales, K., Brandt, W., Rumbak, E. & Lindsey, G. The LEA-like protein HSP 12 in *Saccharomyces cerevisiae* has a plasma membrane location and protects membranes against desiccation and ethanol-induced stress. *Biochim Biophys Acta* 1463, 267-278 (2000)
- [19] Kikawada, T. et al. Dehydration-induced expression of LEA proteins in an anhydrobiotic chironomid. *Biochem Biophys Res Commun* 348, 56-61 (2006)
- [20] Warner, A. H., Chakrabortee, S., Tunnacliffe, A. & Clegg, J. S. Complexity of the heat-soluble LEA proteome in *Artemia* species. *Comp Biochem Physiol Part D Genomics Proteomics* 7, 260-267 (2012)
- [21] Tanaka, S. et al. Novel mitochondria-targeted heat-soluble proteins identified in the anhydrobiotic Tardigrade improve osmotic tolerance of human cells. *PLoS One* 10, e0118272 (2015)
- [22] Garay-Arroyo, A., Colmenero-Flores, J. M., Garcarrubio, A. & Covarrubias, A. A. Highly hydrophilic proteins in prokaryotes and eukaryotes are common during conditions of water deficit. *J Biol Chem* 275, 5668-5674 (2000)
- [23] Campos, F., Cuevas-Velazquez, C., Fares, M. A., Reyes, J. L. & Covarrubias, A. A. Group 1 LEA proteins, an ancestral plant protein group, are also present in other eukaryotes, and in the archaea and bacteria domains. *Mol Genet Genomics* 288, 503-517 (2013)
- [24] Hinton, H. E. A new chironomid from Africa, the larva of which can be dehydrated without injury. *J Zool* 121, 371-380 (1951)
- [25] Watanabe, M., Kikawada, T. & Okuda, T. Increase of internal ion concentration triggers trehalose synthesis associated with cryptobiosis in larvae of *Polypedilum vanderplanki*. *J Exp Biol* 206, 2281-2286 (2003)
- [26] Mitsumasu, K. et al. Enzymatic control of anhydrobiosis-related accumulation of trehalose in the sleeping chironomid. *Polypedilum vanderplanki*. *FEBS J* 277, 4215-4228 (2010)
- [27] Hatanaka, R. et al. An abundant LEA protein in the anhydrobiotic midge, PvLEA4, acts as a molecular shield by limiting growth of aggregating protein particles. *Insect Biochem Mol Biol* 43, 1055-1067 (2013)
- [28] Gusev, O. et al. Anhydrobiosis-associated nuclear DNA damage and repair in the sleeping chironomid: linkage with radioresistance. *PLoS One* 5, e14008 (2010)
- [29] Cornette, R. & Kikawada, T. The Induction of Anhydrobiosis in the Sleeping Chironomid: Current Status of Our Knowledge. *IUBMB Life* 63, 419-429 (2011)
- [30] Watanabe, K., Imanishi, S., Akiduki, G., Cornette, R. & Okuda, T. Air-dried cells from the anhydrobiotic insect, *Polypedilum vanderplanki*, can survive long term preservation at room temperature and retain proliferation potential after rehydration. *Cryobiol* 73, 93-98 (2016)
- [31] Sogame, Y. et al. Establishment of gene transfer and gene silencing methods in a desiccation-tolerant cell line, Pv11. *Extremophiles* 21, 65-72 (2017)
- [32] Tsien, R. Y. The green fluorescent protein. *Annu Rev Biochem* 67, 509-544 (1998)
- [33] Tapia, H. & Koshland, D. E. Trehalose is a versatile and long-lived chaperone for desiccation tolerance. *Curr Biol* 24, 2758-2766 (2014)
- [34] Kikuta, S., Hou, B. H., Sato, R., Frommer, W. B. & Kikawada, T. FRET sensor-based quantification of intracellular trehalose in mammalian cells. *Biosci Biotechnol Biochem* 80, 162-165 (2015)
- [35] Gstraunthaler, G. Alternatives to the use of fetal bovine serum: serum-free cell culture. *ALTEX* 20, 275-281 (2003)

- [36] Borovikova, D., Teparić, R., Mrša, V. & Rapoport, A. Anhydrobiosis in yeast: cell wall mannoproteins are important for yeast *Saccharomyces cerevisiae* resistance to dehydration. *Yeast* 33, 347-53 (2016)
- [37] Singh, J. et al. Transcriptional response of *Saccharomyces cerevisiae* to desiccation and rehydration. *Appl Environ Microbiol* 71, 8752-8763 (2005)
- [38] Gusev, O. et al. Comparative genome sequencing reveals genomic signature of extreme desiccation tolerance in the anhydrobiotic midge. *Nat Commun* 5, 4784, doi: 10.1038/ncomms5784 (2014)
- [39] Nakahara, Y. et al. Cells from an anhydrobiotic chironomid survive almost complete desiccation. *Cryobiol* 60, 138-146 (2010)
- [40] Hatanaka, R. et al. Diversity of the expression profiles of late embryogenesis abundant (LEA) protein encoding genes in the anhydrobiotic midge *Polypedilum vanderplanki*. *Planta* 242, 451-459 (2015)
- [41] Vancha, A. R. et al. Use of polyethyleneimine polymer in cell culture as attachment factor and lipofection enhancer. *BMC Biotechnol* 4, 23 (2004)
- [42] Rice, R. H. & Green, H. Relation of protein synthesis and transglutaminase activity to formation of the cross-linked envelope during terminal differentiation of the cultured human epidermal keratinocyte. *J Cell Biol* 76, 705-711 (1978)
- [43] Kikuta, S. et al. Characterization of a ligand-gated cation channel based on an inositol receptor in the silkworm, *Bombyx mori*. *Insect Biochem Mol Biol* 74, 12-20 (2016).