

Spinal cord molecular and cellular changes induced by adenoviral vector- and cell-mediated triple gene therapy after severe contusion

Izmailov A., Povysheva T., Bashirov F., Sokolov M., Fadeev F., Garifulin R., Naroditsky B., Logunov D., Salafutdinov I., Chelyshev Y., Islamov R., Lavrov I.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Izmailov, Povysheva, Bashirov, Sokolov, Fadeev, Garifulin, Naroditsky, Logunov, Salafutdinov, Chelyshev, Islamov and Lavrov. The gene therapy has been successful in treatment of spinal cord injury (SCI) in several animal models, although it still remains unavailable for clinical practice. Surprisingly, regardless the fact that multiple reports showed motor recovery with gene therapy, little is known about molecular and cellular changes in the post-traumatic spinal cord following viral vector- or cell-mediated gene therapy. In this study we evaluated the therapeutic efficacy and changes in spinal cord after treatment with the genes encoding vascular endothelial growth factor (VEGF), glial cell-derived neurotrophic factor (GDNF), angiogenin (ANG), and neuronal cell adhesion molecule (NCAM) applied using both approaches. Therapeutic genes were used for viral vector- and cell-mediated gene therapy in two combinations: (1) VEGF+GDNF+NCAM and (2) VEGF+ANG+NCAM. For direct gene therapy adenoviral vectors based on serotype 5 (Ad5) were injected intrathecally and for cell-mediated gene delivery human umbilical cord blood mononuclear cells (UCB-MC) were simultaneously transduced with three Ad5 vectors and injected intrathecally 4 h after the SCI. The efficacy of both treatments was confirmed by improvement in behavioral (BBB) test. Molecular and cellular changes following post-traumatic recovery were evaluated with immunofluorescent staining using antibodies against the functional markers of motoneurons (Hsp27, synaptophysin, PSD95), astrocytes (GFAP, vimentin), oligodendrocytes (Olig2, NG2, Cx47) and microglial cells (Iba1). Our results suggest that both approaches with intrathecal delivery of therapeutic genes may support functional recovery of post-traumatic spinal cord via lowering the stress (down regulation of Hsp25) and enhancing the synaptic plasticity (up regulation of PSD95 and synaptophysin), supporting oligodendrocyte proliferation (up regulation of NG2) and myelination (up regulation of Olig2 and Cx47), modulating astrogliosis by reducing number of astrocytes (down regulation of GFAP and vimentin) and microglial cells (down regulation of Iba1).

<http://dx.doi.org/10.3389/fphar.2017.00813>

Keywords

Angiogenin, Gene therapy, Glial cell-derived neurotrophic factor, Glial cells, Human umbilical cord blood mononuclear cell, Neuronal cell adhesion molecule, Spinal cord injury, Vascular endothelial growth factor

References

- [1] Abdellatif, A. A., Pelt, J. L., Benton, R. L., Howard, R. M., Tsoulfas, P., Ping, P., et al. (2006). Gene delivery to the spinal cord: comparison between lentiviral, adenoviral, and retroviral vector delivery systems. *J. Neurosci. Res.* 84, 553-567. doi: 10.1002/jnr.20968
- [2] Bakshi, A., Hunter, C., Swanger, S., Lepore, A., and Fischer, I. (2004). Minimally invasive delivery of stem cells for spinal cord injury: advantages of the lumbar puncture technique. *J. Neurosurg. Spine.* 1, 330-337. doi: 10.3171/spi.2004.1.3.0330
- [3] Chaudhry, N., Desilva, U., and Smith, G. (2006). Cell adhesion molecule L1 modulates nerve-growth-factor-induced CGRP-IR fiber sprouting. *Exp. Neurol.* 202, 238-249. doi: 10.1016/j.expneurol.2006.06.001
- [4] Chen, H. K., Hung, H. F., Shyu, K. G., Wang, B. W., Sheu, J. R., Liang, Y. J., et al. (2005). Combined cord blood stem cells and gene therapy enhances angiogenesis and improves cardiac performance in mouse after acute myocardial infarction. *Eur. J. Clin. Invest.* 35, 677-686. doi: 10.1111/j.1365-2362.2005.01565.x
- [5] Chen, J., Bernreuther, C., Dihné, M., and Schachner, M. (2005). Cell adhesion molecule L1-transfected embryonic stem cells with enhanced survival support regrowth of corticospinal tract axons in mice after spinal cord injury. *J. Neurotrauma.* 22, 896-906. doi: 10.1089/neu.2005.22.896
- [6] Chou, A. K., Yang, M. C., Tsai, H. P., Chai, C. Y., Tai, M. H., Kwan, A. L., et al. (2014). Adenoviral-Mediated glial cell line-derived neurotrophic factor gene transfer has a protective effect on sciatic nerve following constriction-induced spinal cord injury. *PLoS ONE* 9:e92264. doi: 10.1371/journal.pone.0092264
- [7] Colombo, F., and Meldolesi, J. (2015). L1-CAM and N-CAM: from adhesion proteins to pharmacological targets. *Trends Pharmacol. Sci.* 36, 769-781. doi: 10.1016/j.tips.2015.08.004
- [8] Cui, Y.-F., Xu, J.-C., Hargus, G., Jakovcevski, I., Schachner, M., and Bernreuther, C. (2011). Embryonic stem cell-derived L1 overexpressing neural aggregates enhance recovery after spinal cord injury in mice. *PLoS ONE* 6:17126. doi: 10.1371/journal.pone.0017126
- [9] De Laporte, L., des Rieux, A., Tuinstra, H. M., Zelivyanskaya, M. L., De Clerck, N. M., Postnov, A. A., et al. (2011). Vascular endothelial growth factor and fibroblast growth factor 2 delivery from spinal cord bridges to enhance angiogenesis following injury. *J. Biomed. Mater. Res. A* 98, 372-382. doi: 10.1002/jbm.a.33112
- [10] Ding, X., Mao, B., Jiang, S., Li, S., and Deng, Y. (2005). Neuroprotective effect of exogenous vascular endothelial growth factor on rat spinal cord neurons in vitro hypoxia. *Chin. Med. J.* 118, 1644-1650
- [11] Figley, S. A., Liu, Y., Karadimas, S. K., Satkunendrarajah, K., Fettes, P., Spratt, S. K., et al. (2014). Delayed administration of a bio-engineered zinc-finger VEGF-A gene therapy is neuroprotective and attenuates allodynia following traumatic spinal cord injury. *PLoS ONE* 9:96137. doi: 10.1371/journal.pone.0096137
- [12] Franklin, T. B., Krueger-Naug, A. M., Clarke, D. B., Arrigo, A.-P., and Currie, R. W. (2005). The role of heat shock proteins Hsp70 and Hsp27 in cellular protection of the central nervous system. *Int. J. Hyperth.* 21, 379-392. doi: 10.1080/02656730500069955
- [13] Herrera, J. J., Nesic-Taylor, D. O., and Narayana, P. A. (2009). Reduced vascular endothelial growth factor expression in contusive spinal cord injury. *J. Neurotrauma.* 26, 995-1003 doi: 10.1089/neu.2008.0779
- [14] Ichim, T. E., Solano, F., Lara, F., Paris, F., Ugalde, F., Rodriguez, J. P., et al. (2010). Feasibility of combination allogeneic stem cell therapy for spinal cord injury: a case report. *Int. Arch. Med.* 3:30. doi: 10.1186/1755-768-3-30
- [15] Ikeda, Y., Fukuda, N., Wada, M., Matsumoto, T., Satomi, A., Yokoyama, S., et al. (2004). Development of angiogenic cell and gene therapy by transplantation of umbilical cord blood with vascular endothelial growth factor gene. *Hypertens. Res.* 27, 119-128. doi: 10.1291/hyres.27.119
- [16] Islamov, R. R., Hendricks, W. A., Katwa, L. C., McMurray, R. J., Pak, E. S., Spanier, N. S., et al. (2003). Effect of 17 beta-estradiol on gene expression in lumbar spinal cord following sciatic nerve crush injury in ovariectomized mice. *Brain Res.* 966, 65-75. doi: 10.1016/S0006-8993(02)04191-4
- [17] Islamov, R. R., Rizvanov, A. A., Fedotova, V. Y., Izmailov, A. A., Safiulloev, Z. Z., Garanina, E. E., et al. (2016). Tandem delivery of multiple therapeutic genes using umbilical cord blood cells improves symptomatic outcomes in ALS. *Mol. Neurobiol.* 54, 4756-4763. doi: 10.1007/s12035-016-0017-x
- [18] Islamov, R. R., Rizvanov, A. A., Mukhamedyarov, M. A., Salafutdinov, I. I., Garanina, E. E., Fedotova, V. Y., et al. (2015). Symptomatic improvement, increased life-span and sustained cell homing in amyotrophic lateral sclerosis after transplantation of human umbilical cord blood cells genetically modified with adeno-viral vectors expressing a neuro-protective factor and a neural cell adhesion 496 molecule. *Curr. Gene Ther.* 15, 266-276. doi: 10.2174/1566523215666150126122317
- [19] Islamov, R. R., Sokolov, M. E., Bashirov, F. V., Fadeev, F. O., Shmarov, M. M., Naroditskiy, B. S., et al. (2017). A pilot study of cell-mediated gene therapy for spinal cord injury in mini pigs. *Neurosci. Lett.* 644, 67-75. doi: 10.1016/j.neulet.2017.02.034
- [20] Ismailov, S. M., Barykova, I. A., Shmarov, M. M., Tarantul, V. Z., Barskov, I. V., Kucherianu, V. G., et al. (2014). Experimental approach to the gene therapy of motor neuron disease with the use of genes hypoxia-inducible factors. *Genetika* 50, 591-601. doi: 10.1134/S1022795414050056

- [21] Jakovcevski, I., Djogo, N., Hölters, L. S., Szpotowicz, E., and Schachner, M. (2013). Transgenic overexpression of the cell adhesion molecule L1 in neurons facilitates recovery after mouse spinal cord injury. *Neuroscience* 252, 1-12. doi: 10.1016/j.neuroscience.2013.07.067
- [22] Kieran, D., Sebastia, J., Greenway, M. J., King, M. A., Connaughton, D., Concannon, C. G., et al. (2008). Control of motoneuron survival by angiogenin. *J. Neurosci.* 28, 14056-14061. doi: 10.1523/JNEUROSCI.3399-08.2008
- [23] Kim, H. M., Hwang, D. H., Lee, J. E., Kim, S. U., and Kim, B. G. (2009). Ex vivo VEGF delivery by neural stem cells enhances proliferation of glial progenitors, angiogenesis, and tissue sparing after spinal cord injury. *PLoS ONE* 4:4987. doi: 10.1371/journal.pone.0004987
- [24] Lavdas, A. A., Chen, J., Papastefanaki, F., Chen, S., Schachner, M., Matsas, R., et al. (2010). Schwann cells engineered to express the cell adhesion molecule L1 accelerate myelination and motor recovery after spinal cord injury. *Exp. Neurol.* 221, 206-216. doi: 10.1016/j.expneurol.2009.10.024
- [25] Lee, H. J., Bian, S., Jakovcevski, I., Wu, B., Irintchev, A., and Schachner, M. (2012). Delayed applications of L1 and chondroitinase ABC promote recovery after spinal cord injury. *J. Neurotrauma* 29, 1850-1863. doi: 10.1089/neu.2011.2290
- [26] Lepore, A. C., Bakshi, A., Swanger, S. A., Rao, M. S., and Fischer, I. (2005). Neural precursor cells can be delivered into the injured cervical spinal cord by intrathecal injection at the lumbar cord. *Brain Res.* 1045, 206-216. doi: 10.1016/j.brainres.2005.03.050
- [27] Lim, S. T., Airavaara, M., and Harvey, B. K. (2010). Viral vectors for neurotrophic factor delivery: a gene therapy approach for neurodegenerative diseases of the CNS. *Pharmacol. Res.* 61, 14-26. doi: 10.1016/j.phrs.2009.10.002
- [28] Liu, J., Han, D., Wang, Z., Xue, M., Zhu, L., Yan, H., et al. (2013). Clinical analysis of the treatment of spinal cord injury with umbilical cord mesenchymal stem cells. *Cytherapy* 15, 185-191. doi: 10.1016/j.jcyt.2012.09.005
- [29] Liu, Y., Figley, S., Spratt, S. K., Lee, G., Ando, D., Surosky, R., et al. (2010). An engineered transcription factor which activates VEGF-A enhances recovery after spinal cord injury. *Neurobiol. Dis.* 37, 384-393. doi: 10.1016/j.nbd.2009.10.018
- [30] Mukhamedshina, Y. O., Garanina, E. E., Masgutova, G. A., Galieva, L. R., Sanatova, E. R., Chelyshev, Y. A., et al. (2016a). Assessment of glial scar, tissue sparing, behavioral recovery and axonal regeneration following acute transplantation of genetically modified human umbilical cord blood cells in a rat model of spinal cord contusion. *PLoS ONE* 11:e0151745. doi: 10.1371/journal.pone.0151745
- [31] Mukhamedshina, Y. O., Shaymardanova, G. F., Garanina, E. E., Salafutdinov, I. I., Rizvanov, A. A., Islamov, R. R., et al. (2016b). Adenoviral vector carrying glial cell-derived neurotrophic factor for direct gene therapy in comparison with human umbilical cord blood cell-mediated therapy of spinal cord injury in rat. *Spinal Cord* 54, 347-359. doi: 10.1038/sc.2015.161
- [32] Mukhamedshina, Y. O., Shaymardanova, G. F., Muhitov, A. R., Salafutdinov, I., Zarubina, V. N., Rizvanov, A., et al. (2012). Survival and differentiation of endogenous Schwann cells migrating into spinal cord under the influence of neurotrophic factors. *Cell. Transpl. Tissue Eng.* 7, 125-129
- [33] Nakajima, H., Uchida, K., Kobayashi, S., Inukai, T., Horiuchi, Y., Yayama, T., et al. (2007). Rescue of rat anterior horn neurons after spinal cord injury by retrograde transfection of adenovirus vector carrying brain-derived neurotrophic factor gene. *J. Neurotrauma* 24, 703-712. doi: 10.1089/neu.2006.0004
- [34] Niu, J., Li, C., Wu, H., Feng, X., Su, Q., Li, S., et al. (2015). Propidium iodide (PI) stains Nissl bodies and may serve as a quick marker for total neuronal cell count. *Acta Histochem.* 117, 182-187. doi: 10.1016/j.acthis.2014.12.001
- [35] Povysheva, T. V., Shmarov, M. M., Logunov, D. Y., Naroditsky, B., Shulman, I., Ogurcov, S., et al. (2017). Astrocytes mediated post-traumatic spinal cord regeneration in rats after intraspinal injection of recombinant adenoviral vectors Ad5-VEGF and Ad5-ANG. *J. Neurosurg. Spine* 27, 105-115. doi: 10.3171/2016.9.SPINE15959
- [36] Röhl, C., Lucius, R., and Sievers, J. (2007). The effect of activated microglia on astrogliosis parameters in astrocyte cultures. *Brain Res.* 1129, 43-52. doi: 10.1016/j.brainres.2006.10.057
- [37] Romero, M. I., Rangappa, N., Garry, M. G., and Smith, G. M. (2001). Functional regeneration of chronically injured sensory afferents into adult spinal cord after neurotrophin gene therapy. *J. Neurosci.* 21, 8408-8416
- [38] Saini, V., Loers, G., Kaur, G., Schachner, M., and Jakovcevski, I. (2016). Impact of neural cell adhesion molecule deletion on regeneration after mouse spinal cord injury. *Eur. J. Neurosci.* 44, 1734-1746. doi: 10.1111/ejn.13271
- [39] Schnitzer, J., Franke, W. W., and Schachner, M. (1981). Immunocytochemical demonstration of vimentin in astrocytes and ependymal cells of developing and adult mouse nervous system. *J. Cell Biol.* 90, 435-447. doi: 10.1083/jcb.90.2.435
- [40] Shcherbinin, D. N., Esmagambetov, I. B., Noskov, A. N., Selyaninov, Y. O., Tutykhina, I. L., Shmarov, M. M., et al. (2014). Protective immune response against *Bacillus anthracis* induced by intranasal introduction of a recombinant adenovirus expressing the protective antigen fused to the Fc-fragment of IgG2a. *Acta Naturae* 6, 76-84

- [41] Skorupa, A., King, M. A., Aparicio, I. M., Dussmann, H., Coughlan, K., Breen, B., et al. (2012). Motoneurons secrete angiogenin to induce RNA cleavage in astroglia. *J. Neurosci.* 32, 5024-5038. doi: 10.1523/JNEUROSCI.6366-11.2012
- [42] Tai, M.-H., Cheng, H., Wu, J.-P., Liu, Y. L., Lin, P. R., Kuo, J. S., et al. (2003). Gene transfer of glial cell line-derived neurotrophic factor promotes functional recovery following spinal cord contusion. *Exp. Neurol.* 183, 508-515. doi: 10.1016/S0014-4886(03)00130-4
- [43] Thuret, S., Moon, L. D. F., and Gage, F. H. (2006). Therapeutic interventions after spinal cord injury. *Nat. Rev. Neurosci.* 7, 628-643. doi: 10.1038/nrn1955
- [44] Tilleux, S., Berger, J., and Hermans, E. (2007). Induction of astrogliosis by activated microglia is associated with a down-regulation of metabotropic glutamate receptor 5. *J. Neuroimmunol.* 189, 23-30. doi: 10.1016/j.jneuroim.2007.06.011
- [45] van der Weerd, L., Tariq Akbar, M., Aron Badin, R., Valentim, L. M., Thomas, D. L., Wells, D. J., et al. (2010). Overexpression of heat shock protein 27 reduces cortical damage after cerebral ischemia. *J. Cereb. Blood Flow Metab.* 30, 849-856. doi: 10.1038/jcbfm.2009.249
- [46] Walthers, C. M., and Seidlits, S. K. (2015). Gene delivery strategies to promote spinal cord repair. *Biomark. Insights* 10, 11-29. doi: 10.4137/BMI.S20063
- [47] Widenfalk, J., Lipson, A., Jubran, M., Hofstetter, C., Ebendal, T., Cao, Y., et al. (2003). Vascular endothelial growth factor improves functional outcome and decreases secondary degeneration in experimental spinal cord contusion injury. *Neuroscience* 120, 951-960. doi: 10.1016/S0306-4522(03)00399-3
- [48] Xu, J.-C., Bernreuther, C., Cui, Y.-F., Jakovcevski, I., Hargus, G., Xiao, M. F., et al. (2011). Transplanted L1 expressing radial glia and astrocytes enhance recovery after spinal cord injury. *J. Neurotrauma* 28, 1921-1937. doi: 10.1089/neu.2011.1783
- [49] Yao, L., He, C., Zhao, Y., Wang, J., Tang, M., Li, J., et al. (2013). Human umbilical cord blood stem cell transplantation for the treatment of chronic spinal cord injury: Electrophysiological changes and long-term efficacy. *Neural Regen Res.* 8, 397-403. doi: 10.3969/j.issn.1673-5374.2013.05.002
- [50] Yukawa, Y., Lou, J., Fukui, N., and Lenke, L. G. (2002). Optimal treatment timing to attenuate neuronal apoptosis via Bcl-2 gene transfer in vitro and in vivo. *J. Neurotrauma* 19, 1091-1103. doi: 10.1089/089771502760341992
- [51] Zavalishin, I. A., Bochkov, N. P., Suslina, Z. A., Zakharova, M. N., Tarantul, V. Z., Naroditskiy, B. S., et al. (2008). Gene therapy of amyotrophic lateral sclerosis. *Bull. Exp. Biol. Med.* 145, 483-486
- [52] Zhang, J., Wang, H., Zhang, C., and Li, W. (2016). Intrathecal decompression versus epidural decompression in the treatment of severe spinal cord injury in rat model: a randomized, controlled preclinical research. *J. Orthop. Surg. Res.* 11:34. doi: 10.1186/s13018-016-0369-y
- [53] Zhou, Z., Peng, X., Insolera, R., Fink, D. J., and Mata, M. (2009). IL-10 promotes neuronal survival following spinal cord injury. *Exp. Neurol.* 220, 183-190. doi: 10.1016/j.expneurol.2009.08.018
- [54] Zhu, H., Poon, W., Liu, Y., Leung, G. K., Wong, Y., Feng, Y., et al. (2016). Phase III clinical trial assessing safety and efficacy of umbilical cord blood mononuclear cell transplant therapy of chronic complete spinal cord injury. *Cell Transplant.* 25, 1925-1943. doi: 10.3727/096368916X691411