

# Antimicrobial effects of sulfonyl derivative of 2(5H)-furanone against planktonic and biofilm associated methicillin-resistant and -susceptible *Staphylococcus aureus*

Sharafutdinov I., Trizna E., Baidamshina D., Ryzhikova M., Sibgatullina R., Khabibrakhmanova A., Latypova L., Kurbangalieva A., Rozhina E., Klinger-Strobel M., Fakhrullin R., Pletz M., Bogachev M., Kayumov A., Makarewicz O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017 Sharafutdinov, Trizna, Baidamshina, Ryzhikova, Sibgatullina, Khabibrakhmanova, Latypova, Kurbangalieva, Rozhina, Klinger-Strobel, Fakhrullin, Pletz, Bo gachev, Kayumov and Makarewicz. The gram-positive opportunistic bacterium *Staphylococcus aureus* is one of the most common causatives of a variety of diseases including skin and skin structure infection or nosocomial catheter-associated infections. The biofilm formation that is an important virulence factor of this microorganism renders the antibiotic therapy ineffective, because biofilm-embedded bacteria exhibit strongly increased tolerance to antimicrobials. Here, we describe a novel 3-chloro-5(S)-[(1R,2S,5R)-2-isopropyl-5-methylcyclohexyloxy]-4-[4-methylphenylsulfonyl] -2(5H)-furanone (F105), possessing a sulfonyl group and l-menthol moiety. Minimal inhibitory and bactericidal concentration values (MIC and MBC) of F105 were 10 and 40 mg/L, respectively, suggesting F105 biocidal properties. F105 exhibits pronounced activity against biofilm-embedded *S. aureus* and increases the efficacy of aminoglycosides (amikacin, gentamicin, and kanamycin) and benzalkonium chloride with fractional inhibitory concentration index values of 0.33-0.44 and 0.29, respectively, suggesting an alternative external treatment option, e.g., for wound infections. Moreover, low concentrations (0.5-1.3 mg/L) of F105 reduced the MICs of these antimicrobials twofold. By using confocal laser scanning microscopy and CFU counting, we show explicitly that F105 also restores the antimicrobial activity of gentamicin and ampicillin against *S. aureus* biofilms by several orders of magnitude. Biofilm structures were not destroyed but sterilized, with embedded cells being almost completely killed at twofold MBC. While F105 is quite toxic (CC 50 /MBC ratio 0.2), our data suggest that the F105 chemotype might be a promising starting point for the development of complex topical agents for combined anti-staphylococcal biofilm-therapies restoring the efficacy of some antibiotics against difficult to treat *S. aureus* biofilm.

<http://dx.doi.org/10.3389/fmicb.2017.02246>

---

## Keywords

2(5H)-furanones, Antimicrobials synergism, Biofilm, MRSA, *Staphylococcus aureus*, Sulfones

## References

- [1] Akimitsu, N., Hamamoto, H., Inoue, R., Shoji, M., Akamine, A., Takemori, K., et al. (1999). Increase in resistance of methicillin-resistant *Staphylococcus aureus* to beta-lactams caused by mutations conferring resistance to benzalkonium chloride, a disinfectant widely used in hospitals. *Antimicrob. Agents Chemother.* 43, 3042-3043
- [2] Atshan, S. S., Shamsudin, M. N., Sekawi, Z., Lung, L. T. T., Barantalab, F., Liew, Y. K., et al. (2015). Comparative proteomic analysis of extracellular proteins expressed by various clonal types of *Staphylococcus aureus* and during planktonic growth and biofilm development. *Front. Microbiol.* 6:524. doi: 10.3389/fmicb.2015.00524
- [3] Baidamshina, D. R., Trizna, E. Y., Holyavka, M. G., Bogachev, M. I., Artyukhov, V. G., Akhatova, F. S., et al. (2017). Targeting microbial biofilms using Ficin, a nonspecific plant protease. *Sci. Rep.* 7:46068. doi: 10.1038/srep46068
- [4] Bassetti, M., Carnelutti, A., and Righi, E. (2017). The role of methicillin-resistant *Staphylococcus aureus* in skin and soft tissue infections. *Curr. Opin. Infect. Dis.* 30, 150-157. doi: 10.1097/qco.0000000000000353
- [5] Bernstein, I. L. (2000). Is the use of benzalkonium chloride as a preservative for nasal formulations a safety concern? A cautionary note based on compromised mucociliary transport. *J. Allergy Clin. Immunol.* 105, 39-44. doi: 10.1016/s0091-6749(00)90175-1
- [6] Bortolin, M., Bidossi, A., De Vecchi, E., Avvenente, M., and Drago, L. (2017). In vitro antimicrobial activity of chlorquinadol against microorganisms responsible for skin and soft tissue infections: comparative evaluation with gentamicin and fusidic acid. *Front. Microbiol.* 8:10. doi: 10.3389/fmicb.2017.01039
- [7] Brackman, G., and Coenye, T. (2015). Quorum sensing inhibitors as anti-biofilm agents. *Curr. Pharm. Des.* 21, 5-11. doi: 10.2174/1381612820666140905114627
- [8] Bremer, F., Grade, S., Kohorst, P., and Stiesch, M. (2011). In vivo biofilm formation on different dental ceramics. *Quintessence Int.* 42, 565-574
- [9] Conlon, B. P. (2014). *Staphylococcus aureus* chronic and relapsing infections: evidence of a role for persister cells an investigation of persister cells, their formation and their role in *S. aureus* disease. *Bioessays* 36, 991-996. doi: 10.1002/bies.201400080
- [10] Cosgrove, S. E., and Fowler, V. G. Jr. (2008). Management of methicillin-resistant *Staphylococcus aureus* bacteremia. *Clin. Infect. Dis.* 46, S386-S393. doi: 10.1086/533595
- [11] Cosgrove, S. E., Kaye, K. S., Eliopoulos, G. M., and Carmeli, Y. (2002). Health and economic outcomes of the emergence of third-generation cephalosporin resistance in *Enterobacter* species. *Arch. Intern. Med.* 162, 185-190. doi: 10.1001/archinte.162.2.185
- [12] Damour, O., Hua, S. Z., Lasne, F., Villain, M., Rousselle, P., and Collombel, C. (1992). Cytotoxicity evaluation of antibiotics on cultured human fibroblasts and keratinocytes. *Burns* 18, 479-485. doi: 10.1016/0305-4179(92)90180-3
- [13] de Jonge, B. L., and Tomasz, A. (1993). Abnormal peptidoglycan produced in a methicillin-resistant strain of *Staphylococcus aureus* grown in the presence of methicillin: functional role for penicillin-binding protein 2A in cell wall synthesis. *Antimicrob. Agents Chemother.* 37, 342-346. doi: 10.1128/AAC.37.2.342
- [14] den Hollander, J. G., Mouton, J. W., and Verbrugh, H. A. (1998). Use of pharmacodynamic parameters to predict efficacy of combination therapy by using fractional inhibitory concentration kinetics. *Antimicrob. Agents Chemother.* 42, 744-748
- [15] Doll, K., Jongsthaphongpun, K. L., Stumpp, N. S., Winkel, A., and Stiesch, M. (2016). Quantifying implant-associated biofilms: comparison of microscopic, microbiologic and biochemical methods. *J. Microbiol. Methods* 130, 61-68. doi: 10.1016/j.mimet.2016.07.016
- [16] Donlan, R. M. (2002). Biofilms: microbial life on surfaces. *Emerg. Infect. Dis.* 8, 881-890. doi: 10.3201/eid0809.020063
- [17] Donlan, R. M., and Costerton, J. W. (2002). Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin. Microbiol. Rev.* 15, 167-193. doi: 10.1128/cmr.15.2.167-193.2002
- [18] Eliopoulos, G., and Moellering, R. C. Jr. (1996). 'Antimicrobial combinations,' in *Antibiotics in Laboratory Medicine*, 4th Edn, ed. V. Lorian (Baltimore, MD: The Williams & Wilkins Co), 330-396
- [19] European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). (2000). Terminology relating to methods for the determination of susceptibility of bacteria to antimicrobial agents. *Clin. Microbiol. Infect.* 6, 503-508. doi: 10.1046/j.1469-0691.2000.00149.x
- [20] Fagerlund, A., Langsrud, S., Heir, E., Mikkelsen, M. I., and Moretro, T. (2016). Biofilm matrix composition affects the susceptibility of food associated staphylococci to cleaning and disinfection agents. *Front. Microbiol.* 7:856. doi: 10.3389/fmicb.2016.00856
- [21] Fedorova, K. P., Scharafutdinov, I. S., Turbina, E. Y., Bogachev, M. I., Ilinskaja, O. N., and Kayumov, A. R. (2013). C-terminus of transcription factor TnrA from *Bacillus subtilis* controls DNA-binding domain activity but is not required for dimerization. *Mol. Biol.* 47, 293-298. doi: 10.1134/S0026893313020052
- [22] Fenske, D., and Merzweiler, K. (1989). Synthesis of a new chiral phosphine ligand. *J. Chem. Sci.* 44, 879-883

- [23] Forrest, G. N., and Tamura, K. (2010). Rifampin combination therapy for nonmycobacterial infections. *Clin. Microbiol. Rev.* 23, 14-34. doi: 10.1128/cmrr.00034-09
- [24] Fromm-Dornieden, C., Rembe, J. D., Schafer, N., Bohm, J., and Stuermer, E. K. (2015). Cetylpyridinium chloride and miramistin as antiseptic substances in chronic wound management-prospects and limitations. *J. Med. Microbiol.* 64, 407-414. doi: 10.1099/jmm.0.000034
- [25] Garcia, L. G., Lemaire, S., Kahl, B. C., Becker, K., Proctor, R. A., Denis, O., et al. (2013). Antibiotic activity against small-colony variants of *Staphylococcus aureus*: review of in vitro, animal and clinical data. *J. Antimicrob. Chemother.* 68, 1455-1464. doi: 10.1093/jac/dkt072
- [26] Heck, R., and Stuetz, A. (1988). Antimycotic-6-phenyl-2-hexen-4-ynamines. U.S. Patent No. EP 0254677 A1
- [27] Hentzer, M., Riedel, K., Rasmussen, T. B., Heydorn, A., Andersen, J. B., Parsek, M. R., et al. (2002). Inhibition of quorum sensing in *Pseudomonas aeruginosa* biofilm bacteria by a halogenated furanone compound. *Microbiology* 148, 87-102. doi: 10.1099/00221287-148-1-87
- [28] Hentzer, M., Wu, H., Andersen, J. B., Riedel, K., Rasmussen, T. B., Bagge, N., et al. (2003). Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* 22, 3803-3815. doi: 10.1093/emboj/cdg366
- [29] Herigstad, B., Hamilton, M., and Heersink, J. (2001). How to optimize the drop plate method for enumerating bacteria. *J. Microbiol. Methods* 44, 121-129. doi: 10.1016/s0167-7012(00)00241-4
- [30] Hodille, E., Rose, W., Diep, B. A., Goutelle, S., Lina, G., and Dumitrescu, O. (2017). The role of antibiotics in modulating virulence in *Staphylococcus aureus*. *Clin. Microbiol. Rev.* 30, 887-917. doi: 10.1128/CMR.00120-16
- [31] Kali, A., Bhuvaneshwar, D., Charles, P. M., and Seetha, K. S. (2016). Antibacterial synergy of curcumin with antibiotics against biofilm producing clinical bacterial isolates. *J. Basic Clin. Pharm.* 7, 93-96. doi: 10.4103/0976-0105.183265
- [32] Kalita, S., Devi, B., Kandimalla, R., Sharma, K. K., Sharma, A., Kalita, K., et al. (2015). Chloramphenicol encapsulated in poly-epsilon-caprolactone-pluronic composite: nanoparticles for treatment of MRSA-infected burn wounds. *Int. J. Nanomed.* 10, 2971-2984. doi: 10.2147/ijn.s75023
- [33] Kamatou, G. P., Vermaak, I., Viljoen, A. M., and Lawrence, B. M. (2013). Menthol: a simple monoterpenoid with remarkable biological properties. *Phytochemistry* 96, 15-25. doi: 10.1016/j.phytochem.2013.08.005
- [34] Kayumov, A. R., Khakimullina, E. N., Sharafutdinov, I. S., Trizna, E. Y., Latypova, L. Z., Hoang Thi, L., et al. (2015a). Inhibition of biofilm formation in *Bacillus subtilis* by new halogenated furanones. *J. Antibiotics* 68, 297-301. doi: 10.1038/ja.2014.143
- [35] Kayumov, A. R., Nureeva, A. A., Trizna, E. Y., Gazizova, G. R., Bogachev, M. I., Shtyrlin, N. V., et al. (2015b). New derivatives of pyridoxine exhibit high antibacterial activity against biofilm-embedded staphylococcus cells. *Biomed Res. Int.* 2015:890968. doi: 10.1155/2015/890968
- [36] Kifer, D., Muzinic, V., and Klaric, M. S. (2016). Antimicrobial potency of single and combined mupirocin and monoterpenes, thymol, menthol and 1,8-cineole against *Staphylococcus aureus* planktonic and biofilm growth. *J. Antibiot.* 69, 689-696. doi: 10.1038/ja.2016.10
- [37] Kitty, H., Samuel, K., Daniel, C., Renxun, C., Mark, W., and Naresh, K. (2015). 'Development of fimbriolides, halogenated furanones and their derivatives as antimicrobial agents,' in *Antibacterial Surfaces*, ed. R. C. Elena Ivanova (Cham: Springer International Publishing), 149-170
- [38] Klinger-Strobel, M., Suesse, H., Fischer, D., Pletz, M. W., and Makarewicz, O. (2016). A novel computerized cell count algorithm for biofilm analysis. *PLOS ONE* 11:e0154937. doi: 10.1371/journal.pone.0154937
- [39] Kudryavtsev, K. V., Bentley, M. L., and McCafferty, D. G. (2009). Probing of the cis-5-phenyl proline scaffold as a platform for the synthesis of mechanism-based inhibitors of the *Staphylococcus aureus* sortase SrtA isoform. *Bioorg. Med. Chem.* 17, 2886-2893. doi: 10.1016/j.bmc.2009.02.008
- [40] Kuehl, R., Al-Bataineh, S., Gordon, O., Luginbuehl, R., Otto, M., Textor, M., et al. (2009). Furanone at subinhibitory concentrations enhances staphylococcal biofilm formation by luxS repression. *Antimicrob. Agents Chemother.* 53, 4159-4166. doi: 10.1128/aac.01704-08
- [41] Kurbangalieva, A. R., Devyatova, N. F., Bogdanov, A. V., Berdnikov, E. A., Mannafov, T. G., Krivolapov, D. B., et al. (2007). Synthesis of novel arylthio derivatives of mucochloric acid. *Phosphorus Sulfur Silicon Relat. Elem.* 182, 607-630. doi: 10.1080/10426500601015989
- [42] Lattmann, E., Dunn, S., Niemann, S., and Sattayasai, N. (2005). Synthesis and antibacterial activities of 5-hydroxy-4-amino-2(5H)-furanones. *Bioorg. Med. Chem. Lett.* 15, 919-921. doi: 10.1016/j.bmcl.2004.12.051
- [43] Latypova, L. Z., Saigitbatalova, E. S., Chulakova, D. R., Lodochnikova, O. A., Kurbangalieva, A. R., Berdnikov, E. A., et al. (2014). Sulfides, sulfones, and sulfoxides of the furan-2(5H)-one series. synthesis and structure. *Russ. J. Organ. Chem.* 50, 521-534. doi: 10.1134/s1070428014040149
- [44] Leclercq, R., Canton, R., Brown, D. F., Giske, C. G., Heisig, P., MacGowan, A. P., et al. (2013). EUCAST expert rules in antimicrobial susceptibility testing. *Clin. Microbiol. Infect.* 19, 141-160. doi: 10.1111/j.1469-0691.2011.03703.x
- [45] Lewis, K. (2001). Riddle of biofilm resistance. *Antimicrob. Agents Chemother.* 45, 999-1007. doi: 10.1128/AAC.45.4.999-1007.2001

- [46] Li, S. S., Skov, R. L., Han, X., Larsen, A. R., Larsen, J., Sorum, M., et al. (2011). Novel types of staphylococcal cassette chromosome *mec* elements identified in clonal complex 398 methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob. Agents Chemother.* 55, 3046-3050. doi: 10.1128/aac.01475-10
- [47] Lonn-Stensrud, J., Landin, M. A., Benneche, T., Petersen, F. C., and Scheie, A. A. (2009). Furanones, potential agents for preventing *Staphylococcus epidermidis* biofilm infections? *J. Antimicrob. Chemother.* 63, 309-316. doi: 10.1093/jac/dkn501
- [48] Low, E., Kim, B., Francavilla, C., Shiao, T. P., Turtle, E. D., O'Mahony, D. J. R., et al. (2011). Structure stability/activity relationships of sulfone stabilized N,N-dichloroamines. *Bioorg. Med. Chem. Lett.* 21, 3682-3685. doi: 10.1016/j.bmcl.2011.04.084
- [49] McCann, J., and Ames, B. N. (1976). A simple method for detecting environmental carcinogens as mutagens. *Ann. N. Y. Acad. Sci.* 271, 5-13. doi: 10.1111/j.1749-6632.1976.tb23086.x
- [50] McCarthy, H., Rudkin, J. K., Black, N. S., Gallagher, L., O'Neill, E., and O'Gara, J. P. (2015). Methicillin resistance and the biofilm phenotype in *Staphylococcus aureus*. *Front. Cell. Infect. Microbiol.* 5:1. doi: 10.3389/fcimb.2015.00001
- [51] Meadows, D. C., and Gervay-Hague, J. (2006). Vinyl sulfones: synthetic preparations and medicinal chemistry applications. *Med. Res. Rev.* 26, 793-814. doi: 10.1002/med.20074
- [52] Merritt, J. H., Kadouri, D. E., and O'Toole, G. A. (2005). Growing and analyzing static biofilms. *Curr. Protoc. Microbiol.* Chapter 1, Unit 1B. 1. doi: 10.1002/9780471729259.mc01b01s00
- [53] Mohan, N. R., Sreenivasa, S., Manojkumar, K. E., Rao, T. M. C., Thippeswamy, B. S., and Suchetan, P. A. (2014). Synthesis, antibacterial, anthelmintic and anti-inflammatory studies of novel methylpyrimidine sulfonyl piperazine derivatives. *J. Braz. Chem. Soc.* 25, 1012-1020. doi: 10.5935/0103-5053.20140073
- [54] Naicker, P. R., Karayem, K., Hoek, K. G. P., Harvey, J., and Wasserman, E. (2016). Biofilm formation in invasive *Staphylococcus aureus* isolates is associated with the clonal lineage. *Microb. Pathog.* 90, 41-49. doi: 10.1016/j.micpath.2015.10.023
- [55] Narishetty, S. T., and Panchagnula, R. (2005). Effect of L-menthol and 1,8-cineole on phase behavior and molecular organization of SC lipids and skin permeation of zidovudine. *J. Control. Release* 102, 59-70. doi: 10.1016/j.jconrel.2004.09.016
- [56] Odds, F. C. (2003). Synergy, antagonism, and what the chequerboard puts between them. *J. Antimicrob. Chemother.* 52:1. doi: 10.1093/jac/dkg301
- [57] OECD (1997). Test No. 471: Bacterial Reverse Mutation Test. Paris: OECD
- [58] Parra-Ruiz, J., Vidaillac, C., Rose, W. E., and Rybak, M. J. (2010). Activities of high-dose daptomycin, vancomycin, and moxifloxacin alone or in combination with clarithromycin or rifampin in a novel in vitro model of *Staphylococcus aureus* biofilm. *Antimicrob. Agents Chemother.* 54, 4329-4334. doi: 10.1128/aac.00455-10
- [59] Peeters, E., Nelis, H. J., and Coenye, T. (2008). Comparison of multiple methods for quantification of microbial biofilms grown in microtiter plates. *J. Microbiol. Methods* 72, 157-165. doi: 10.1016/j.mimet.2007.11.010
- [60] Percival, S. L., Finnegan, S., Donelli, G., Vuotto, C., Rimmer, S., and Lipsky, B. A. (2016). Antiseptics for treating infected wounds: efficacy on biofilms and effect of pH. *Crit. Rev. Microbiol.* 42, 293-309. doi: 10.3109/1040841x.2014.940495
- [61] Ren, D. C., Bedzyk, L. A., Setlow, P., England, D. F., Kjelleberg, S., Thomas, S. M., et al. (2004). Differential gene expression to investigate the effect of (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone on *Bacillus subtilis*. *Appl. Environ. Microbiol.* 70, 4941-4949. doi: 10.1128/aem.70.8.4941-4949.2004
- [62] Ren, D. C., Sims, J. J., and Wood, T. K. (2001). Inhibition of biofilm formation and swarming of *Escherichia coli* by (5Z)-4-bromo-5(bromomethylene)-3-butyl-2(5H)-furanone. *Environ. Microbiol.* 3, 731-736. doi: 10.1046/j.1462-2920.2001.00249.x
- [63] Rogers, S. A., Whitehead, D. C., Mullikin, T., and Melander, C. (2010). Synthesis and bacterial biofilm inhibition studies of ethyl N-(2-phenethyl) carbamate derivatives. *Organ. Biomol. Chem.* 8, 3857-3859. doi: 10.1039/c0ob00063a
- [64] Roy, R., Tiwari, M., Donelli, G., and Tiwari, V. (2017). Strategies for combating bacterial biofilms: a focus on anti-biofilm agents and their mechanisms of action. *Virulence* doi: 10.1080/21505594.2017.1313372 [Epub ahead of print]
- [65] Sanchez-Vizcute, P., Orgaz, B., Aymerich, S., Le Coq, D., and Briandet, R. (2015). Pathogens protection against the action of disinfectants in multispecies biofilms. *Front. Microbiol.* 6:705. doi: 10.3389/fmich.2015.00705
- [66] Sharafutdinov, I., Shigapova, Z., Baltin, M., Akhmetov, N., Bogachev, M., and Kayumov, A. (2016). HtrA protease from *Bacillus subtilis* suppresses the bacterial fouling of the rat skin injuries. *Bionanoscience* 6, 564-567. doi: 10.1007/s12668-016-0281-2
- [67] Siala, W., Mingeot-Leclercq, M. P., Tulkens, P. M., Hallin, M., Denis, O., and Van Bambeke, F. (2014). Comparison of the antibiotic activities of daptomycin, vancomycin, and the investigational fluoroquinolone delafloxacin against biofilms from *Staphylococcus aureus* clinical isolates. *Antimicrob. Agents Chemother.* 58, 6385-6397. doi: 10.1128/AAC.03482-14

- [68] Singh, R., Ray, P., Das, A., and Sharma, M. (2010). Penetration of antibiotics through *Staphylococcus aureus* and *Staphylococcus epidermidis* biofilms. *J. Antimicrob. Chemother.* 65, 1955-1958. doi: 10.1093/jac/dkq257
- [69] Stein, C., Makarewicz, O., Forstner, C., Weis, S., Hagel, S., Loffler, B., et al. (2016). Should daptomycin-rifampin combinations for MSSA/MRSA isolates be avoided because of antagonism? *Infection* 44, 499-504. doi: 10.1007/s15010-016-0874-2
- [70] Thirukovela, N. S., Kankala, S., Kankala, R. K., Paidakula, S., Gangula, M. R., Vasam, C. S., et al. (2017). Regioselective synthesis of some new 1,4-disubstituted sulfonyl-1,2,3-triazoles and their antibacterial activity studies. *Med. Chem. Res.* 26, 2190-2195. doi: 10.1007/s00044-017-1926-6
- [71] Trizna, E., Latypova, L., Kurbangalieva, A., Bogachev, M., Kayumov, A., et al. (2016). 2(5H)-Furanone derivatives as inhibitors of staphylococcal biofilms. *BioNanoScience* 6, 423-426. doi: 10.1007/s12668-01-0258-1
- [72] Trizna, E. Y., Khakimullina, E. N., Latypova, L. Z., Kurbangalieva, A. R., Sharafutdinov, I. S., Evtyugin, V. G., et al. (2015). Thio derivatives of 2(5H)-furanone as inhibitors against *Bacillus subtilis* biofilms. *Acta Naturae* 7, 102-107
- [73] Yujie, L., Geng, X., Huang, Y. C., Li, Y., Yang, K. Y., Ye, L. H., et al. (2013). The Effect of brominated furanones on the formation of *Staphylococcus aureus* biofilm on PVC. *Cell Biochem. Biophys.* 67, 1501-1505. doi: 10.1007/s12013-013-9652-2