

Effect of membrane-active microbial autoregulators on the growth of cultured ras-transformed fibroblasts

Il'inskaya O., Kolpakov A., Mulyukin A., Dreyer F., El'-Registan G.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Differential effects on the proliferation of individual vs. combined administration of high- and low-molecular-weight microbial autoregulators (extracellular RNase from *Bacillus intermedius* and anabiosis-inducing factor d 1) are reported for the first time for cultured cells of higher eukaryotes. Proliferation of ras-transformed mouse fibroblasts was affected by both autoregulators dose-dependently. The cytotoxic activity of individual regulators was directly related to their concentration. Unlike RNase, factor d 1 (which functions as a chemical chaperone) exerted reversible effects. Studies of the effects of combined administration of the autoregulators demonstrated that pretreatment of the cells with low-dose d 1 decreased the toxicity of RNase. Higher doses of d 1 were required to attenuate the effects of toxic agents with more pronounced membrane tropism. The results obtained suggest that a universal system regulating the physiological activity of cells is operative in taxonomically remote organisms. The operation of the system is based on sequential changes in the structural organization and function of subcellular structures induced by low- and high-molecular-weight autoregulators.
