

## Length and time scales of entanglement and confinement effects constraining polymer chain dynamics

Kimmich R., Fatkullin N.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

With characteristic time constants for polymer dynamics, namely  $\tau_s$  (the segment fluctuation time),  $\tau_e$  (the entanglement time), and  $\tau_R$  (the longest Rouse relaxation time), the time scales of particular interest (i)  $\tau < \tau_{ss}$  (ii)  $\tau_S < t < \tau_e$ , and (iii)  $\tau_e < t < \tau_R$  will be discussed and compared with experimental data. These ranges correspond to the chain-mode length scales (i)  $l < b$ , (ii)  $b < l < d^2/b$ , and (iii)  $d^2/b < l < L$ , where  $b$  is the statistical segment length,  $d$  is the dimension of constraints by entanglements and/or confinement, and  $L$  is the chain contour length. Based on Langevin-type equations-of-motion coarse-grained predictions for the mean-squared segment displacement and the spin-lattice relaxation dispersion will be outlined for the scenarios "freely-draining", "entangled", and "confined". In the discussion we will juxtapose "local" versus "global" dynamics on the one hand, and "bulk" versus "confined" systems on the other. © 2010 Materials Research Society.

---