

Complexes of N-(thio)phosphorylthioureas AdNHC(S)NHP(X)(OiPr)₂ (X = O, S) with cobalt(II), zinc(II), and cadmium(II). Crystal structure of Co[AdNHC(S)NP(S)(OiPr)₂]₂

Safin D., Bolte M., Babashkina M.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Reaction of the potassium salts of N-(thio)phosphorylated thioureas of common formula AdNHC(S)N(H)P(X)(OiPr)₂ (Ad = adamantyl; X = O, HL I ; X = S, HL II) with Co(II), Zn(II), and Cd(II) cations in aqueous EtOH leads to M(L I,II -X, S)₂ chelate complexes. The Cd(II) complex Cd(L I)₂ could not be isolated under analogous conditions because of its hydrolytic lability. The structure of the resulting compounds was studied by means of spectroscopy and microanalysis; in addition, the molecular structure of the complex Co(L II)₂ was elucidated by single crystal X-ray diffraction analysis. The cobalt atom is in a tetrahedral S₄ environment formed by the C=S and P=S sulfur atoms of two deprotonated ligands L II . The magnetic properties of Co(L I,II)₂ were investigated and the photoluminescent properties of the complexes are also reported. © 2008 Springer Science+Business Media B.V.

<http://dx.doi.org/10.1007/s11243-008-9157-9>
