

Coordination mode of the nickel(II) cation with N-diisopropoxyphosphinyl-p- bromothiobenzamide

Safin D., Sokolov F., Baranov S., Szyrwił Ł., Babashkina M., Shakirova E., Hahn F., Kozłowski H.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Reaction of the potassium salt of N-diisopropoxyphosphinyl-p- bromothiobenzamide $p\text{-BrC}_6\text{H}_4\text{C(S)NHP(O)(OiPr)}_2$ (HL) with $\text{Ni(NO}_3)_2$ in aqueous EtOH leads to complex of formula $[\text{Ni}(\text{HL-O})_2(\text{L-O,S})_2]$ (1). The structure of 1 was investigated by single crystal X-ray diffraction analysis, IR, ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy, MALDI and microanalysis. The nickel(II) ion in 1 has a tetragonal-bipyramidal environment, $(\text{O}_{\text{ax}})_2(\text{O}_{\text{eq}})_2(\text{Seq})_2$, with two neutral ligand molecules coordinated in axial positions through the oxygen atoms of the P=O groups. The equatorial plane of bipyramide is formed by two anionic ligands involving 1,5-O,S-coordination mode. The chelating ligands are bound in trans configuration. © 2008 Wiley-VCH Verlag GmbH & Co. KGaA.

<http://dx.doi.org/10.1002/zaac.200700502>

Keywords

Amidophosphate, Chelates, Crystal structure, Nickel