



Contents lists available at ScienceDirect

J. Chem. Thermodynamics

journal homepage: www.elsevier.com/locate/jct

Thermodynamic properties of methylprednisolone aceponate



A.V. Knyazev^{a,*}, V.N. Emel'yanenko^{b,c,*}, N.N. Smirnova^a, O.V. Stepanova^d, A.S. Shipilova^a,
A.V. Markin^a, Ya.S. Samosudova^a, E.V. Gusarova^a, S.S. Knyazeva^a, S.P. Verevkin^{b,*}

^aLobachevsky State University of Nizhni Novgorod, Gagarin Prospekt 23/2, 603950 Nizhni Novgorod, Russia

^bDepartment of Physical Chemistry, University of Rostock, Dr-Lorenz-Weg 1, D-18059 Rostock, Germany

^cDepartment of Physical Chemistry, Kazan Federal University, Kremlevskaya str. 18, 420008 Kazan, Russia

^dOpen Joint Stock Company Chemico-Pharmaceutical Works Akrikhin, Kosmodamianskaya Naberezhnaya, 52/5, 115054 Moscow, Russia

ARTICLE INFO

Article history:

Received 29 July 2016

Received in revised form 12 August 2016

Accepted 16 August 2016

Available online 17 August 2016

Keywords:

Steroid hormone

Methylprednisolone aceponate

Adiabatic vacuum calorimetry

Heat capacity

Combustion calorimetry

Thermodynamic functions

ABSTRACT

In the present work, temperature dependence of heat capacity of methylprednisolone aceponate has been measured for the first time over the range from 7 K to 346 K by precision adiabatic vacuum calorimetry. Based on the experimental results, the thermodynamic functions of the methylprednisolone aceponate, namely, the heat capacity, enthalpy $H^\circ(T) - H^\circ(0)$, entropy $S^\circ(T) - S^\circ(0)$ and Gibbs function $G^\circ(T) - H^\circ(0)$ have been determined for the range from $T/K \rightarrow 0 - 333$. The enthalpy of combustion (-14304.4 ± 9.1) $\text{kJ}\cdot\text{mol}^{-1}$ of the methylprednisolone aceponate was determined for the first time using high-precision combustion calorimeter. The standard molar enthalpy of formation in the crystalline state (-1465.3 ± 9.8) $\text{kJ}\cdot\text{mol}^{-1}$ of compound at 298.15 K was derived from the combustion experiments. Using a combination of the adiabatic and combustion calorimetry results the thermodynamic functions of formation of the folic acid dihydrate at $T = 298.15$ K and $p = 0.1$ MPa have been calculated.

© 2016 Elsevier Ltd.

1. Introduction

Methylprednisolone aceponate (CAS: 86401-95-8) is a non-halogenated diester of 6 α -methylprednisolone. The introduction of two ester groups results in a molecule with increased lipophilicity and enhanced penetration into skin. Estimations of anti-inflammatory activity in humans suggest that it is a protein glucocorticosteroid. In humans its atrophogenic potential, as assessed by skin thinning and telangiectasia, appears lower than, or similar to, that of other agents belonging to this class such as betamethasone valerate and mometasone furoate [1].

This work is a continuation of systematic studies of bioactive compounds. Earlier in the articles [2–6], we have investigated the thermodynamic properties of vitamins and hydrocortisone acetate. It should be noted that the temperature dependence of the heat capacity of the steroid hormones has not been studied previously. The goals of this work include calorimetric determination of the standard thermodynamic functions of the methylprednisolone aceponate (steroid hormone) with the purpose of describing biochemical and industrial processes with its participation.

* Corresponding authors at: Department of Physical Chemistry, University of Rostock, Dr-Lorenz-Weg 1, D-18059 Rostock, Germany.

E-mail addresses: knyazevav@gmail.com (A.V. Knyazev), vladimir.yemelyanenko@uni-rostock.de (V.N. Emel'yanenko).

2. Experimental

2.1. Sample

Methylprednisolone aceponate was purchased from Fluka. For phase identification, an X-ray diffraction pattern of the methylprednisolone aceponate sample was recorded on a Shimadzu X-ray diffractometer XRD-6000 (CuK α radiation, geometry θ -2 θ) in the 2 θ range from 5° to 60° with scan increment of 0.02°. The absence of water in methylprednisolone aceponate was determined by Karl Fischer titration. The X-ray data, certificate (purity control performed by mass spectrometry) and Karl Fischer titration led us to conclude that the methylprednisolone aceponate sample studied (the content of impurities 0.1 wt%) was an individual crystalline compound (trigonal modification [7]).

2.2. Apparatus and measurement procedure

To measure the heat capacity C_p° of the tested substance in the range from 7 K to 346 K, a BKT-3.0 automatic precision adiabatic vacuum calorimeter with discrete heating was used. The calorimeter design and the operation procedure were described earlier [8]. The calorimeter was tested by measuring the heat capacity of high-purity copper and reference samples of synthetic corundum and K-2 benzoic acid. The analysis of the results showed that