

Mechanisms for nonmitotic activation of Aurora-A at cilia

Korobeynikov V., Deneka A., Golemis E.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 The Author(s). Overexpression of the Aurora kinase A (AURKA) is oncogenic in many tumors. Many studies of AURKA have focused on activities of this kinase in mitosis, and elucidated the mechanisms by which AURKA activity is induced at the G2/M boundary through interactions with proteins such as TPX2 and NEDD9. These studies have informed the development of small molecule inhibitors of AURKA, of which a number are currently under preclinical and clinical assessment. While the first activities defined for AURKA were its control of centrosomal maturation and organization of the mitotic spindle, an increasing number of studies over the past decade have recognized a separate biological function of AURKA, in controlling disassembly of the primary cilium, a small organelle protruding from the cell surface that serves as a signaling platform. Importantly, these activities require activation of AURKA in early G1, and the mechanisms of activation are much less well defined than those in mitosis. A better understanding of the control of AURKA activity and the role of AURKA at cilia are both important in optimizing the efficacy and interpreting potential downstream consequences of AURKA inhibitors in the clinic. We here provide a current overview of proteins and mechanisms that have been defined as activating AURKA in G1, based on the study of ciliary disassembly.

<http://dx.doi.org/10.1042/BST20160142>

References

- [1] Glover, D. M., Leibowitz, M. H., McLean, D. A. and Parry, H. (1995) Mutations in aurora prevent centrosome separation leading to the formation of monopolar spindles. *Cell* 81, 95-105 doi:10.1016/0092-8674(95)90374-7
- [2] Carmena, M. and Earnshaw, W. C. (2003) The cellular geography of aurora kinases. *Nat. Rev. Mol. Cell. Biol.* 4, 842-854 doi:10.1038/nrm1245
- [3] Brown, J. R., Koretke, K. K., Birkeland, M. L., Sanseau, P. and Patrick, D. R. (2004) Evolutionary relationships of Aurora kinases: implications for model organism studies and the development of anti-cancer drugs. *BMC Evol. Biol.* 4, 39 doi:10.1186/1471-2148-4-39
- [4] Bischoff, J. R., Anderson, L., Zhu, Y., Mossie, K., Ng, L., Souza, B. et al. (1998) A homologue of *Drosophila* aurora kinase is oncogenic and amplified in human colorectal cancers. *EMBO J.* 17, 3052-3065 doi:10.1093/emboj/17.11.3052
- [5] Li, Z. and Rana, T. M. (2012) A kinase inhibitor screen identifies small-molecule enhancers of reprogramming and iPS cell generation. *Nat. Commun.* 3, 1085 doi:10.1038/ncomms2059
- [6] Lee, D.-F., Su, J., Ang, Y.-S., Carvajal-Vergara, X., Mulero-Navarro, S., Pereira, C. F. et al. (2012) Regulation of embryonic and induced pluripotency by aurora kinase-p53 signaling. *Cell Stem Cell* 11, 179-194 doi:10.1016/j.stem.2012.05.020
- [7] Ochi, T., Fujiwara, H. and Yasukawa, M. (2009) Aurora-A kinase: a novel target both for cellular immunotherapy and molecular target therapy against human leukemia. *Expert Opin. Ther. Targets* 13, 1399-1410 doi:10.1517/14728220903307483

- [8] Park, H.-S., Park, W. S., Bondaruk, J., Tanaka, N., Katayama, H., Lee, S. et al. (2008) Quantitation of Aurora kinase A gene copy number in urine sediments and bladder cancer detection. *J. Natl Cancer Inst.* 100, 1401-1411 doi:10.1093/jnci/djn304
- [9] Tanaka, T., Kimura, M., Matsunaga, K., Fukada, D., Mori, H. and Okano, Y. (1999) Centrosomal kinase AIK1 is overexpressed in invasive ductal carcinoma of the breast. *Cancer Res.* 59, 2041-2044 PMID:10232583
- [10] Watanabe, T., Imoto, I., Katahira, T., Hirasawa, A., Ishiwata, I., Emi, M. et al. (2002) Differentially regulated genes as putative targets of amplifications at 20q in ovarian cancers. *Jpn J. Cancer Res.* 93, 1114-1122 doi:10.1111/j.1349-7006.2002.tb01213.x
- [11] Li, D., Zhu, J., Firozi, P. F., Abbruzzese, J. L., Evans, D. B., Cleary, K. et al. (2003) Overexpression of oncogenic STK15/BTAK/Aurora A kinase in human pancreatic cancer. *Clin. Cancer Res.* 9, 991-997 PMID:12631597
- [12] Kamada, K., Yamada, Y., Hirao, T., Fujimoto, H., Takahama, Y., Ueno, M. et al. (2004) Amplification/overexpression of Aurora-A in human gastric carcinoma: potential role in differentiated type gastric carcinogenesis. *Oncol. Rep.* 12, 593-599 PMID:15289843
- [13] Rojanala, S., Han, H., Muñoz, R. M., Browne, W., Nagle, R., Von Hoff, D. D. et al. (2004) The mitotic serine threonine kinase, Aurora-2, is a potential target for drug development in human pancreatic cancer. *Mol. Cancer Ther.* 3, 451-457 PMID:15078988
- [14] Cammareri, P., Scopelliti, A., Todaro, M., Eterno, V., Francescangeli, F., Moyer, M. P. et al. (2010) Aurora-A is essential for the tumorigenic capacity and chemoresistance of colorectal cancer stem cells. *Cancer Res.* 70, 4655-4665 doi:10.1158/0008-5472.CAN-09-3953
- [15] Lee, H. H., Zhu, Y., Govindasamy, K. M. and Gopalan, G. (2008) Downregulation of Aurora-A overrides estrogen-mediated growth and chemoresistance in breast cancer cells. *Endocr. Relat. Cancer* 15, 765-775 doi:10.1677/ERC-07-0213
- [16] Wang, X., Zhou, Y.-X., Qiao, W., Tominaga, Y., Ouchi, M., Ouchi, T. et al. (2006) Overexpression of aurora kinase A in mouse mammary epithelium induces genetic instability preceding mammary tumor formation. *Oncogene* 25, 7148-7158 doi:10.1038/sj.onc.1209707
- [17] Zhou, H., Kuang, J., Zhong, L., Kuo, W.-L., Gray, J., Sahin, A. et al. (1998) Tumour amplified kinase STK15/BTAK induces centrosome amplification, aneuploidy and transformation. *Nat. Genet.* 20, 189-193 doi:10.1038/2496
- [18] Gorgun, G., Calabrese, E., Hideshima, T., Ecsedy, J., Perrone, G., Mani, M. et al. (2010) A novel Aurora-A kinase inhibitor MLN8237 induces cytotoxicity and cell-cycle arrest in multiple myeloma. *Blood* 115, 5202-5213 doi:10.1182/blood-2009-12-259523
- [19] Huck, J. J., Zhang, M., McDonald, A., Bowman, D., Hoar, K. M., Stringer, B. et al. (2010) MLN8054, an inhibitor of Aurora A kinase, induces senescence in human tumor cells both in vitro and in vivo. *Mol. Cancer Res.* 8, 373-384 doi:10.1158/1541-7786.MCR-09-0300
- [20] Nikonova, A. S., Astsaturov, I., Serebriiskii, I. G., Dunbrack, R. L. and Golemis, E. A. (2013) Aurora A kinase (AURKA) in normal and pathological cell division. *Cell Mol. Life Sci.* 70, 661-687 doi:10.1007/s00018-01-1073-7
- [21] Margulis, L. (1980) Undulipodia, flagella and cilia. *Biosystems* 12, 105-108 doi:10.1016/0303-2647(80)90041-6
- [22] Lewin, B. 2007 *Cells*. Sudbury, MA: Jones and Bartlett Publishers. XIX, 863 p
- [23] Singh, M., Chaudhry, P. and Merchant, A. A. (2016) Primary cilia are present on human blood and bone marrow cells and mediate Hedgehog signaling. *Exp. Hematol.* doi:10.1016/j.exphem.2016.08.009
- [24] Bangs, F. K., Schrode, N., Hadjantonakis, A.-K. and Anderson, K. V. (2015) Lineage specificity of primary cilia in the mouse embryo. *Nat. Cell. Biol.* 17, 113-122 doi:10.1038/ncb3091
- [25] Kowalevsky, A. O. 1867 *Entwicklungsgeschichte des Amphioxus lanceolatus*. Mémoires de l'Académie impériale des sciences de St-Pétersbourg VIIe série, p. 17. Eggers, St.-Pétersbourg
- [26] Sorokin, S. (1962) Centrioles and the formation of rudimentary cilia by fibroblasts and smooth muscle cells. *J. Cell Biol.* 15, 363-377 doi:10.1083/jcb.15.2.363
- [27] Wong, S. Y., Seol, A. D., So, P.-L., Ermilov, A. N., Bichakjian, C. K., Epstein, E. H. et al. (2009) Primary cilia can both mediate and suppress Hedgehog pathway-dependent tumorigenesis. *Nat. Med.* 15, 1055-1061 doi:10.1038/nm.2011
- [28] Suizu, F., Hirata, N., Kimura, K., Edamura, T., Tanaka, T., Ishigaki, S. et al. (2016) Phosphorylation-dependent Akt-Inversin interaction at the basal body of primary cilia. *EMBO J.* 35, 1346-1363 doi:10.15252/embj.201593003
- [29] Simons, M., Gloy, J., Ganner, A., Bullerkotte, A., Bashkurov, M., Krönig, C. et al. (2005) Inversin, the gene product mutated in nephronophthisis type II, functions as a molecular switch between Wnt signaling pathways. *Nat. Genet.* 37, 537-543 doi:10.1038/ng1552
- [30] Ezratty, E. J., Stokes, N., Chai, S., Shah, A. S., Williams, S. E. and Fuchs, E. (2011) A role for the primary cilium in Notch signaling and epidermal differentiation during skin development. *Cell* 145, 1129-1141 doi:10.1016/j.cell.2011.05.030

- [31] Clement, D. L., Mally, S., Stock, C., Lethan, M., Satir, P., Schwab, A. et al. (2013) PDGFR α signaling in the primary cilium regulates NHE1-dependent fibroblast migration via coordinated differential activity of MEK1/2-ERK1/2-p90RSK and AKT signaling pathways. *J. Cell Sci.* 126(Pt 4), 953-965 doi:10.1242/jcs.116426
- [32] Hou, X., Mrug, M., Yoder, B. K., Lefkowitz, E. J., Kremmidiotis, G., D'Eustachio, P. et al. (2002) Cystin, a novel cilia-associated protein, is disrupted in the cpk mouse model of polycystic kidney disease. *J. Clin. Invest.* 109, 533-540 doi:10.1172/JCI0214099
- [33] Pazour, G. J., San Agustin, J. T., Follit, J. A., Rosenbaum, J. L. and Witman, G. B. (2002) Polycystin-2 localizes to kidney cilia and the ciliary level is elevated in orpk mice with polycystic kidney disease. *Curr. Biol.* 12, R378-R380 doi:10.1016/S0960-9822(02)00877-1
- [34] Sorokin, S. P. (1968) Centriole formation and ciliogenesis. *Aspen Emphysema Conf.* 11, 213-216 PMID:5751848
- [35] Archer, F. L. and Wheatley, D. N. (1971) Cilia in cell-cultured fibroblasts. II. Incidence in mitotic and post-mitotic BHK 21-C13 fibroblasts. *J. Anat.* 109(Pt 2), 277-292 PMID:5105129
- [36] Pazour, G. J., Dickert, B. L. and Witman, G. B. (1999) The DHC1b (DHC2) isoform of cytoplasmic dynein is required for flagellar assembly. *J. Cell Biol.* 144, 473-481 doi:10.1083/jcb.144.3.473
- [37] Pazour, G. J., Dickert, B. L., Vucica, Y., Seeley, E. S., Rosenbaum, J. L., Witman, G. B. et al. (2000) *Chlamydomonas* IFT88 and its mouse homologue, polycystic kidney disease gene Tg737, are required for assembly of cilia and flagella. *J. Cell Biol.* 151, 709-718 doi:10.1083/jcb.151.3.709
- [38] Pugacheva, E. N., Jablonski, S. A., Hartman, T. R., Henske, E. P. and Golemis, E. A. (2007) HEF1-dependent Aurora A activation induces disassembly of the primary cilium. *Cell* 129, 1351-1363 doi:10.1016/j.cell.2007.04.035
- [39] Pan, J. and Snell, W. J. (2000) Regulated targeting of a protein kinase into an intact flagellum. An Aurora/Ipl1p-like protein kinase translocates from the cell body into the flagella during gamete activation in *Chlamydomonas*. *J. Biol. Chem.* 275, 24106-24114 doi:10.1074/jbc.M002686200
- [40] Davis, E. E., Brueckner, M. and Katsanis, N. (2006) The emerging complexity of the vertebrate cilium: new functional roles for an ancient organelle. *Dev. Cell* 11, 9-19 doi:10.1016/j.devcel.2006.06.009
- [41] Marshall, W. F. (2008) Basal bodies platforms for building cilia. *Curr. Top. Dev. Biol.* 85, 1-22 doi:10.1016/S0070-2153(08)00801-6
- [42] Ostrowski, L. E., Dutcher, S. K. and Lo, C. W. (2011) Cilia and models for studying structure and function. *Proc. Am. Thorac. Soc.* 8, 423-429 doi:10.1513/pats.201103-027SD
- [43] Yuan, S. and Sun, Z. (2013) Expanding horizons: ciliary proteins reach beyond cilia. *Annu. Rev. Genet.* 47, 353-376 doi:10.1146/annurev-genet-111212-133243
- [44] Sánchez, I. and Dynlacht, B. D. (2016) Cilium assembly and disassembly. *Nat Cell Biol.* 18, 711-717 doi:10.1038/ncb3370
- [45] Miyamoto, T., Hosoba, K., Ochiai, H., Royba, E., Izumi, H., Sakuma, T. et al. (2015) The microtubule-depolymerizing activity of a mitotic kinesin protein KIF2A drives primary cilia disassembly coupled with cell proliferation. *Cell Rep.* 10, 664-673 doi:10.1016/j.celrep.2015.01.003
- [46] Kim, S., Lee, K., Choi, J.-H., Ringstad, N. and Dynlacht, B. D. (2015) Nek2 activation of Kif24 ensures cilium disassembly during the cell cycle. *Nat Commun.* 6, 8087 doi:10.1038/ncomms9087
- [47] Ding, X.-F., Zhou, J., Hu, Q.-Y., Liu, S.-C. and Chen, G. (2015) The tumor suppressor pVHL down-regulates never-in-mitosis A-related kinase 8 via hypoxia-inducible factors to maintain cilia in human renal cancer cells. *J. Biol. Chem.* 290, 1389-1394 doi:10.1074/jbc.M114.589226
- [48] Grampa, V., Delous, M., Zaidan, M., Ody, G., Thomas, S. and Elkhartoufi, N. (2016) Novel NEK8 mutations cause severe syndromic renal cystic dysplasia through YAP dysregulation. *PLoS Genet* 12, e1005894 doi:10.1371/journal.pgen.1005894
- [49] Pugacheva, E. N. and Golemis, E. A. (2005) The focal adhesion scaffolding protein HEF1 regulates activation of the Aurora-A and Nek2 kinases at the centrosome. *Nat Cell Biol.* 7, 937-946 doi:10.1038/ncb1309
- [50] Shida, T., Cueva, J. G., Xu, Z., Goodman, M. B. and Nachury, M. V. (2010) The major α -tubulin K40 acetyltransferase α TAT1 promotes rapid ciliogenesis and efficient mechanosensation. *Proc. Natl. Acad. Sci. U. S. A.* 107, 21517-21522 doi:10.1073/pnas.1013728107
- [51] Kinzel, D., Boldt, K., Davis, E. E., Bartscher, I., Trümbach, D., Diplas, B. et al. (2010) Pitchfork regulates primary cilia disassembly and left-right asymmetry. *Dev. Cell.* 19, 66-77 doi:10.1016/j.devcel.2010.06.005
- [52] Jung, B., Padula, D., Bartscher, I., Landerer, C., Lutter, D., Theis, F. et al. (2016) Pitchfork and Gprasp2 target smoothed to the primary cilium for Hedgehog pathway activation. *PLoS ONE* 11, e0149477 doi:10.1371/journal.pone.0149477
- [53] Nishizawa, M., Izawa, I., Inoko, A., Hayashi, Y., Nagata, K.-I., Yokoyama, T. et al. (2005) Identification of trichoplein, a novel keratin filament-binding protein. *J. Cell Sci.* 118(Pt 5), 1081-1090 doi:10.1242/jcs.01667

- [54] Inoko, A., Matsuyama, M., Goto, H., Ohmuro-Matsuyama, Y., Hayashi, Y., Enomoto, M. et al. (2012) Trichoplein and Aurora A block aberrant primary cilia assembly in proliferating cells. *J. Cell Biol.* 197, 391-405 doi:10.1083/jcb.201106101
- [55] Inaba, H., Goto, H., Kasahara, K., Kumamoto, K., Yonemura, S., Inoko, A. et al. (2016) Nde1 suppresses ciliogenesis in proliferating cells by regulating the trichoplein-Aurora A pathway. *J. Cell Biol.* 212, 409-423 doi:10.1083/jcb.201507046
- [56] Kim, S., Zaghloul, N. A., Bubenshchikova, E., Oh, E. C., Rankin, S., Katsanis, N. et al. (2011) Nde1-mediated inhibition of ciliogenesis affects cell cycle re-entry. *Nat Cell Biol.* 13, 351-360 doi:10.1038/ncb2183
- [57] Doobin, D. J., Kemal, S., Dantas, T. J. and Vallee, R. B. (2016) Severe NDE1-mediated microcephaly results from neural progenitor cell cycle arrests at multiple specific stages. *Nat Commun.* 7, 12551 doi:10.1038/ncomms12551
- [58] Gabriel, E., Wason, A., Ramani, A., Gooi, L. M., Keller, P., Pozniakovsky, A. et al. (2016) CPAP promotes timely cilium disassembly to maintain neural progenitor pool. *EMBO J.* 35, 803-819 doi:10.15252/embj.201593679
- [59] Plotnikova, O. V., Nikonova, A. S., Loskutov, Y. V., Kozyulina, P. Y., Pugacheva, E. N. and Golemis, E. A. (2012) Calmodulin activation of Aurora-A kinase (AURKA) is required during ciliary disassembly and in mitosis. *Mol. Biol. Cell* 23, 2658-2670 doi:10.1091/mbc.E11-12-1056
- [60] Plotnikova, O. V., Pugacheva, E. N., Dunbrack, R. L. and Golemis, E. A. (2010) Rapid calcium-dependent activation of Aurora-A kinase. *Nat Commun.* 1, Article Number: 64 (2010) doi:10.1038/ncomms1061
- [61] Giet, R. and Prigent, C. (2001) The non-catalytic domain of the *Xenopus laevis* auroraA kinase localises the protein to the centrosome. *J. Cell Sci.* 114(Pt 11), 2095-2104 PMID11493645
- [62] Pugacheva, E. N. and Golemis, E. A. (2006) HEF1-aurora A interactions: points of dialog between the cell cycle and cell attachment signaling networks. *Cell Cycle* 5, 384-391 doi:10.4161/cc.5.4.2439
- [63] Littlepage, L. E. and Ruderman, J. V. (2002) Identification of a new APC/C recognition domain, the A box, which is required for the Cdh1-dependent destruction of the kinase Aurora-A during mitotic exit. *Genes Dev.* 16, 2274-2285 doi:10.1101/gad.1007302
- [64] Nielsen, B. S., Malinda, R. R., Schmid, F. M., Pedersen, S. F., Christensen, ST., Pedersen, L. B. (2015) PDGFR β and oncogenic mutant PDGFR α D842V promote disassembly of primary cilia through a PLC γ - and AURKA-dependent mechanism. *J. Cell Sci.* 128, 3543-3549 doi:10.1242/jcs.173559
- [65] Plotnikova, O. V., Seo, S., Cottle, D. L., Conduit, S., Hakim, S., Dyson, J. M. et al. (2015) INPP5E interacts with AURKA, linking phosphoinositide signaling to primary cilium stability. *J. Cell Sci.* 128, 364-372 doi:10.1242/jcs.161323
- [66] Lee, K. H., Johmura, Y., Yu, L.-R., Park, J.-E., Gao, Y., Bang, J. K. et al. (2012) Identification of a novel Wnt5a-CK1 α -Dvl2-Plk1-mediated primary cilia disassembly pathway. *EMBO J.* 31, 3104-3117 doi:10.1038/emboj.2012.144
- [67] Milella, M., Falcone, I., Conciatori, F., Cesta Incani, U., Del Curatolo, A., Inzerilli, N. et al. (2015) PTEN: multiple functions in human malignant tumors. *Front Oncol.* 5, 24 doi:10.3389/fonc.2015.00024
- [68] Tesio, M., Trinquand, A., Macintyre, E. and Asnafi, V. (2016) Oncogenic PTEN functions and models in T-cell malignancies. *Oncogene* 35, 3887-3896 doi:10.1038/onc.2015.462
- [69] Shnitsar, I., Bashkurov, M., Masson, G. R., Ogunjimi, A. A., Mosessian, S., Cabeza, E. A. et al. (2015) PTEN regulates cilia through Dishevelled. *Nat Commun.* 6, 8388 doi:10.1038/ncomms9388
- [70] Jang, C.-Y., Coppinger, J. A., Seki, A., Yates, J. R. and Fang, G. (2009) Plk1 and Aurora A regulate the depolymerase activity and the cellular localization of Kif2a. *J. Cell Sci.* 122(Pt 9), 1334-1341 doi:10.1242/jcs.044321
- [71] Gong, F., Liu, H., Li, J., Xue, L. and Zhang, M. (2014) Peroxiredoxin 1 is involved in disassembly of flagella and cilia. *Biochem. Biophys. Res. Commun.* 444, 420-426 doi:10.1016/j.bbrc.2014.01.081
- [72] Ji, H., Ramsey, M. R., Hayes, D. N., Fan, C., McNamara, K., Kozlowski, P. et al. (2007) LKB1 modulates lung cancer differentiation and metastasis. *Nature* 448, 807-810 doi:10.1038/nature06030
- [73] Spalluto, C., Wilson, D. I. and Hearn, T. (2012) Nek2 localises to the distal portion of the mother centriole/basal body and is required for timely cilium disassembly at the G2/M transition. *Eur. J. Cell Biol.* 91, 675-686 doi:10.1016/j.ejcb.2012.03.009
- [74] Esteban, M. A. (2006) Formation of primary cilia in the renal epithelium is regulated by the von hippel-Lindau tumor suppressor protein. *J. Am. Soc. Nephrol.* 17, 1801-1806 doi:10.1681/ASN.2006020181
- [75] Kuehn, E. W., Walz, G. and Benzing, T. (2007) von Hippel-Lindau: a tumor suppressor links microtubules to ciliogenesis and cancer development. *Cancer Res.* 67, 4537-4540 doi:10.1158/0008-5472.CAN-07-0391
- [76] Lutz, M. S. and Burk, R. D. (2006) Primary cilium formation requires von Hippel-Lindau gene function in renal-derived cells. *Cancer Res.* 66, 6903-6907 doi:10.1158/0008-5472.CAN-06-0501
- [77] Dere, R., Perkins, A. L., Bawa-Khalfe, T., Jonasch, D. and Walker, C. L. (2015) β -catenin links von Hippel-Lindau to aurora kinase A and loss of primary cilia in renal cell carcinoma. *J. Am. Soc. Nephrol.* 26, 553-564 doi:10.1681/ASN.2013090984

- [78] Otto, E. A., Trapp, M. L., Schultheiss, U. T., Helou, J., Quarmby, L. M. and Hildebrandt, F. (2008) NEK8 mutations affect ciliary and centrosomal localization and may cause nephronophthisis. *J. Am. Soc. Nephrol.* 19, 587-592 doi:10.1681/ASN.2007040490
- [79] Trapp, M. L., Galtseva, A., Manning, D. K., Beier, D. R., Rosenblum, N. D. and Quarmby, L. M. (2008) Defects in ciliary localization of Nek8 is associated with cystogenesis. *Pediatr. Nephrol.* 23, 377-387 doi:10.1007/s00467-007-0692-y
- [80] Sohara, E., Luo, Y., Zhang, J., Manning, D. K., Beier, D. R. and Zhou, J. (2008) Nek8 regulates the expression and localization of polycystin-1 and polycystin-2. *J. Am. Soc. Nephrol.* 19, 469-476 doi:10.1681/ASN.2006090985
- [81] Zalli, D., Bayliss, R. and Fry, A. M. (2012) The Nek8 protein kinase, mutated in the human cystic kidney disease nephronophthisis, is both activated and degraded during ciliogenesis. *Hum. Mol. Genet.* 21, 1155-1171 doi:10.1093/hmg/ddr544
- [82] Liu-Chittenden, Y., Huang, B., Shim, J. S., Chen, Q., Lee, S.-J., Anders, R. A. et al. (2012) Genetic and pharmacological disruption of the TEAD-YAP complex suppresses the oncogenic activity of YAP. *Genes Dev.* 26, 1300-1305 doi:10.1101/gad.192856.112
- [83] Izawa, I., Goto, H., Kasahara, K. and Inagaki, M. (2015) Current topics of functional links between primary cilia and cell cycle. *Cilia* 4, 12 doi:10.1186/s13630-015-0021-1
- [84] Tanos, B. E., Yang, H.-J., Soni, R., Wang, W.-J., Macaluso, F. P., Asara, J. M. et al. (2013) Centriole distal appendages promote membrane docking, leading to cilia initiation. *Genes Develop.* 27, 163-168 doi:10.1101/gad.207043.112
- [85] Kobayashi, T., Tsang, W. Y., Li, J., Lane, W. and Dynlacht, B. D. (2011) Centriolar kinesin Kif24 interacts with CP110 to remodel microtubules and regulate ciliogenesis. *Cell* 145, 914-925 doi:10.1016/j.cell.2011.04.028
- [86] Goetz, S. C., Liem, Jr, K. F. and Anderson, K. V. (2012) The spinocerebellar ataxia-associated gene Tau tubulin kinase 2 controls the initiation of ciliogenesis. *Cell* 151, 847-858 doi:10.1016/j.cell.2012.10.010
- [87] Kuhns, S., Schmidt, K. N., Reymann, J., Gilbert, D. F., Neuner, A., Hub, B. et al. (2013) The microtubule affinity regulating kinase MARK4 promotes axoneme extension during early ciliogenesis. *J. Cell Biol.* 200, 505-522 doi:10.1083/jcb.201206013
- [88] Xu, Q., Zhang, Y., Wei, Q., Huang, Y., Hu, J. and Ling, K. (2016) Phosphatidylinositol phosphate kinase PIPKly and phosphatase INPP5E coordinate initiation of ciliogenesis. *Nat. Commun.* 7, 10777 doi:10.1038/ncomms10777
- [89] Schmidt, K. N., Kuhns, S., Neuner, A., Hub, B., Zentgraf, H. and Pereira, G. (2012) Cep164 mediates vesicular docking to the mother centriole during early steps of ciliogenesis. *J. Cell Biol.* 199, 1083-1101 doi:10.1083/jcb.201202126
- [90] Knodler, A., Feng, S., Zhang, J., Zhang, X., Das, A., Peranen, J. et al. (2010) Coordination of Rab8 and Rab11 in primary ciliogenesis. *Proc. Natl Acad. Sci. U. S. A.* 107, 6346-6351 doi:10.1073/pnas.1002401107
- [91] Yoshimura, S.-i., Egerer, J., Fuchs, E., Haas, A. K. and Barr, F. A. (2007) Functional dissection of Rab GTPases involved in primary cilium formation. *J. Cell Biol.* 178, 363-369 doi:10.1083/jcb.200703047
- [92] Westlake, C. J., Baye, L. M., Nachury, M. V., Wright, K. J., Ervin, K. E., Phu, L. et al. (2011) Primary cilia membrane assembly is initiated by Rab11 and transport protein particle II (TRAPP II) complex-dependent trafficking of Rabin8 to the centrosome. *Proc. Natl Acad. Sci. U. S. A.* 108, 2759-2764 doi:10.1073/pnas.1018823108
- [93] Franco, I., Gulluni, F., Campa, C. C., Costa, C., Margaria, J. P., Ciraolo, E. et al. (2014) PI3K class II alpha controls spatially restricted endosomal PtdIns3P and Rab11 activation to promote primary cilium function. *Dev. Cell* 28, 647-658 doi:10.1016/j.devcel.2014.01.022
- [94] Zhang, B., Zhang, T., Wang, G., Wang, G., Chi, W., Jiang, Q. et al. (2015) GSK3 β -Dzip1-Rab8 cascade regulates ciliogenesis after mitosis. *PLoS Biol.* 13, e1002129 doi:10.1371/journal.pbio.1002129
- [95] Walter, A. O., Seghezzi, W., Korver, W., Sheung, J. and Lees, E. (2000) The mitotic serine/threonine kinase Aurora2/AIK is regulated by phosphorylation and degradation. *Oncogene* 19, 4906-4916 doi:10.1038/sj.onc.1203847
- [96] Mergen, M., Engel, C., Muller, B., Follo, M., Schafer, T., Jung, M. et al. (2013) The nephronophthisis gene product NPHP2/Inversin interacts with Aurora A and interferes with HDAC6-mediated cilia disassembly. *Nephrol. Dial. Transplant.* 28, 2744-2753 doi:10.1093/ndt/gft316
- [97] Kasahara, K., Kawakami, Y., Kiyono, T., Yonemura, S., Kawamura, Y., Era, S. et al. (2014) Ubiquitin-proteasome system controls ciliogenesis at the initial step of axoneme extension. *Nat. Commun.* 5, 5081 doi:10.1038/ncomms6081
- [98] Kim, J., Jo, H., Hong, H., Kim, M. H., Kim, J. M., Lee, J.-K. et al. (2015) Actin remodelling factors control ciliogenesis by regulating YAP/TAZ activity and vesicle trafficking. *Nat. Commun.* 6, 6781 doi:10.1038/ncomms7781
- [99] Kim, M., Kim, M., Lee, M.-S., Kim, C.-H. and Lim, D.-S. (2014) The MST1/2-SAV1 complex of the Hippo pathway promotes ciliogenesis. *Nat. Commun.* 5, 5370 doi:10.1038/ncomms6370

- [100] Brown, J. M. and Witman, G. B. (2014) Cilia and diseases. *BioScience* 64, 1126-1137 doi:10.1093/biosci/biu174
- [101] Habbig, S. and Liebau, M. C. (2015) Ciliopathies - from rare inherited cystic kidney diseases to basic cellular function. *Mol. Cell. Pediatr.* 2, 8 doi:10.1186/s40348-015-0019-1
- [102] Waters, A. M. and Beales, P. L. (2011) Ciliopathies: an expanding disease spectrum. *Pediatr. Nephrol.* 26, 1039-1056 doi:10.1007/s00467-010-1731-7
- [103] Igarashi, P. and Somlo, S. (2002) Genetics and pathogenesis of polycystic kidney disease. *J. Am. Soc. Nephrol.* 13, 2384-2398 doi:10.1097/01.ASN.0000028643.17901.42
- [104] Torres, V. E. and Harris, P. C. (2009) Autosomal dominant polycystic kidney disease: the last 3 years. *Kidney Int.* 76, 149-168 doi:10.1038/ki.2009.128
- [105] Deltas, C. and Papagregoriou, G. (2010) Cystic diseases of the kidney: molecular biology and genetics. *Arch. Pathol. Lab Med.* 134, 569-582 doi:10.1043/1543-2165-134.4.569
- [106] Gunay-Aygun, M. (2009) Liver and kidney disease in ciliopathies. *Am. J. Med. Genet. C Semin. Med. Genet.* 151, 296-306 doi:10.1002/ajmg.c.30225
- [107] Luijten, M. N. H., Basten, S. G., Claessens, T., Vernooij, M., Scott, C. L., Janssen, R. et al. (2013) Birt-Hog-Dube syndrome is a novel ciliopathy. *Hum. Mol. Genet.* 22, 4383-4397 doi:10.1093/hmg/ddt288
- [108] Barker, A. R., Thomas, R. and Dawe, H. R. (2014) Meckel-Gruber syndrome and the role of primary cilia in kidney, skeleton, and central nervous system development. *Organogenesis* 10, 96-107 doi:10.4161/org.27375
- [109] Ben-Salem, S., Al-Shamsi, A. M., Gleeson, J. G., Ali, B. R. and Al-Gazali, L. (2014) Mutation spectrum of Joubert syndrome and related disorders among Arabs. *Hum. Genome Var.* 1, 14020 doi:10.1038/hgv.2014.20
- [110] Franco, B. and Thauvin-Robinet, C. (2016) Update on oral-facial-digital syndromes (OFDS). *Cilia* 5, 12 doi:10.1186/s13630-016-0034-4
- [111] Sattar, S. and Gleeson, J. G. (2011) The ciliopathies in neuronal development: a clinical approach to investigation of Joubert syndrome and Joubert syndrome-related disorders. *Dev. Med. Child Neurol.* 53, 793-798 doi:10.1111/j.1469-8749.2011.04021.x
- [112] Schock, E. N., Chang, C.-F., Youngworth, I. A., Davey, M. G., Delany, M. E., Brugmann, S. A. (2016) Utilizing the chicken as an animal model for human craniofacial ciliopathies. *Dev. Biol.* 415, 326-337 doi:10.1016/j.ydbio.2015.10.024
- [113] Riaz, M. and Baird, P. N. (2016) Genetics in retinal diseases. *Dev. Ophthalmol.* 55, 57-62 doi:10.1159/000431142
- [114] Ronquillo, C. C., Bernstein, P. S. and Baehr, W. (2012) Senior-Loken syndrome: a syndromic form of retinal dystrophy associated with nephronophthisis. *Vision Res.* 75, 88-97 doi:10.1016/j.visres.2012.07.003
- [115] Khan, S. A., Muhammad, N., Khan, M. A., Kamal, A., Rehman, Z. U. and Khan, S. (2016) Genetics of human Bardet-Biedl syndrome, an updates. *Clin. Genet.* 90, 3-15 doi:10.1111/cge.12737
- [116] Suspitsin, E. N. and Imyanitov, E. N. (2016) Bardet-Biedl syndrome. *Mol. Syndromol.* 7, 62-71 doi:10.1159/000445491
- [117] Álvarez-Satta, M., Castro-Sánchez, S. and Valverde, D. (2015) Alström syndrome: current perspectives. *Appl. Clin. Genet.* 8, 171-179 doi:10.2147/TACG.S56612
- [118] Stokman, M., Lilien, M. and Knoers, N. (1993) Nephronophthisis. In *GeneReviews®* (R. A. Pagon et al. ed.), University of Washington, Seattle, WA
- [119] Keppler-Noreuil, K. M., Adam, M. P., Welch, J., Muilenburg, A. and Willing, M. C. (2011) Clinical insights gained from eight new cases and review of reported cases with Jeune syndrome (asphyxiating thoracic dystrophy). *Am. J. Med. Genet. A* 155, 1021-1032 doi:10.1002/ajmg.a.33892
- [120] Huber, C. and Cormier-Daire, V. (2012) Ciliary disorder of the skeleton. *Am. J. Med. Genet. C Semin. Med. Genet.* 160C, 165-174 doi:10.1002/ajmg.c.31336
- [121] Mariman, E. C. M., Vink, R. G., Roumans, N. J. T., Bouwman, F. G., Stumpel, C. T. R. M., Aller, E. E. J. G. et al. (2016) The cilium: a cellular antenna with an influence on obesity risk. *Br. J. Nutr.* 116, 576-592 doi:10.1017/S0007114516002282
- [122] Lee, H., Song, J., Jung, J. H. and Ko, H. W. (2015) Primary cilia in energy balance signaling and metabolic disorder. *BMB Rep.* 48, 647-654 doi:10.5483/BMBRep.2015.48.12.229
- [123] Shalata, A., Ramirez, M. C., Desnick, R. J., Priedigkeit, N., Buettner, C., Lindtner, C. et al. (2013) Morbid obesity resulting from inactivation of the ciliary protein CEP19 in humans and mice. *Am. J. Hum. Genet.* 93, 1061-1071 doi:10.1016/j.ajhg.2013.10.025
- [124] Dinsmore, C. and Reiter, J. F. (2016) Endothelial primary cilia inhibit atherosclerosis. *EMBO Rep.* 17, 156-166 doi:10.15252/embr.201541019
- [125] Willemarck, N., Rysman, E., Brusselmans, K., Van Imschoot, G., Vanderhoydonc, F., Moerloose, K. et al. (2010) Aberrant activation of fatty acid synthesis suppresses primary cilium formation and distorts tissue development. *Cancer Res.* 70, 9453-9462 doi:10.1158/0008-5472.CAN-10-2324

- [126] Seeger-Nukpezah, T., Geynisman, D. M., Nikonova, A. S., Benzing, T. and Golemis, E. A. (2015) The hallmarks of cancer: relevance to the pathogenesis of polycystic kidney disease. *Nat. Rev. Nephrol.* 11, 515-534 doi:10.1038/nrneph.2015.46
- [127] Xu, J., Li, H., Wang, B., Xu, Y., Yang, J., Zhang, X. et al. (2010) VHL inactivation induces HEF1 and Aurora kinase A. *J. Am. Soc. Nephrol.* 21, 2041-2046 doi:10.1681/ASN.2010040345
- [128] Nikonova, A. S., Plotnikova, O. V., Serzhanova, V., Efimov, A., Bogush, I., Cai, K. Q. et al. (2014) Nedd9 restrains renal cystogenesis in Pkd1^{-/-} mice. *Proc. Natl Acad. Sci. U. S. A.* 111, 12859-12864 doi:10.1073/pnas.1405362111
- [129] Nikonova, A. S., Deneka, A. Y., Eckman, L., Kopp, M. C., Hensley, H. H., Egleston, B. L. et al. (2015) Opposing effects of inhibitors of Aurora-A and EGFR in autosomal-dominant polycystic kidney disease. *Front. Oncol.* 5, 228 doi:10.3389/fonc.2015.00228
- [130] Badano, J. L., Mitsuma, N., Beales, P. L. and Katsanis, N. (2006) The ciliopathies: an emerging class of human genetic disorders. *Annu. Rev. Genomics. Hum. Genet.* 7, 125-148 doi:10.1146/annurev.genom.7.080505.115610
- [131] Hampshire, D. J., Ayub, M., Springell, K., Roberts, E., Jafri, H., Rashid, Y. et al. (2006) MORM syndrome (mental retardation, truncal obesity, retinal dystrophy and micropenis), a new autosomal recessive disorder, links to 9q34. *Eur. J. Hum. Genet.* 14, 543-548 doi:10.1038/sj.ejhg.5201577
- [132] Egeberg, D. L., Lethan, M., Manguso, R., Schneider, L., Awan, A., Jørgensen, T. S. et al. (2012) Primary cilia and aberrant cell signaling in epithelial ovarian cancer. *Cilia* 1, 15 doi:10.1186/2046-2530-1-15
- [133] Johnson, C. A. and Collis, S. J. (2016) Ciliogenesis and the DNA damage response: a stressful relationship. *Cilia* 5, 19 doi:10.1186/s13630-016-0040-6
- [134] Hassounah, N. B., Bunch, T. A. and McDermott, K. M. (2012) Molecular pathways: the role of primary cilia in cancer progression and therapeutics with a focus on Hedgehog signaling. *Clin. Cancer Res.* 18, 2429-2435 doi:10.1158/1078-0432.CCR-11-0755
- [135] Plotnikova, O. V., Golemis, E. A. and Pugacheva, E. N. (2008) Cell cycle-dependent ciliogenesis and cancer. *Cancer Res.* 68, 2058-2061 doi:10.1158/0008-5472.CAN-07-5838
- [136] Seeger-Nukpezah, T., Little, J. L., Serzhanova, V. and Golemis, E. A. (2013) Cilia and cilia-associated proteins in cancer. *Drug Discov. Today Dis. Mech.* 10, e135-e142 doi:10.1016/j.ddmec.2013.03.004
- [137] Shagisultanova, E., Gaponova, A. V., Gabbasov, R., Nicolas, E. and Golemis, E. A. (2015) Preclinical and clinical studies of the NEDD9 scaffold protein in cancer and other diseases. *Gene* 567, 1-11 doi:10.1016/j.gene.2015.04.086
- [138] Janecek, M., Rossmann, M., Sharma, P., Emery, A., Huggins, D. J., Stockwell, S. R. et al. (2016) Allosteric modulation of AURKA kinase activity by a small-molecule inhibitor of its protein-protein interaction with TPX2. *Sci. Rep.* 6, 28528 doi:10.1038/srep28528
- [139] Shagisultanova, E., Dunbrack, Jr, R. L. and Golemis, E. A. (2015) Issues in interpreting the in vivo activity of Aurora-A. *Expert Opin. Ther. Targets* 19, 187-200 doi:10.1517/14728222.2014.981154