

# The relative contribution of nmdars to excitatory postsynaptic currents is controlled by $Ca^{2+}$ -induced inactivation

Valiullina F., Zakharova Y., Mukhtarov M., Draguhn A., Burnashev N., Rozov A.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2016 Valiullina, Zakharova, Mukhtarov, Draguhn, Burnashev and Rozov. NMDA receptors (NMDARs) are important mediators of excitatory synaptic transmission and plasticity. A hallmark of these channels is their high permeability to  $Ca^{2+}$ . At the same time, they are themselves inhibited by the elevation of intracellular  $Ca^{2+}$  concentration. It is unclear however, whether the  $Ca^{2+}$  entry associated with single NMDAR mediated synaptic events is sufficient to self-inhibit their activation. Such auto-regulation would have important effects on the dynamics of synaptic excitation in several central neuronal networks. Therefore, we studied NMDAR-mediated synaptic currents in mouse hippocampal CA1 pyramidal neurons. Postsynaptic responses to subthreshold Schaffer collateral stimulation depended strongly on the absence or presence of intracellular  $Ca^{2+}$  buffers. Loading of pyramidal cells with exogenous  $Ca^{2+}$  buffers increased the amplitude and decay time of NMDAR mediated EPSCs (EPSPs) and prolonged the time window for action potential (AP) generation. Our data indicate that the  $Ca^{2+}$  influx mediated by unitary synaptic events is sufficient to produce detectable self-inhibition of NMDARs even at a physiological  $Mg^{2+}$  concentration. Therefore, the contribution of NMDARs to synaptic excitation is strongly controlled by both previous synaptic activity as well as by the  $Ca^{2+}$  buffer capacity of postsynaptic neurons.

<http://dx.doi.org/10.3389/fncel.2016.00012>

---

## Keywords

Action potentials, Calcium, Excitation, Firing properties, Modulation, NMDA receptor