

Carbon and nitrogen availability in paddy soil affects rice photosynthate allocation, microbial community composition, and priming: combining continuous ^{13}C labeling with PLFA analysis

Ziwei Zhao · Tida Ge  · Anna Gunina · Yuhong Li · Zhenke Zhu · Peiqin Peng · Jinshui Wu · Yakov Kuzyakov

Received: 26 July 2018 / Accepted: 2 November 2018
© Springer Nature Switzerland AG 2018

Abstract

Background and aims Carbon (C) and nitrogen (N) availability in soil change microbial community composition and activity and so, might affect soil organic matter (SOM) decomposition as well as allocation of plant assimilates. The study was focused on interactions between C and N availability and consequences for rhizodeposition and microbial community structure in paddy soil.

Methods Rice continuously labeled in a $^{13}\text{C}\text{O}_2$ atmosphere was fertilized with either carboxymethyl cellulose (CMC) (+C), ammonium sulfate (+N), or their combination (+CN), and unfertilized soil was used as a

control. ^{13}C was traced in aboveground and belowground plant biomass, soil organic matter, and microbial biomass. Microbial community composition was analyzed by phospholipid fatty acids (PLFAs).

Results +CN application led to a higher yield and lower root C and N content: ^{13}C assimilated in shoots increased by 1.39-fold and that in roots decreased by 0.75-fold. Correspondingly, after +CN addition, ^{13}C from rhizodeposits incorporated into SOM and microorganisms decreased by 0.68-fold and 0.53-fold, respectively, as compared with that in the unfertilized soil. The application of +C or +N alone resulted in smaller changes. CMC led to a 3% of total N mobilized from

Responsible Editor: Eric Paterson.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11104-018-3873-5>) contains supplementary material, which is available to authorized users.

Z. Zhao · Y. Kuzyakov
Department of Agricultural Soil Science, Department of Soil Science of Temperate Ecosystems, Georg-August University of Göttingen, 37077 Göttingen, Germany

Z. Zhao · T. Ge (✉) · Y. Li · Z. Zhu · J. Wu
Key Laboratory of Agro-ecological Processes in Subtropical Region & Changsha Research Station for Agricultural and Environmental Monitoring, Institute of Subtropical Agriculture, Chinese Academy of Sciences, Hunan 410125, China
e-mail: gtd@isa.ac.cn

A. Gunina
Department of Soil Biology and Biochemistry, Dokuchaev Soil Science Institute, Moscow, Russia

P. Peng
College of Environmental Science and Engineering, Central South University of Forestry and Technology, Changsha 410004, China

Y. Kuzyakov
Agro-Technology Institute, RUDN University, Moscow, Russia

Y. Kuzyakov
Institute of Environmental Sciences, Kazan Federal University, 420049 Kazan, Russia