

The antioxidant system in the anhydrobiotic midge as an essential, adaptive mechanism for desiccation survival

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© Springer Nature Singapore Pte Ltd. 2018. One of the major damaging factors for living organisms experiencing water insufficiency is oxidative stress. Loss of water causes a dramatic increase in the production of reactive oxygen species (ROS). Thus, the ability for some organisms to survive almost complete desiccation (called anhydrobiosis) is tightly related to the ability to overcome extraordinary oxidative stress. The most complex anhydrobiotic organism known is the larva of the chironomid *Polypedilum vanderplanki*. Its antioxidant system shows remarkable features, such as an expansion of antioxidant genes, their overexpression, as well as the absence or low expression of enzymes required for the synthesis of ascorbate and glutathione and their antioxidant function. In this chapter, we summarize existing data about the antioxidant system of this insect, which is able to cope with substantial oxidative damage, even in an intracellular environment that is severely disturbed due to water loss.

http://dx.doi.org/10.1007/978-981-13-1244-1_14

Keywords

Anhydrobiosis, Antioxidant, Glutathione peroxidase, *P. Vanderplanki*, Superoxide dismutase, Thioredoxin

References

- [1] Bannai H, Tamada Y, Maruyama O, Nakai K, Miyano S (2002) Extensive feature detection of N-terminal protein sorting signals. *Bioinformatics* 18:298-305
- [2] Benaroudj N, Lee DH, Goldberg AL (2001) Trehalose accumulation during cellular stress protects cells and cellular proteins from damage by oxygen radicals. *J Biol Chem* 276:24261-24267
- [3] Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F, Gallo Cassarino T, Bertoni M, Bordoli L et al (2014) SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res* 42:W252-W258
- [4] Brown NM, Torres AS, Doan PE, O'Halloran TV (2004) Oxygen and the copper chaperone CCS regulate post-translational activation of Cu, Zn superoxide dismutase. *Proc Natl Acad Sci U S A* 101:5518-5523
- [5] Buitink J, Leprince O (2004) Glass formation in plant anhydrobiotes: survival in the dry state. *Cryobiology* 48:215-228
- [6] Carroll MC, Girouard JB, Ulloa JL, Subramaniam JR, Wong PC, Valentine JS, Culotta VC (2004) Mechanisms for activating Cu- and Zn-containing superoxide dismutase in the absence of the CCS Cu chaperone. *Proc Natl Acad Sci U S A* 101:5964-5969
- [7] Cornette R, Kikawada T (2011) The induction of anhydrobiosis in the sleeping chironomid: current status of our knowledge. *IUBMB Life* 63:419-429

- [8] Cornette R, Kanamori Y, Watanabe M, Nakahara Y, Gusev O, Mitsumasu K, Kadono-Okuda K, Shimomura M, Mita K, Kikawada T et al (2010) Identification of anhydrobiosis-related genes from an expressed sequence tag database in the cryptobiotic midge *Polypedilum vanderplanki* (Diptera; Chironomidae). *J Biol Chem* 285:35889-35899
- [9] Cornette R, Yamamoto N, Yamamoto M, Kobayashi T, Petrova NA, Gusev O, Shimura S, Kikawada T, Pemba D, Okuda T (2017) A new anhydrobiotic midge from Malawi, *Polypedilum pembai* sp n. (Diptera: Chironomidae), closely related to the desiccation tolerant midge, *Polypedilum vanderplanki* Hinton. *Syst Entomol* 42:814-825
- [10] Corona M, Robinson GE (2006) Genes of the antioxidant system of the honey bee: annotation and phylogeny. *Insect Mol Biol* 15:687-701
- [11] Cranston PS (2014) A new putatively cryptobiotic midge, *Polypedilum ovahimba* sp nov (Diptera: Chironomidae), from southern Africa. *Aust Entomol* 53:373-379
- [12] Crowe JH (2007) Trehalose as a "chemical chaperone": fact and fantasy. In: Csermely P, Vigh L (eds) *Molecular aspects of the stress response: chaperones, membranes and networks*. Springer, New York, pp 143-158
- [13] da Costa Morato Nery D, da Silva CG, Mariani D, Fernandes PN, Pereira MD, Panek AD, Eleutherio EC (2008) The role of trehalose and its transporter in protection against reactive oxygen species. *Biochim Biophys Acta* 1780:1408-1411
- [14] Emanuelsson O, Nielsen H, Brunak S, von Heijne G (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol* 300:1005-1016
- [15] Finkel T, Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. *Nature* 408:239-247
- [16] Foyer CH, Noctor G (2011) Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol* 155:2-18
- [17] França MB, Panek AD, Eleutherio EC (2007) Oxidative stress and its effects during dehydration. *Comp Biochem Physiol A Mol Integr Physiol* 146:621-631
- [18] Furukawa Y, Torres AS, O'Halloran TV (2004) Oxygen-induced maturation of SOD1: a key role for disulfide formation by the copper chaperone CCS. *EMBO J* 23:2872-2881
- [19] Grubor-Lajsic G, Block W, Jovanovic A, Worland R (1996) Antioxidant enzymes in larvae of the Antarctic fly, *Belgica antarctica*. *CryoLetters* 17:39-42
- [20] Gusev O, Nakahara Y, Vanyagina V, Malutina L, Cornette R, Sakashita T, Hamada N, Kikawada T, Kobayashi Y, Okuda T (2010) Anhydrobiosis-associated nuclear DNA damage and repair in the sleeping chironomid: linkage with radioresistance. *PLoS One* 5:e14008
- [21] Gusev O, Suetsugu Y, Cornette R, Kawashima T, Logacheva MD, Kondrashov AS, Penin AA, Hatanaka R, Kikuta S, Shimura S et al (2014) Comparative genome sequencing reveals genomic signature of extreme desiccation tolerance in the anhydrobiotic midge. *Nat Commun* 5:4784
- [22] Herdeiro RS, Pereira MD, Panek AD, Eleutherio EC (2006) Trehalose protects *Saccharomyces cerevisiae* from lipid peroxidation during oxidative stress. *Biochim Biophys Acta* 1760:340-346
- [23] Hinton H (1951) A new Chironomid from Africa, the larva of which can be dehydrated without injury. *Proc Zool Soc Lond* 121:371-380
- [24] Hinton H (1960) A fly larva that tolerates dehydration and temperatures of -270° to $+102^{\circ}$ C. *Nature* 188:336-337
- [25] Holmgren A, Sengupta R (2010) The use of thiols by ribonucleotide reductase. *Free Radic Biol Med* 49:1617-1628
- [26] Indo HP, Davidson M, Yen HC, Suenaga S, Tomita K, Nishii T, Higuchi M, Koga Y, Ozawa T, Majima HJ (2007) Evidence of ROS generation by mitochondria in cells with impaired electron transport chain and mitochondrial DNA damage. *Mitochondrion* 7:106-118
- [27] Jensen LT, Culotta VC (2005) Activation of CuZn superoxide dismutases from *Caenorhabditis elegans* does not require the copper chaperone CCS. *J Biol Chem* 280:41373-41379
- [28] Lartigue A, Burlat B, Coutard B, Chaspoul F, Claverie JM, Abergel C (2015) The Megavirus chilensis Cu, Zn-superoxide dismutase: the first viral structure of a typical cellular copper chaperone-independent hyper-stable dimeric enzyme. *J Virol* 89:824-832
- [29] Leitch JM, Yick PJ, Culotta VC (2009) The right to choose: multiple pathways for activating copper, zinc superoxide dismutase. *J Biol Chem* 284:24679-24683
- [30] Leprince O, Atherton NM, Deltour R, Hendry G (1994) The involvement of respiration in free radical processes during loss of desiccation tolerance in germinating *Zea mays* L. (an electron paramagnetic resonance study). *Plant Physiol* 104:1333-1339
- [31] Lopez-Martinez G, Elnitsky MA, Benoit JB, Lee RE Jr, Denlinger DL (2008) High resistance to oxidative damage in the Antarctic midge *Belgica antarctica*, and developmentally linked expression of genes encoding superoxide dismutase, catalase and heat shock proteins. *Insect Biochem Mol Biol* 38:796-804
- [32] Maiorino M, Ursini F, Bosello V, Toppo S, Tosatto SC, Mauri P, Becker K, Roveri A, Bulato C, Benazzi L et al (2007) The thioredoxin specificity of *Drosophila* GPx: a paradigm for a peroxiredoxin-like mechanism of many glutathione peroxidases. *J Mol Biol* 365:1033-1046

- [33] Meyer Y, Siala W, Bashandy T, Riondet C, Vignols F, Reichheld JP (2008) Glutaredoxins and thioredoxins in plants. *Biochim Biophys Acta* 1783:589-600
- [34] Nair PM, Park SY, Chung JW, Choi J (2013) Transcriptional regulation of glutathione biosynthesis genes, gamma-glutamyl-cysteine ligase and glutathione synthetase in response to cadmium and nonylphenol in *Chironomus riparius*. *Environ Toxicol Pharmacol* 36:265-273
- [35] Nakahara Y, Imanishi S, Mitsumasu K, Kanamori Y, Iwata K, Watanabe M, Kikawada T, Okuda T (2010) Cells from an anhydrobiotic chironomid survive almost complete desiccation. *Cryobiology* 60:138-146
- [36] Nesmelov A, Devatiyarov R, Voronina T, Kondratyeva S, Cherkasov A, Cornette R, Kikawada T, Shagimardanova E (2016) New antioxidant genes from an anhydrobiotic insect: unique structural features in functional motifs of thioredoxins. *BioNanoScience* 6:568-570
- [37] Oku K, Watanabe H, Kubota M, Fukuda S, Kurimoto M, Tsujisaka Y, Komori M, Inoue Y, Sakurai M (2003) NMR and quantum chemical study on the OH $\cdots\pi$ and CH \cdots O interactions between trehalose and unsaturated fatty acids: implication for the mechanism of antioxidant function of trehalose. *J Am Chem Soc* 125:12739-12748
- [38] Patenaude A, Ven Murthy MR, Mirault ME (2004) Mitochondrial thioredoxin system: effects of TrxR2 overexpression on redox balance, cell growth, and apoptosis. *J Biol Chem* 279:27302-27314
- [39] Pereira Ede J, Panek AD, Eleutherio EC (2003) Protection against oxidation during dehydration of yeast. *Cell Stress Chaperones* 8:120-124
- [40] Petersen TN, Brunak S, von Heijne G, Nielsen H (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8:785-786
- [41] Pompella A, Visvikis A, Paolicchi A, De Tata V, Casini AF (2003) The changing faces of glutathione, a cellular protagonist. *Biochem Pharmacol* 66:1499-1503
- [42] Reuter S, Gupta SC, Chaturvedi MM, Aggarwal BB (2010) Oxidative stress, inflammation, and cancer: how are they linked? *Free Radic Biol Med* 49:1603-1616
- [43] Sakurai M, Furuki T, Akao K, Tanaka D, Nakahara Y, Kikawada T, Watanabe M, Okuda T (2008) Vitrification is essential for anhydrobiosis in an African chironomid, *Polypedilum vanderplanki*. *Proc Natl Acad Sci U S A* 105:5093-5098
- [44] Scheerer P, Borchert A, Krauss N, Wessner H, Gerth C, Hohne W, Kuhn H (2007) Structural basis for catalytic activity and enzyme polymerization of phospholipid hydroperoxide glutathione peroxidase-4 (GPx4). *Biochemistry* 46:9041-9049
- [45] Sun WQ, Leopold AC (1995) The Maillard reaction and oxidative stress during aging of soybean seeds. *Physiol Plant* 94:94-104
- [46] Uttara B, Singh AV, Zamboni P, Mahajan R (2009) Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol* 7:65-74
- [47] Watanabe M, Kikawada T, Minagawa N, Yukuhiro F, Okuda T (2002) Mechanism allowing an insect to survive complete dehydration and extreme temperatures. *J Exp Biol* 205:2799-2802
- [48] Watanabe M, Kikawada T, Fujita A, Okuda T (2005) Induction of anhydrobiosis in fat body tissue from an insect. *J Insect Physiol* 51:727-731
- [49] Watanabe M, Nakahara Y, Sakashita T, Kikawada T, Fujita A, Hamada N, Horikawa DD, Wada S, Kobayashi Y, Okuda T (2007) Physiological changes leading to anhydrobiosis improve radiation tolerance in *Polypedilum vanderplanki* larvae. *J Insect Physiol* 53:573-579
- [50] Weisiger RA, Fridovich I (1973) Mitochondrial superoxide dismutase site of synthesis and intramitochondrial localization. *J Biol Chem* 248:4793-4796
- [51] Zelko IN, Mariani TJ, Folz RJ (2002) Superoxide dismutase multigene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free Radic Biol Med* 33:337-349