

A new, large species of *Gomphonema* Ehrenberg from ancient Lake Matano, Indonesia

Kociolek J., Kapustin D., Kulikovskiy M.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018, © 2018 The International Society for Diatom Research. A large species of the diatom genus *Gomphonema* Ehrenberg, *G. matanensis* sp. nov., is described as new to science from ancient Lake Matano, Indonesia. Light and scanning electron microscopic observations of the valve morphology of this species are presented. The morphology of this species is similar to other large species that have ovoid areolar openings without external flaps or occlusions, such as *G. gomphopleuroides* Amossé ex Kociolek, Glushchenko & Kulikovskiy and members of the *G. kaznakowi* species complex, both known from Southeast Asia. This species has the special features of bi- and triradiate siliceous growths at the valve mantle junction, continuing onto the mantle. Though an analysis of the phylogenetic relationships of the diatom genus *Gomphonema* is currently wanting, it would appear that gigantism occurred several times within the genus.

<http://dx.doi.org/10.1080/0269249X.2018.1513868>

Keywords

ancient lakes, Bacillariophyceae, diatoms, *Gomphonema*, Indonesia, morphology, systematics

References

- [1] Amossé, A., 1969. Note sur des Diatomées récoltées en Indochine. *Revue Algologique, Nouvelle Série* 9: 326-344
- [2] Blanco S., Alvarez-Blanco I., Cejudo-Figueiras C., & Becares E., 2012. Contribution to the diatom flora of Cambodia: five new recent freshwater taxa. *Journal of Systematics and Evolution* 50: 258-266. doi: 10.1111/j.1759-6831.2012.00189.x
- [3] Brun J., 1899. Tafel 216. In: *Atlas der Diatomaceenkunde* (Ed. by A., Schmidt). O. R. Reiland, Leipzig
- [4] Cleve P.T., 1894. Synopsis of the naviculoid diatoms. Part 1. *Kongliga Svenska-Vetenskaps Akademiens Handlingar* 26(2): 1-194
- [5] Ehrenberg C.G., 1854. *Mikrogeologie. Das Erden und Felsen schaffende Wirken des unsichtbar kleinen selbstständigen Lebens auf der Erde. Text and Atlas.* Leopold Voss, Leipzig
- [6] Foged N., 1971. Freshwater diatoms in Thailand. *Nova Hedwigia* 22: 267-369
- [7] Foged N., 1976. Freshwater diatoms in Sri Lanka (Ceylon). *Bibliotheca Phycologica* 23: 1-112
- [8] Fournanier E., & Kociolek J.P., 2011. Catalogue of diatom names. Available from: <http://researcharchive.calacademy.org/research/diatoms/names/index.asp>
- [9] Fricke F., 1904. Tafel 248. In: *Atlas der Diatomaceenkunde* (Ed. by A., Schmidt). O.R. Reiland, Leipzig
- [10] Fujita Y., & Ohtsuka T., 2005. Diatoms from paddy fields in northern Laos. *Diatom* 21: 71-89
- [11] Glushchenko A., & Kulikovskiy M., 2015. Species of the genus *Luticola* (Bacillariophyceae) in waterbodies of Laos and Vietnam. *Botanical Journal* 100(8): 799-804 (In Russian)

- [12] Glushchenko A.M., & Kulikovskiy M.S., 2017a. *Amphipleura vavilovii*: a new diatom species of the family Amphipleuraceae from Laos. *Inland Water Biology* 10: 17–21. doi: 10.1134/S1995082916040064
- [13] Glushchenko A.M., & Kulikovskiy M.S., 2017b. Taxonomy and distribution of the genus *Eunotia* Ehrenberg in water ecosystems of Vietnam. *Inland Water Biology* 10: 130–139. doi: 10.1134/S1995082917020055
- [14] Glushchenko A., Genkal S.I., & Kulikovskiy M., 2016. *Aulacoseira konstantinovii* sp. nov. and *Aulacoseira krylovii* sp. nov.—two new centric diatoms from South east Asia. *Diatom Research* 31: 367–378. doi: 10.1080/0269249X.2016.1253617
- [15] Glushchenko A.M., Kulikovskiy M., & Kociolek J.P., 2017. New and interesting species from the genus *Luticola* (Bacillariophyceae) in waterbodies of Southeastern Asia. *Nova Hedwigia*, Beihefte 146: 157–173. doi: 10.1127/1438-9134/2017/157
- [16] Glushchenko A., Kulikovskiy M., Kuznetsova I., Dorofeyuk N., & Kociolek J.P., 2018. New species and combinations in the genus *Eunotia* Ehrenberg 1837 (Bacillariophyceae: Eunotiaceae) from waterbodies of Southeastern Asia. *Nova Hedwigia* Beihefte 147: 69–103
- [17] Grunow A., 1865. Über die von Herrn Gerstenberger in Rabenhorst's Decaden ausgegeben Süßwasser Diatomaceen und Desmidiaceen von der Insel Banka, nebst Untersuchungen über die Gattungen *Ceratoneis* und *Frustulia*. In: *Beiträge zur näheren Kenntniss und Verbreitung der Algen* (Ed. by L., Rabenhorst). Heft II, pp. 1–16. Verlag von Eduard Kummer, Leipzig
- [18] Gusev E.S., & Kulikovskiy M., 2014. Centric diatoms from Vietnam reservoirs with description of one new *Urosolenia* species. *Nova Hedwigia* Beihefte 143: 111–126
- [19] Gutwinski R., 1903. De algis, praecipue diatomaceis a Dre J. Holderer anno 1898: in Asia centrali atque in China collectis. *Bulletin international de l'Académie Polonaise des sciences et des lettres* (Cracovie). Classe des Sciences Mathématiques et Naturelles 1903: 201–227
- [20] Hirano H., 1967. Freshwater algae collected by the joint Thai-Japanese biological expedition to Southeast Asia 1961–62. In: *Nature and life in Southeast Asia* (Ed. by T., Kira & K., Iwata), Vol. 5, pp. 1–71. Flora and Fauna Research Society, Kyoto
- [21] Hustedt F., 1922. Bacillariales aus Innerasien. Gesammelt von Dr. Sven Hedin. In: *Southern Tibet, discoveries in former times compared with my own researches in 1906–1908* (Ed. by S., Hedin), 6, pp. 107–152. Lithographic Institute of the General Staff of the Swedish Army, Stockholm
- [22] Hustedt F., 1927. Bacillariales aus dem Aokikosee in Japan. *Archiv für Hydrobiologie* 18: 155–172
- [23] Hustedt F., 1942. Süßwasser-Diatomeen des indomalayischen archipels und der hawaii-inseln. nach dem material der wallacea-expedition. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 42: 1–252. doi: 10.1002/iroh.19420420102
- [24] Iwahashi Y., 1936. Studies on fresh water diatoms of Western Japan. I. *Journal of Japanese Botany* 1: 390–401
- [25] Iwahashi, Y., 1937. Studies on fresh-water diatoms of western Japan. III. *Journal of Japanese Botany* 13: 360–369
- [26] Jüttner, I., Krammer K., Van de Vijver B., Tuji A., Simkhada B., Gurung S., Sharma S., Sharma C., & Cox E.J., 2010. *Oricymba* (Cymbellales, Bacillariophyceae), a new cymbelloid genus and three new species from the Nepalese Himalaya. *Phycologia* 49: 407–423. doi: 10.2216/09-77.1
- [27] Kapustin D.A., Kulikovskiy M., & Kociolek J.P., 2017. *Celebesia* gen. nov., a new cymbelloid diatom genus from the ancient Lake Matano (Sulawesi Island, Indonesia). *Nova Hedwigia*, Beihefte 146: 147–155. doi: 10.1127/1438-9134/2017/147
- [28] Karthick B., Kociolek J.P., Taylor J.C., & Cocquyt C., 2016. *Gomphonema grande* sp. nov., a new diatom (Bacillariophyta) from the Democratic Republic of the Congo, Tropical Africa. *Phytotaxa* 245: 187–196. doi: 10.11646/phytotaxa.245.3.1
- [29] Kociolek J.P., 1992. Valve ultrastructure and systematic position of *Gomphonema kaznakowi* Mereschkowsky. *Diatom Research* 7: 259–265. doi: 10.1080/0269249X.1992.9705218
- [30] Kociolek J.P., & Kingston J.C., 1999. Taxonomy, ultrastructure, and distribution of some gomphonemoid diatoms (Bacillariophyceae: Gomphonemataceae) from rivers in the United States. *Canadian Journal of Botany* 77: 686–705. doi: 10.1139/b99-007
- [31] Kociolek J.P., & Stoermer E.F., 1990. Diatoms from the Upper Miocene Hot Springs Limestone, Snake River Plain, Idaho (U.S.A.). *Micropaleontology* 36: 331–352. doi: 10.2307/1485474
- [32] Kociolek J.P., & Stoermer E.F., 1993. Freshwater gomphonemoid diatom phylogeny: preliminary results. *Hydrobiologia* 269–270: 31–38. doi: 10.1007/BF00028001
- [33] Kociolek J.P., Kulikovskiy M.S., & Solak C.N., 2013. The diatom genus *Gomphoneis* Cleve (Bacillariophyceae) from Lake Baikal, Russia. *Phytotaxa* 154: 1–37. doi: 10.11646/phytotaxa.154.1.1
- [34] Kociolek J.P., Glushchenko A., & Kulikovskiy M., 2015. Typification, valve ultrastructure, and systematic position of *Gomphonema gomphopleuroides* Amossé ex Kociolek, Glushchenko & Kulikovskiy, an endemic diatom from Southeast Asia. *Diatom Research* 30: 247–255. doi: 10.1080/0269249X.2015.1072583

- [35] Kociolek J.P., Woodward J.C., & Graeff C., 2016a. New and endemic Gomphonema C.G. Ehrenberg (Bacillariophyceae) species from Hawaii. *Nova Hedwigia* 102: 141–171. doi: 10.1127/nova_hedwigia/2015/0296
- [36] Kociolek J.P., You Q., Stepanek J.G., Lowe R.L., & Wang Q., 2016b. A new freshwater diatom genus, *Edtheriotia* gen. nov. of the Stephanodiscaceae Glezer & Makarova (Bacillariophyta) from south-central China. *Phycological Research* 64: 274–280. doi: 10.1111/pre.12145
- [37] Kociolek J.P., Hamsher S.E., Kulikovskiy M., & Bramberger A.J., 2017. Are there species flocks in freshwater diatoms? A review of past reports and a look to the future. *Hydrobiologia* 792: 17–35. doi: 10.1007/s10750-016-3075-1
- [38] Kulikovskiy M.S., & Kociolek J.P., 2014. The diatom genus *Gomphonema* Ehrenberg in Lake Baikal. I. Morphology and taxonomic history of two endemic species. *Nova Hedwigia*, Beiheft 143: 507–518
- [39] Kulikovskiy M.S., & Kuznetsova I.V., 2014. Biogeography of Bacillariophyta. Main concepts and approaches. *International Journal on Algae* 16: 207–228. doi: 10.1615/InterJAlgae.v16.i3.10
- [40] Kulikovskiy M.S., & Kuznetsova I.V., 2016. Morphology, taxonomical position, and distribution of the genera of diatoms *Ochigma* and *Khursevichia* from Lake Baikal. *Inland Water Biology* 9: 226–233. doi: 10.1134/S1995082916030147
- [41] Kulikovskiy M.S., Lange-Bertalot H., Witkowski A., & Khursevich G.K., 2011. *Achnanthidium sibiricum* (Bacillariophyta), a new species from bottom sediments in Lake Baikal. *Algological Studies* 136/137: 77–87. doi: 10.1127/1864-1318/2011/0136-0077
- [42] Kulikovskiy M.S., Lange-Bertalot H., Metzeltin D., & Witkowski A., 2012a. Lake Baikal: hotspot of endemic diatoms I. In: *Iconographia diatomologica. annotated diatom micrographs* (Ed. by H., Lange-Bertalot), Vol. 23, pp. 7–608. A.R.G. Gantner Verlag K.G., Ruggell
- [43] Kulikovskiy M.S., Witkowski A., & Khursevich G.K., 2012b. *Encyonema horstii* sp. nov., a species of unusual valve outline from Pleistocene deposits of Lake Baikal. *Nova Hedwigia*, Beiheft 141: 365–374
- [44] Kulikovskiy M., Lange-Bertalot H., & Witkowski A., 2013. *Gliwiczia* gen. nov., a new achnanthoid diatom genus with description of four species new for science. *Phytotaxa* 109: 1–16. doi: 10.11646/phytotaxa.109.1.1
- [45] Kulikovskiy M.S., Gusev E., Andreeva S., & Annenkova N., 2014a. Phylogenetic position of the diatom genus *Geissleria* Lange-Bertalot & Metzeltin and description of two new species from Siberian mountain lakes. *Phytotaxa* 177: 249–260. doi: 10.11646/phytotaxa.177.5.1
- [46] Kulikovskiy M.S., Lange-Bertalot H., Witkowski A., & Kuznetsova I., 2014b. Description of four species belonging in *Cavinula* D.G. Mann & Stickle from Lake Baikal with notes on family *Cavinulaceae* D.G. Mann in Round et al. 1990. *Nova Hedwigia* 99: 487–499. doi: 10.1127/0029-5035/2014/0196
- [47] Kulikovskiy M., Glushchenko A., & Kociolek J.P., 2015a. The diatom genus *Oricymba* in Vietnam and Laos with description of one new species, and a consideration of its systematic placement. *Phytotaxa* 227: 120–134. doi: 10.11646/phytotaxa.227.2.2
- [48] Kulikovskiy M.S., Kociolek J.P., Solak C.N., & Kuznetsova I., 2015b. The diatom genus *Gomphonema* Ehrenberg in Lake Baikal. II. Revision of taxa from *Gomphonema acuminatum* and *Gomphonema truncatum-capitatum* complexes. *Phytotaxa* 233: 251–272. doi: 10.11646/phytotaxa.233.3.3
- [49] Kulikovskiy M.S., Lange-Bertalot H., & Kuznetsova I.V., 2015c. Lake Baikal: hotspot of endemic diatoms II. In: *Iconographia diatomologica. Annotated diatom micrographs* (Ed. by H., Lange-Bertalot), Vol. 26. Koeltz Scientific Books, Oberreifenberg. 1–657
- [50] Kulikovskiy M., Lange-Bertalot H., Kuznetsova I., & Khursevich G., 2015d. Three new species of *Eolimna* Lange-Bertalot & Schiller (Bacillariophyta) from Lake Baikal. *Nova Hedwigia*, Beiheft 144: 199–209
- [51] Kulikovskiy M., Lange-Bertalot H., Witkowski A., Khursevich G.K., & Kociolek P., 2015e. Description of *Eunotia* species (Bacillariophyta) new to science from Lake Baikal with comments on morphology and biogeography of the genus. *Phycologia* 54: 248–260. doi: 10.2216/14-98.1
- [52] Kulikovskiy M.S., Lange-Bertalot H., Witkowski A., Khursevich G., & Kuznetsova I., 2015f. Typification of diatoms from Lake Baikal. I. Some species described by A.P. Skabitschewsky. *Nova Hedwigia* 100: 215–223. doi: 10.1127/nova_hedwigia/2014/0214
- [53] Kulikovskiy M., Lange-Bertalot H., Annenkova N.V., Gusev E.S., & Kociolek J.P., 2016a. Morphological and molecular evidence support description of two new diatom species from the genus *Ulnaria* in Lake Baikal. *Fottea* 16: 34–42. doi: 10.5507/fot.2015.011
- [54] Kulikovskiy M.S., Lange-Bertalot H., & Kuznetsova I.V., 2016b. *Cocconeis nanoburyatica* sp. nov.–a new monoraphid diatom species from Lake Baikal. *Inland Water Biology* 9: 112–115. doi: 10.1134/S1995082916020103
- [55] Kulikovskiy M., Glushchenko A., Dorofeyuk N., & Kociolek J.P., 2018a. Morphology and distribution of *Ninastrelnikovia laosica* sp. nov.–a new species from the previously monospecific genus. *Nova Hedwigia*, Beiheft 147: 119–126
- [56] Kulikovskiy M., Glushchenko A., Kuznetsova I., & Kociolek J.P., 2018b. Description of the new freshwater diatom genus *Okhapkinia* gen. nov. from Laos (Southeast Asia), with notes on family *Sellaphoraceae* Mereschkowsky 1902. *Fottea* 18: 120–129. doi: 10.5507/fot.2017.021

- [57] Leelahakriengkrai P., Pruetiworanan S., & Peerapornpisal Y., 2009. Diversity of benthic diatoms and macroalgae and water quality in the Mekong River passing Chiang Rai Province, Thailand. *KKU Science Journal* 37: 143-152
- [58] Levkov Z., Mitic-Kopanja D., & Reichardt E., 2016. The diatom genus *Gomphonema* from the Republic of Macedonia. In: *Diatoms of Europe* (Ed. by H., Lange-Bertalot), Vol. 8. Koeltz Botanical Books, Oberreifenberg. 1-552
- [59] Li Y., Xie P., Gong Z., & Shi Z., 2003. Cymbellaceae and Gomphonemataceae (Bacillariophyta) from the Hengduan Mountains region (southwestern China). *Nova Hedwigia* 76: 507-536. doi: 10.1127/0029-5035/2003/0076-0507
- [60] Li Y., Gong Z., Xie P., & Shen J., 2006. Distribution and morphology of two endemic gomphonemoid species, *Gomphonema kaznakowi* Mereschkowsky and *G. yangtzensis* Li nov. sp. in China. *Diatom Research* 21: 313-324. doi: 10.1080/0269249X.2006.9705672
- [61] Li Y., Williams D.M., Metzeltin D., Patrick Kocielek J.P. & Gong Z., 2010. *Tibetiana pulchra* gen. nov. et sp. nov., a new freshwater epilithic diatom from River Nujiang in Tibet, China. *Journal of Phycology* 46: 325-330. doi: 10.1111/j.1529-8817.2009.00776.x
- [62] Li Y., Lange-Bertalot H., & Metzeltin D., 2013. *Sichuaniella* Li Yanling, Lange-Bertalot et Metzeltin nom. nov.-a new name for *Sichuania* Li Yanling et al. In: *Diatoms of Europe* (Ed. by H., Lange-Bertalot), Vol. 7, p. 698. Koeltz Scientific Books, Königstein
- [63] Liu Y., Kocielek J.P., Fan Y., & Wang Q., 2012. *Pseudofallacia* gen. nov., a new freshwater diatom (Bacillariophyceae) genus based on *Navicula occulta* Krasske. *Phycologia* 51: 620-626. doi: 10.2216/11-098.1
- [64] Liu Q., Kocielek J.P., Li B., You Q., & Wang Q., 2017. The diatom genus *Neidium* Pfitzer (Bacillariophyceae) from Zoige Wetland, China. *Bibliotheca Diatomologica* 63: 1-120
- [65] Liu Q., Kocielek J.P., Glushchenko A., Kulikovskiy M., & Fan Y., 2018. A new genus of Eunotiales (Bacillariophyta, Bacillariophyceae: Peroniaceae), *Sinoperonia*, from Southeast Asia, exhibiting remarkable phenotypic plasticity with regard to the raphe system. *Phycologia* 57: 147-158. doi: 10.2216/17-21.1
- [66] Maltsev E., Andreeva S., Kulikovskiy M., Podunaj J., & Kocielek J.P., 2017. Molecular phylogeny of the diatom genus *Envekadea* (Bacillariophyceae, Naviculales). *Nova Hedwigia, Beihefte* 146: 241-252. doi: 10.1127/1438-9134/2017/241
- [67] Meister F., 1913. Beiträge zur Bacillariaceenflora Japans. *Archiv für Hydrobiologie und Planktonkunde* 8: 305-312
- [68] Meister F., 1914. Beiträge zur Bacillariaceenflora Japans. *Archiv für Hydrobiologie und Planktonkunde* 9: 226-232
- [69] Mereschkowsky C., 1906. Diatomées du Tibet. *Imperial Russkoe geograficheskoe obshchestvo*. St. Petersburg. 40 pp
- [70] Metzeltin D., & Lange-Bertalot H., 1998. Tropical diatoms of South America I: about 700 predominantly rarely known or new taxa representative of the neotropical flora. In: *Iconographia diatomologica. Annotated diatom micrographs* (Ed. by H., Lange-Bertalot), Vol. 5. A.R.G. Gantner Verlag K.G., Ruggell. 695 p
- [71] Ohno M., Fukushima H., & Kobayashi T., 1972. Diatom flora of the Mekong water system, Cambodia. *Natural Science* 20: 1-24
- [72] Ohtaka A., Watanabe R., Im S., Chhay R., & Tsukawaki S., 2010. Spatial and seasonal changes of net plankton and zoobenthos in Lake Tonle Sap, Cambodia. *Limnology* 11: 85-94. doi: 10.1007/s10201-009-0283-7
- [73] Prowse G.A., 1962. Diatoms of Malayan freshwaters. *Gardens Bulletin, Singapore* 19: 1-104
- [74] Reichardt E., 2009. Silikataauswüchse an den inneren Stigmenöffnungen bei *Gomphonema*-Arten. *Diatom Research* 24: 159-173. doi: 10.1080/0269249X.2009.9705788
- [75] Shi Z., 1984. *Flora Algarum Sinicarum Aquae Dulcis*. Tomus XII. Bacillariophyta, Gomphonemaceae. Science Press, Beijing. 147p
- [76] Simonsen R., 1987. Atlas and catalogue of the diatom types of Friedrich Hustedt, Vol. 1-3. J. Cramer, Berlin
- [77] Skabitschewsky A.P., 1984. Species *Gomphonematis* Ag. (Bacillariophyta) Lacus Baical. *News of the Systematic of Lower Plants* 21: 51-62
- [78] Skuja H., 1937. Algae. In: *Symbolae Sinicae. Botanische Ergebnisse der expedition der Akademik der Wissenschaften in Wien nach Südwest-China 1914-1918* (Ed. by H., Handel-Mazzetti), 1. J, pp. 1-106. Springer, Vienna
- [79] Skvortzow B.V., 1930: Notes on Ceylon diatoms. I. *The Ceylon Journal of Science, Section A, Botany*, 11: 251-260
- [80] Skvortzow B.V., 1932. Notes on Ceylon diatoms. II. *The Ceylon Journal of Science, Section A, Botany*, 11: 333-338
- [81] Skvortzow B.W., 1937. Bottom diatoms from Olhon Gate of Baikal Lake, Siberia. *Philippine Journal of Science* 62: 293-377

- [82] Skvortzow B.V., 1946. Species novae et minus cognitae Algarum, Flagellatarum et Phycomicetarum Asiae, Africae, Americae et Japoniae nec non Ceylon anno 1931–1945, descripto et illustrato per tab. 1–18. Proceedings of the Harbin Society of Natural History and Ethnography. Botany. Harbin 2: 1–34
- [83] Skvortzow B.W., 1976. Moss diatom flora from River Gan in the northern part of Great Khingan Mountains, China, with description of a new genus *Porosularia* gen. nov. from Northern and Southern China. Quarterly Journal of the Taiwan Museum 29: 111–152
- [84] Skvortzow B.W., & Meyer K.I., 1928. A contribution to the diatoms of Lake Baikal. Proceedings of the Sungari River Biological Station 1: 1–55
- [85] Suphan S., & Peerapornpisal Y., 2009. Scanning electron microscope of benthic diatoms in Mekong River and its tributaries, Thailand. Journal of Microscopy Society of Thailand 23: 42–46
- [86] Suphan S., & Peerapornpisal Y., 2010. Fifty-three new record species of benthic diatoms from Mekong River and its tributaries in Thailand. Chiang Mai Journal of Science 37: 326–343
- [87] Van Heurck H., 1881. Synopsis des Diatomées de Belgique Atlas. pls XXXI–LXXVII. Anvers, Ducaju et Cie
- [88] Vishnyakov V.S., Kulikovskiy M.S., Genkal S.I., Dorofeyuk N.I., Lange-Bertalot H., & Kuznetsova I.V., 2014. Taxonomy and geographical distribution of the diatom genus *Epithemia* Kützing in water bodies of Central Asia. Inland Water Biology 7: 318–330. doi: 10.1134/S199508291404018X
- [89] Vishnyakov V.S., Kulikovskiy M.S., Genkal S.I., & Kuznetsova I.V., 2015. Comparative morphological characteristic of diatoms of genus *Hannaea* Patrick of the two largest lakes of the Baikal rift zone with a description of the new species. Inland Water Biology 8: 222–231. doi: 10.1134/S1995082915030165
- [90] Zhu H., & Chen J., 2000. Bacillariophyta of the Xizang Plateau. Science Press, Beijing. 353 p