

In-vitro antitumor activity of new quaternary phosphonium salts, derivatives of 3-hydroxypyridine

Akhunzyanov A., Pugachev M., Vafiva R., Shtyrlin N., Balakin K., Shtyrlin Y.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Wolters Kluwer Health, Inc. All rights reserved. This work presents the results of in-vitro biological activity studies of three novel anticancer agents, phosphonium salts based on the 3-hydroxypyridine scaffold, including one derivative of 4-deoxypyridoxine. Proliferation and viability of cells treated with these compounds was assessed by the colony formation and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assays. Effects of the compounds on apoptosis and cell cycle were studied by flow cytometry using annexin V-FITC/propidium iodide and propidium iodide staining, respectively. The influence of the compounds on mitochondrial membrane potential and intracellular reactive oxygen species was evaluated using tetramethyl rhodamine ethyl and DCFHA staining. Western blot analysis was used to study the changes in the expression of Bcl-xL, Bax, and caspase-3 apoptotic proteins. The treatment of ovarian adenocarcinoma cells OVCAR-4 with the tested compounds inhibited the growth and induced cell cycle arrest in the G1 phase. 3-Hydroxypyridine derivatives induced apoptosis by hyperexpression of Bax and caspase-3, whereas 4-deoxypyridoxine derivative induced cell death partly by reactive oxygen species generation and caspase-3 hyperexpression. These results indicate that the quaternary phosphonium salts studied represent potential therapeutic agents for the treatment of ovarian cancer.

<http://dx.doi.org/10.1097/CAD.0000000000000642>

Keywords

3-hydroxypyridine, 4-deoxypyridoxine, apoptosis, cell cycle, cytotoxicity, OVCAR-4

References

- [1] Malvezzi M, Carioli G, Rodriguez T, Negri E, La Vecchia C. Global trends and predictions in ovarian cancer mortality. *Ann Oncol* 2016; 27:2017-2025.
- [2] Ferraz R, Costa-Rodrigues J, Fernandes MH, Santos MM, Marrucho IM, Rebelo LPN, et al. Antitumor activity of ionic liquids based on ampicillin. *Chem Med Chem* 2015; 10:1480-1483.
- [3] Millard M, Pathania D, Shabaik Y, Taheri L, Deng J, Neamati N. Preclinical evaluation of novel triphenylphosphonium salts with broad-spectrum activity. *PLoS ONE* 2010; 5:1-18.
- [4] Spivak AY, Nedopekina DA, Khalitova RR, Gubaidullin RR, Odinokov VN, Bel'skii YP, et al. Triphenylphosphonium cations of betulonic acid derivatives: synthesis and antitumor activity. *Med Chem Res* 2017; 26:518-531.
- [5] Kim H, Yang J, Kim MJ, Choi S, Chung JR, Kim JM, et al. Tumor necrosis factor receptor-associated protein 1 (TRAP1) mutation and TRAP1 inhibitor gamitrinib-triphenylphosphonium (G-TPP) induce a forkhead box O (FOXO)-dependent cell protective signal from mitochondria. *J Biol Chem* 2016; 291:1841-1853.

- [6] Son JK, Zhao LX, Basnet A, Thapa P, Karki R, Na Y, et al. Synthesis of 2, 6-diaryl-substituted pyridines and their antitumor activities. *Eur J Med Chem* 2008; 43:675-682.
- [7] Wang GT, Wang X, Wang W, Hasvold LA, Sullivan G, Hutchins CW, et al. Design and synthesis of o-trifluoromethylbiphenyl substituted 2-amino-nicotinonitriles as inhibitors of farnesyltransferase. *Bioorg Med Chem Lett* 2005; 15:153-158.
- [8] Cocco MT, Congiu C, Lilliu V, Onnis V. Synthesis and antiproliferative activity of 2, 6-dibenzylamino-3, 5-dicyanopyridines on human cancer cell lines. *Eur J Med Chem* 2005; 40:1365-1372.
- [9] Radhamani S, Bradley C, Meehan-Andrews T, Ihmaid SK, Al-Rawi J. Radiosensitizing activity of a novel benzoxazine through the promotion of apoptosis and inhibition of DNA repair. *Invest New Drugs* 2014; 32:424-435.
- [10] Shtyrilin NV, Vafina RM, Pugachev MV, Khaziev RM, Nikitina EV, Zeldi MI, et al. Synthesis and biological activity of quaternary phosphonium salts based on 3-hydroxypyridine and 4-deoxyribose. *Russ Chem Bull* 2016; 65:537-545.
- [11] Rittie LFG, Fisher GJ. Isolation and culture of skin fibroblasts. *Fibrosis research: methods and protocols. Methods Mol Med* 2005; 117:83-98.
- [12] Wang J, Yang CT, Kim YS, Sreerama SG, Cao Q, Li ZB, et al. Cu-Labeled triphenylphosphonium and triphenylarsonium cations as highly tumor-selective imaging agents. *J Med Chem* 2007; 50:5057-5069.
- [13] Modica-Napolitano J, Aprile J. Delocalized lipophilic cations selectively target the mitochondria of carcinoma cells. *Adv Drug Deliv Rev* 2001; 49:63-70.
- [14] Evan GI, Vousden KH. Proliferation, cell cycle and apoptosis in cancer. *Nature* 2001; 411:342-348.
- [15] Strasser A, Cory S, Adams JM. Deciphering the rules of programmed cell death to improve therapy of cancer and other diseases. *EMBO J* 2011; 30:3667-3683.
- [16] Gottlieb E, Armour SM, Harris MH, Thompson CB. Mitochondrial membrane potential regulates matrix configuration and cytochrome c release during apoptosis. *Cell Death Differ* 2003; 10:709-717.