

Features of Reaction of 2-(5-Methyl-2-phenyl-2*H*-1,2,3-diazaphosphol-4-yl)-4*H*-benzo[*e*]-1,3,2-dioxaphosphorin-4-one with 1,2-Dicarbonyl Compounds¹

V. F. Mironov^{a,b,*}, G. A. Ivkova^b, L. M. Abdrakhmanova^a,
E. V. Mironova^{a,b}, R. Z. Musin^a, and V. K. Cherkasov^c

^a Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Center of the Russian Academy of Sciences,
ul. Akademika Arbuzova 8, Kazan, Tatarstan, 420088 Russia

*e-mail: mironov@iopc.ru

^b Kazan (Volga Region) Federal University, Kazan, Tatarstan, Russia

^c Razuvaev Institute of Organometallic Chemistry, Russian Academy of Sciences, Nizhny Novgorod, Russia

Received December 21, 2017

Abstract—2-(5-Methyl-2-phenyl-2*H*-1,2,3-diazaphosphol-4-yl)-4*H*-benzo[*e*]-1,3,2-dioxaphosphorin-4-one reacts with perfluorodiacetyl, 3,6-di(*tert*-butyl)-1,2-benzoquinone and phenanthrenequinone only with the participation of a three-coordinated phosphorus atom to form spirophosphoranes containing acyclic 5-methyl-2-phenyl-2*H*-1,2,3-diazaphosphol-4-yl substituent, whereas the interaction with tetrachloro-1,2-benzoquinone proceeds via expanding the six-membered heterocycle to the nine-membered one to form 2-(2-phenyl-2*H*-1,2,3-diazaphosphol-4-yl)-2,9-dioxo-4,5,6,7-tetrachlorodibenzo[*d,h*]-1,3,8-trioxaphosphonine.

Keywords: diazaphosphol, tetrachloro-1,2-benzoquinone, phenanthrenequinone, perfluorodiacetyl, phosphorane


DOI: 10.1134/S107036321803009X

The reaction of 1,2-dicarbonyl compounds with phosphorus(III) derivatives [1–4] is a convenient method for synthesizing derivatives of pentacoordinated phosphorus, important intermediates of phosphorylation and dephosphorylation processes occurring in living cells [5]. On the basis of this reaction, a variety of phospholenes **1** with a pentacoordinated phosphorus atom has been obtained (Scheme 1), the features of their molecular structure have been studied, and the

main directions of their chemical transformations into the more stable derivatives of tetracoordinated phosphorus have been determined [6–8].

Among the unusual chemical properties of phospholenes **1** the oxaphosphorane condensation should be mentioned [9, 10]. Recently, it has been shown [11] that the P-pentacoordinated products of the [4+1]-cycloaddition of 3,5-di(*tert*-butyl)-1,2-benzoquinone with

Scheme 1.



R, R¹ = Alk.

¹ Supporting materials are available from authors.