

Hydrothermal transformations of organic matter of low permeability rocks from domanic formation of the romashkino oil field

Kayukova G., Mikhailova A., Kosachev I., Pronin N., Sotnikov O., Evdokimov A., Khisamov R.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018, © 2018 Taylor & Francis Group, LLC. The oil generating potential of Domanic rocks from Dankov-Lebedyan horizon of the Zelenogorsk area of Romashkino oil field was evaluated by Rock-Eval pyrolysis technique. The result of given method depends on the content, composition, and thermal stability of organic matter in rocks. During hydrothermal processes, the distinctive conversion behavior of organic matter at temperatures of 200°C, 250°C, 300°C, and 350°C in CO₂environment was revealed. The yield of obtained aquathermolysis products and their quality were evaluated. The results of the studies suggest that low-permeability carbonate rocks of the Dankov-Lebedyan horizon contain productive beds with content of Corg 1.89–3.03%, which when developed using thermal methods, can become an additional source of liquid hydrocarbons.

<http://dx.doi.org/10.1080/10916466.2018.1490764>

Keywords

carbonate rock, domanic deposit, hydrothermal conversions, kerogen, organic matter, shale oil

References

- [1] Bushnev, D. A., and N. S., Burdel'naya. 2013. Modeling of oil generation by domanik carbonaceous shale. *Petroleum Chemistry* 53 (3):145–51. doi:10.1134/S096554411303002X.
- [2] Disnar, J. R., B., Guillet, D., Keravis, C., Di-Giovanni, and D., Sebag. 2003. Soil organic matter (SOM) characterization by rock-eval pyrolysis: Scope and limitations. *Organic Geochemistry* 34 (3):327–43. doi:10.1016/S0146-6380(02)00239-5.
- [3] Feoktistov, D. A., G. P., Kayukova, A. V., Vakhin, and S. A., Sitnov. 2018. Catalytic aquathermolysis of high-viscosity oil using iron, cobalt and copper tallowates. *Chemistry and Technology of Fuels and Oils* 53 (6):905–12. doi:10.1007/s10553-018-0880-4.
- [4] Ivanova, A. G., A. V., Vakhin, E. V., Voronina, A. V., Pyataev, D. K., Nurgaliev, and S. A., Sitnov. 2017. Mössbauer study of products of the thermocatalytic treatment of kerogen-containing rocks. *Bulletin of the Russian Academy of Sciences: Physics* 81 (7):817–21. doi:10.3103/S1062873817070139.
- [5] Kayukova, G. P., A. M., Kiyamova, A. N., Mikhailova, I. P., Kosachev, S. M., Petrov, G. V., Romanov, L. M., Sildikova, I. N., Plotnikova, and A. V., Vakhin. 2016. Generation of hydrocarbons by hydrothermal transformation of organic matter of domanik rocks. *Chemistry and Technology of Fuels and Oils* 52 (2):149–61. doi:10.1007/s10553-016-0685-2.
- [6] Kayukova, G. P., A. M., Mikhailova, D. A., Feoktistov, V. P., Morozov, and A. V., Vakhin. 2017. Conversion of the organic matter of domanic shale and permian bituminous rocks in hydrothermal catalytic processes. *Energy & Fuels* 31 (8):7789–99. doi:10.1021/acs.energyfuels.7b00612.

- [7] Khisamov, R. S., V. G., Bazarevskaya, T. I., Tarasova, N. A., Badurtdinova, and O. G., Gibadullina. 2017. Hydrocarbon potential of domanic pay zones of volga-ural oil-and-gas basin. *Neftyanoe Khozyaystvo - Oil Industry* (6):10-4. doi:10.24887/0028-2448-2017-6-10-14.
- [8] Kontorovich, A. E., V. N., Melenevskii, Y. N., Zanin, A. G., Zamirailova, V. A., Kazanenkov, V. Y., Kazarbin, E. N., Makhneva, and L. S., Yamkovaya. 1998. Lithology, organic geochemistry, and formation conditions of basic rocks in the bazhenovka formation (west siberia.). *Geologiya i Geofizika* 39 (11):1477-91.
- [9] Kontorovich, A. E., V. I., Moskvin, O. I., Bostrikov, V. P., Danilova, A. N., Fomin, A. S., Fomichev, E. A., Kostyрева, and V. N., Melenevsky. 1997. Main oil source formations of the west siberian basin. *Petroleum Geoscience* 3 (4):343-58. doi:10.1144/petgeo.3.4.343.
- [10] Petrov, S. M., D. A., Ibragimova, A. G., Safiulina, B., Tohidi, A. V., Vakhin, R. C., Okekwe, and E. A., Karalin. 2017. Conversion of organic matter in the carbonaceous medium in the supercritical water. *Journal of Petroleum Science and Engineering* 159:497-505. doi:10.1016/j.petrol.2017.09.060.
- [11] Skibitskaya, N. A., Somov, Yu, P., Faizrakhmanov, R. R., Domanova, E. G., Batalin, O., Yu, and Mishuta M. N., 2008. Recovery of heavy organics from gas-bearing sedimentary rocks. *Solid Fuel Chemistry* 42 (3):165-70. doi:10.3103/S0361521908030099.
- [12] Tissot, B. P., and D. H., Welte. 1984. *Petroleum formation and occurrence*. 2nd ed., 151. Berlin, Germany: Springer Verlag.
- [13] Vakhin, A. V., Y. V., Onishchenko, N. A., Nazimov, and R. U., Kadyrov. 2017. Thermal transformation of the mobile-hydrocarbon composition of domanik deposits of volga-ural oil-and gas-bearing province. *Chemistry and Technology of Fuels and Oils* 53 (4):511-9. doi:10.1007/s10553-017-0830-6.
- [14] Varfolomeev, M. A., R. N., Nagrimanov, A. A., Samatov, I. T., Rakipov, A. D., Nikanshin, A. V., Vakhin, D. K., Nurgaliev, and M. V., Kok. 2016. Chemical evaluation and kinetics of siberian, North regions of russia and republic of Tatarstan crude oils. *Energy Sources, Part A: Recovery, Utilization and Environmental Effects* 38 (8):1031-8. doi:10.1080/15567036.2015.1107866.
- [15] Wilson, M. A., D. E., Lambert, and P. J., Collin. 1985. Chemical transformations during pyrolysis of rundle oil shale. *Fuel* 64 (12):1647-54. doi:10.1016/0016-2361(85)90388-6.
- [16] Yang, F., T., Wang, and M., Li. 2017. Geochemistry characteristics of high-mature source rock extracts and using various approaches to discriminate the indigenity. *Petroleum Science and Technology* 35 (12):1296-303. doi:10.1080/10916466.2017.1327968.