

Sellaphora balashovae (Bacillariophyta), a new species from siberian mountain lake frolikha (Baikal region), Russia

Andreeva S., Kocolek J., Maltsev E., Dorofeyuk N., Kezlya E., Shkurina N., Kuznetsova I., Gusev E., Kulikovskiy M.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Magnolia Press. A new species, *Sellaphora balashovae* sp. nov., from Lake Frolikha is described based on light (LM) and scanning electron microscopy (SEM). The systematic position of this new species is determined based on molecular data. This species is placed in the genus *Sellaphora* based on chloroplast morphology, areolae covered by external hymenes, and uniserial striae that are typical for the genus. Molecular data support this taxonomic assignment, and the new species is part of a monophyletic group with some other members of the genus *Sellaphora*. The new species is distinguishable from other *Sellaphora* species by its small size, valve shape, striae and areolae densities, striae orientation and distinctly large areolae bordering the sternum.

<http://dx.doi.org/10.11646/phytotaxa.371.2.1>

Keywords

Diatoms, Morphology, Phylogeny, S: baikal region, Sellaphoraceae

References

- [1] Alverson, A.J., Jansen, R.K. & Theriot, E.C. (2007) Bridging the Rubicon: Phylogenetic analysis reveals repeated colonization's of marine and fresh waters by thalassiosiroid diatoms. *Molecular Phylogenetics and Evolution* 45:193-210. <https://doi.org/10.1016/j.ympev.2007.03.024>
- [2] Cleve, P.T. & Grunow, A. (1880) Beiträge zur Kenntniss der Arctischen Diatomeen. *Kongliga Svenska-Vetenskaps Akademiens Handlingar* 17 (2): 1-121.
- [3] Cleve, P.T. & Möller, J.D. (1882) Diatoms. Part VI, Esatas Edquist's Boktryckeri, Upsala, 48 pp. [pp. 277-324]
- [4] Enache, M.D. & Potapova, M. (2009) A new species of *Sellaphora* (Sellaphoraceae) from Hannaberry Lake, Arkansas, U.S.A. *Acta Botanica Croatica* 68: 231-237.
- [5] Evans, K.M., Wortley, A.H. & Mann, D.G. (2007) An assessment of potential diatom "barcode" genes (cox1, rbcL, 18S and ITS rDNA) and their effectiveness in determining relationships in *Sellaphora* (Bacillariophyta). *Protist* 158: 349-364. <https://doi.org/10.1016/j.protis.2007.04.001>
- [6] Evans, K.M., Wortley, A.H., Simpson, G.E., Chepurnov, V.A. & Mann, D.G. (2008) A molecular systematic approach to explore diversity within the *Sellaphora pupula* species complex (Bacillariophyta). *Journal of Phycology* 44: 215-231. <https://doi.org/10.1111/j.1529-8817.2007.00454.x>
- [7] Guillard, R.R.L., Carpenter, E.J. & Reimann, B.E.F. (1974) *Skeletonema menzelii* sp. nov., a new diatom from the western Atlantic Ocean. *Phycologia* 13 (2):131-138. <https://doi.org/10.2216/i0031-8884-13-2-131.1>
- [8] Guillard, R.R.L. & Lorenzen, C.J. (1972) Yellow-green algae with chlorophyllide c. *Journal of Phycology* 8: 10-14.

- [9] Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. (2010) New algorithms and methods to estimate Maximum-Likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* 59 (3): 307-321. <https://doi.org/10.1093/sysbio/syq010>
- [10] Gusev, E.S. & Kulikovskiy, M.S. (2013) A new species of the genus *Mallomonas* (Chrysophyceae: Synurales), *Mallomonas kuzminii* sp. nov., from lake Frolikha (Russia, Baikal region). *Phytotaxa* 155 (1): 66-70. <https://doi.org/10.11646/phytotaxa.155.1.6>
- [11] Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95-98.
- [12] Katoh, K. & Toh, H. (2010) Parallelization of the MAFFT multiple sequence alignment program. *Bioinformatics* 26: 1899-1900. <https://doi.org/10.1093/bioinformatics/btq224>
- [13] Kociolek, J.P., Balasubramanian, K., Blanco, S., Coste, M., Ector, L., Liu, Y., Kulikovskiy, M., Lundholm, N., Ludwig, T., Potapova, M., Rimet, F., Sabbe, K., Sala, S., Sar, E., Taylor, J., Van de Vijver, B., Wetzel, C.E., Williams, D.M., Witkowski, A. & Witkowski, J. (2017) DiatomBase. Available from: <http://diatombase.org/> (accessed 12 November 2017)
- [14] Kociolek, J.P., Hamsher, S.E., Kulikovskiy, M. & Bramberger, A.S. (2017) Are there species flocks in freshwater diatoms? A review of past reports and a look to the future. *Hydrobiologia* 792: 17-35. <https://doi.org/10.1007/s10750-016-3075-1>
- [15] Kociolek, J.P., Kulikovskiy, M., Kuznetsova, I., Glushchenko, A.M., Solak, C.N. (2018) A putative species flock in the diatom genus *Gomphonema* Ehrenberg (Bacillariophyta: Gomphonemataceae) from Lake Baikal, Russia: description of six new species similar to *G. ventricosum* W. Gregory. *Cryptogamie, Algologie* 39 (3): 365-388. <https://doi.org/10.7872/crya/v39.iss3.2018.365>
- [16] Kociolek, J.P., Kulikovskiy, M.S. & Solak, C.N. (2013) The diatom genus *Gomphoneis* Cleve (Bacillariophyceae) from Lake Baikal, Russia. *Phytotaxa* 154 (1): 1-37. <https://doi.org/10.11646/phytotaxa.154.1.1>
- [17] Kozhov, M.M. (1950) Fresh waters of East Siberia. Irkutskoe oblastnoye izdatelstvo, Irkutsk, 367 pp.
- [18] Kulikovskiy, M., Glushchenko, A., Kuznetsova, I. & Kociolek, J.P. (2018) Description of the new freshwater diatom genus *Okhaphinia* gen. nov. from Laos (Southeast Asia), with notes on family Sellaphoraceae Mereschkowsky 1902. *Fottea* 1: 120-129. <https://doi.org/10.5507/fot.2017.021>
- [19] Kulikovskiy, M.S., Gusev, E.S., Andreeva, S.A. & Annenkova, N.V. (2014a) Phylogenetic position of the diatom genus *Geissleria* Lange-Bertalot & Metzeltin and description of two new species from Siberian mountain lakes. *Phytotaxa* 177 (5): 249-260. <https://doi.org/10.11646/phytotaxa.177.5.1>
- [20] Kulikovskiy, M.S. & Kociolek, J.P. (2014) The diatom genus *Gomphonema* Ehrenberg in Lake Baikal. I. Morphology and taxonomic history of two endemic species. *Nova Hedwigia, Beiheft* 143: 507-518. <https://doi.org/10.1127/1438-9134/2014/027>
- [21] Kulikovskiy, M.S., Kociolek, J.P., Solak, C.N. & Kuznetsova, I.V. (2015a) The diatom genus *Gomphonema* Ehrenberg in Lake Baikal. II. Revision of taxa from *Gomphonema acuminatum* and *Gomphonema truncatum-capitatum* complexes. *Phytotaxa* 233 (3): 251-272. <https://doi.org/10.11646/phytotaxa.233.3.3>
- [22] Kulikovskiy, M.S. & Kuznetsova, I.V. (2014) Biogeography of Bacillariophyta. Main concepts and approaches. *International Journal on Algae* 16 (3): 207-228. <https://doi.org/10.1615/IJterAlgae.v16.i3.10>
- [23] Kulikovskiy, M.S. & Kuznetsova, I.V. (2016) Morphology, taxonomical position, and distribution of the genera of diatoms *Ochigma* and *Khursevichia* from Lake Baikal. *Inland Water Biology* 9 (3): 226-233. <https://doi.org/10.1134/S1995082916030147>
- [24] Kulikovskiy, M.S., Lange-Bertalot, H., Annenkova, N.V., Gusev, E.S. & Kociolek, J.P. (2016a) Morphological and molecular evidence support description of two new diatom species from the genus *Ulnaria* in Lake Baikal. *Fottea* 16: 34-42. <https://doi.org/10.5507/fot.2015.011>
- [25] Kulikovskiy, M.S., Lange-Bertalot, H. & Kuznetsova, I.V. (2016b) *Cocconeis nanoburyatica* sp. nov.—a new monoraphid diatom species from Lake Baikal. *Inland Water Biology* 9 (2): 112-115. <https://doi.org/10.1134/S1995082916020103>
- [26] Kulikovskiy, M.S., Lange-Bertalot, H., Kuznetsova, I.V. & Khursevich, G.K. (2015b) Three new species of *Eolimna* Lange-Bertalot & Schiller (Bacillariophyta) from Lake Baikal. *Nova Hedwigia, Beiheft* 144: 199-209. https://doi.org/10.1127/nova_supp1/2015/0040
- [27] Kulikovskiy, M.S., Lange-Bertalot, H., Metzeltin, D. & Witkowski, A. (2012a) Lake Baikal: hotspot of endemic diatoms I. *Iconographia Diatomologica* 23: 1-608.
- [28] Kulikovskiy, M.S., Lange-Bertalot, H. & Witkowski, A. (2013) *Gliwiczia* gen. nov., a new achanthoid diatom genus with description of four species new for science. *Phytotaxa* 109 (1): 1-16. <https://doi.org/10.11646/phytotaxa.109.1.1>
- [29] Kulikovskiy, M.S., Lange-Bertalot, H., Witkowski, A. & Khursevich, G.K. (2011) *Achnanthidium sibiricum* (Bacillariophyta), a new species from bottom sediments in Lake Baikal. *Algological Studies* 136/137: 77-87. <https://doi.org/10.1127/1864-1318/2011/0136-0077>

- [30] Kulikovskiy, M.S., Lange-Bertalot, H., Witkowski, A., Khursevich, G.K. & Kociolek, J.P. (2015c) Description of Eunotia species (Bacillariophyta) new to science from Lake Baikal with comments on morphology and biogeography of the genus. *Phycologia* 54 (3): 248-260. <https://doi.org/10.2216/14-98.1>
- [31] Kulikovskiy, M.S., Lange-Bertalot, H., Witkowski, A., Khursevich, G.K. & Kuznetsova, I.V. (2015d) Typification of diatoms from Lake Baikal. I. Some species described by A.P. Skabitschewsky. *Nova Hedwigia* 100 (1-2): 215-223. https://doi.org/10.1127/nova_hedwigia/2014/0214
- [32] Kulikovskiy, M.S., Lange-Bertalot, H., Witkowski, A. & Kuznetsova, I.V. (2014b) Description of four species belonging in Cavinula D.G. Mann & Stickle from Lake Baikal with notes on family Cavinulaceae D.G. Mann in Round et al. 1990. *Nova Hedwigia* 99 (3-4): 487-499. <https://doi.org/10.1127/0029-5035/2014/0196>
- [33] Kulikovskiy, M.S., Witkowski, A. & Khursevich, G.K. (2012b) Encyonema horstii sp. nov., a species of unusual valve outline from Pleistocene deposits of Lake Baikal. *Nova Hedwigia*, Beiheft 141: 365-374.
- [34] Kuzmina, A.E., Ignatova, N.V. & Mizandronzev, I.B. (2004) Species composition, ecological and geographical characteristics of the diatoms from Lake Frolikha. Proceedings of the international symposium "Living cells of the diatoms" (Irkutsk) 1: 55.
- [35] Lange-Bertalot, H., Cavacini, P., Tagliaventi, N. & Alfinito, S. (2003) Diatoms of Sardinia. Rare and 76 new species in rock pools and other ephemeral waters. *Iconographia Diatomologica* 12: 1-438.
- [36] Lange-Bertalot, H., Küblst, K., Lauser, T., Nörpel-Schempp, M. & Willmann, M. (1996) Diatom Taxa introduced by Georg Krasske Documentation and Revision. *Iconographia Diatomologica* 3: 1-358.
- [37] Lange-Bertalot, H. & Moser, G. (1994) Brachysira. Monographie der Gattung. *Bibliotheca Diatomologica* 29: 1-212.
- [38] Levkov, Z., Krstić, S., Metzeltin, D. & Nakov, T. (2007) Diatoms of lakes Prespa and Ohrid. About 500 taxa from ancient lake system. *Iconographia Diatomologica* 16: 1-611.
- [39] Mann, D.G. (1989) The Diatom genus Sellaphora: separation from Navicula. *British Phycological Journal* 24 (1): 1-20. <https://doi.org/10.1080/00071618900650011>
- [40] Mann, D.G., Thomas, S.J. & Evans, K.M. (2008) Revision of the diatom genus Sellaphora: a first account of the larger species in the British Isles. *Fottea* 8 (1): 15-78. <https://doi.org/10.5507/fot.2008.002>
- [41] Mann, D.G., Evans, K.M., Chepurnov, V.A. & Nagai, S. (2009) Morphology and formal description of Sellaphora bisexualis sp. nov. (Bacillariophyta). *Fottea* 9: 199-209. <https://doi.org/10.5507/fot.2009.021>
- [42] Mann, D.G., McDonald, S.M., Bayer, M.M., Droop, S.J.M., Chepurnov, V.A., Loke, R.E., Ciobanu, A. & du Buf, J.M.H. (2004) Morphometric analysis, ultrastructure and mating data provide evidence for five new species of Sellaphora (Bacillariophyceae). *Phycologia* 43: 459-482. <https://doi.org/10.2216/I0031-8884-43-4-459.1>
- [43] Mereschkowsky, C. (1902) On Sellaphora, a new genus of diatoms. *Annals and Magazine of Natural History* 7-9: 185-195. <https://doi.org/10.1080/00222930208678565>
- [44] Moser, G., Lange-Bertalot, H. & Metzeltin, D. (1998) Insel der Endemiten Geobotanisches Phänomen Neukaledonien (Island of endemics New Caledonia—a geobotanical phenomenon). *Bibliotheca Diatomologica* 38: 1-464.
- [45] Podzorski, A.C. (1984) Caponea caribbea Podorski, a structurally unique new diatom from Jamaica. *Nova Hedwigia* 40: 1-8.
- [46] Posada, D. (2006) Modeltest Server: a web-based tool for the statistical selection of models of nucleotide substitution online. *Nucleic Acids Research* 34: 700-703. <https://doi.org/10.1093/nar/gkl042>
- [47] Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572- 1574. <https://doi.org/10.1093/bioinformatics/btg180>
- [48] Round, F.E., Crawford, R.M. & Mann, D.G. (1990) The diatoms. Biology and morphology of the genera. Cambridge University Press, 747 pp.
- [49] Ruck, E.C. & Theriot, E.C. (2011) Origin and evolution of the canal raphe system in diatoms. *Protist* 162: 723-737. <https://doi.org/10.1016/j.protis.2011.02.003>
- [50] Schiller, W. & Lange-Bertalot, H. (1997) Eolimna martinii n. gen., n. sp. (Bacillariophyceae) aus dem Unter-Oligozän von Sieblos/Rhon im Vergleich mit ähnlichen rezenten Taxa. *Paläontologische Zeitschrift* 71 (3): 163-172. <https://doi.org/10.1007/BF02988485>
- [51] Skabitschewsky, A.P. (1953) Phytoplankton and diatoms of Lake Frolikha (Transbaikalia). Proceedings of the Irkutsk State University. Biological Series 1: 145-152.
- [52] Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30: 2725-2729. <https://doi.org/10.1093/molbev/mst197>
- [53] Timoshkin, O.A. (2001) Lake Baikal: diversity of fauna, problems of its immiscibility and origin, ecology and "exotic" communities. - Index of animal species inhabiting Lake Baikal and its catchment area. *Lake Baikal* 1 (1): 16-73.
- [54] Van de Vijver, B., Frenot, Y. & Beyens, L. (2002) Freshwater diatoms from Ile de la Possession (Crozet archipelago, subantarctica). *Bibliotheca Diatomologica* 46: 1-412.

- [55] Voigt, M. (1960) Some New Diatoms from the Far East. *Journal of the Royal Microscopical Society* 78: 92-94.
<https://doi.org/10.1111/j.1365-2818.1958.tb02048.x>
- [56] Werum, M. & Lange-Bertalot, H. (2004) Diatoms in springs from Central Europe and elsewhere under the influence of hydrogeology and anthropogenic impacts. *Iconographia Diatomologica* 13: 3-417.
- [57] Wetzel, C.E., Ector, L., Van de Vijver, B., Compère, P. & Mann, D.G. (2015) Morphology, typification and critical analysis of some ecologically important small naviculoid species (Bacillariophyta). *Fottea* 15 (2): 203-234.
<https://doi.org/10.5507/fot.2015.020>
- [58] Zimmermann, J., Jahn, R. & Gemeinholzer, B. (2011) Barcoding diatoms: evaluation of the V4 subregion on the 18S rRNA gene, including new primers and protocols. *Organisms Diversity and Evolution* 11: 173-192.
<https://doi.org/10.1007/s13127-011-0050-6>