

Palynological analysis of bottom sediments of lake rubskoe (Ivanovo region, Russia)

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2018. The results of the palynological analysis of the sediment core recovered in Lake Rubskoe (Ivanovo Region, Russia) in 2015 are presented. An environmental reconstruction of Holocene and late Pleistocene vegetation, climate and lake dynamic was inferred. Pollen analysis has shown that vegetation of the area was dominated by *Betula pendula*, *Pinus* and *Alnaster*. Herbaceous were mostly represented by *Artemisia*, *Chenopodiaceae*. *Sphagnum* spores finding in the in the upper layers of core indicates a significant moisture content of the region and a warm climate, which explains the development of peat bogs in the study area. Four pollen zones have been identified in the diagram. Bottom of the core (zone PZ I) was characterized of low pollen concentrations. A decrease of forest vegetation and an increase of dry steppes vegetation were indicated of the last phase of the late Pleistocene. In upper layers of the core was presented a higher concentration of arboreal pollen. The concentration of herbs a simultaneous decrease. Zone PZ II corresponds to the late Dryas Pleistocene and early Holocene. The slight increase of *Betula pendula* pollen coincided with Allerod warming. Atlantic, boreal and preboreal periods (zone PZ III) characterized by a predominance of *Betula pendula* and *Pinus* pollen. The development of broad-leaved trees indicates the climatic optimum of the Holocene. Pollen zone IV coincided with the late Holocene. Here warming and cooling of the Subboreal and Subatlantic chronozone were noticeable.

<http://dx.doi.org/10.5593/sgem2018/5.1/S20.081>

Keywords

Lake Rubskoe, Palynological analysis, Pollen

References

- [1] Nigamatzyanova G., Frolova L. Specific structural features of zooplankton of polygonal pond Lena River Delta, Russia, Arctic, 17th International multidisciplinary scientific geoconference SGEM, Ecology, economics education and legislation, Issue 51, vol. 17, 2017, p. 891-898.
- [2] Frolova L.A., Ibragimova A.G., Fedorova I.V. Stratigraphy of Cladocera in a core from Yamal Peninsula Lake (Arctic Russia), 16th International Multidisciplinary Scientific GeoConference SGEM, book4, vol. 2, 2016, p. 579-586.
- [3] Frolova L.A., Frolova A.A. Cladocera from bottom deposits as an indicator of changes in climate and ecological conditions, IOP Conf. Series: Earth and Environmental Science, vol. 107, issue 1, 201, no 012084.
- [4] Borisova E.A., Shilov M.P., Markov D.S., Melnikov V.N., Tikhomirov A.M., Mineeva L.Y., Barinov S.N., Lasareva O.G., Chudnenko D.E. Water bodies located in Ivanovo region specially protected natural areas, issue 1, 2013, p. 88.

- [5] Grichuk V.P. An attempt to reconstruct some elements of the climate of the northern hemisphere in the Atlantic period of the Holocene. In M. I. Neishtadt, ed., *The Holocene*, 1969, pp. 41-57.
- [6] Kupriyanova L.A., Alyoshina L.A. Pollen and Spores of Plants from the Flora of European Part of USSR, vol. I, 1972, p.171.
- [7] Kupriyanova L.A., Alyoshina L.A. Pollen and Spores of Plants from the Flora of European Part of USSR, vol. II, 1978, p. 183.
- [8] Reille M. Pollen et spores d'Europe et d'Afrique du nord. *Laboratoire de botanique historique et palynologie*, 1992, p. 520.
- [9] Reille M. Pollen et spores d'Europe et d'Afrique du nord Supplement 1.-*Laboratoire de botanique historique et palynologie*, 1995, p. 520.
- [10] Grimm E. *Tilia software 2.0.2*. Illinois State Museum Research and Collection Center, Springfield, 2004.
- [11] Velichko A.A. *The natural process in the Pleistocene*, 1973, p. 256.
- [12] Khotinsky N.A. *Holocene of Northern Eurasia*, 1977, p. 200.
- [13] Novenko E.Yu., Velichko A.A., Suganova I.S., Jung F.W., Boettger T. Dynamiks of vegetation at the Late Pleistocene glacial-interglacial transition (new data from the center of the East European Plain), *Polish Geological Institute Special Papers, Proceedings of the Workshop Reconstruction of Quaternary palaeoclimate and palaeoenvironments and their abrupt changes*, 2005, 16, p. 77-82.
- [14] Velichko A.A., Novenko E.Y., Pisareva V.V., Zelikson E.M., Boettger T., Jung F.W. Vegetation and climate changes during the Eemian interglacial in Central and Eastern Europe: comparative analysis of pollen data, *Boreas*, 34(2), p. 207-219.
- [15] Payne R.J, Malysheva E., Tsyganov A., Pampura T., Novenko E., Elena Volkova, Babeshko K., Mazei Yu. A multi-proxy record of Holocene environmental change, peatland development and carbon accumulation from Staroselsky Moch peatland, Russia, *The Holocene*, 2016, vol. 26 (2), p. 314-326.