

Dynamics of the hypoxia—induced tissue edema in the rat barrel cortex in vitro

Juzekaeva E., Gainutdinov A., Mukhtarov M., Khazipov R.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Juzekaeva, Gainutdinov, Mukhtarov and Khazipov. Cerebral edema is a major, life threatening complication of ischemic brain damage. Previous studies using brain slices have revealed that cellular swelling and a concomitant increase in tissue transparency starts within minutes of the onset of metabolic insult in association with collective anoxic spreading depolarization (aSD). However, the dynamics of tissue swelling in brain slices under ischemia-like conditions remain elusive. Here, we explored the dynamics of brain tissue swelling induced by oxygen-glucose deprivation (OGD) in submerged rat barrel cortex slices. Video monitoring of the vertical and horizontal position of fluorescent dye-filled neurons and contrast slice surface imaging revealed elevation of the slice surface and a horizontal displacement of the cortical tissue during OGD. The OGD-induced tissue movement was also associated with an expansion of the slice borders. Tissue swelling started several minutes after aSD and continued during reperfusion with normal solution. Thirty minutes after aSD, slice borders had expanded by ~130 μm and the slice surface had moved up to attain a height of ~70 μm above control levels, which corresponded to a volume increase of ~30%. Hyperosmotic sucrose solution partially reduced the OGD-induced slice swelling. Thus, OGD-induced cortical slice tissue swelling in brain slices in vitro recapitulates many features of ischemic cerebral edema in vivo, its onset is tightly linked to aSD and it develops at a relatively slow pace after aSD. We propose that this model of cerebral edema in vitro could be useful for the exploration of the pathophysiological mechanisms underlying ischemic cerebral edema and in the search for an efficient treatment to this devastating condition.

<http://dx.doi.org/10.3389/fncel.2018.00502>

Keywords

Brain ischemia, Cortex, Edema, Hypoxia, Swelling

References

- [1] Andrew R.D., Labron M. W., Boehnke S. E., Carnduff L., Kirov S. A., (2007). Physiological evidence that pyramidal neurons lack functional water channels. *Cereb. Cortex* 17, 787–802. 10.1093/cercor/bhk03216723408.
- [2] Balestrino M., Somjen G. G., (1986). Chlorpromazine protects brain tissue in hypoxia by delaying spreading depression-mediated calcium influx. *Brain Res.* 385, 219–226. 10.1016/0006-8993(86)91067-X3022872.
- [3] Brisson C.D., Andrew R., (2012). A neuronal population in hypothalamus that dramatically resists acute ischemic injury compared to neocortex. *J. Neurophysiol.* 108, 419–430. 10.1152/jn.00090.201222514289.

- [4] Charpak S., Audinat E., (1998). Cardiac arrest in rodents: maximal duration compatible with a recovery of neuronal activity. *Proc Natl Acad Sci U.S.A.* 95, 4748–4753. 10.1073/pnas.95.8.47489539810.
- [5] Croning M. D., Haddad G. G., (1998). Comparison of brain slice chamber designs for investigations of oxygen deprivation in vitro. *J. Neurosci. Methods* 81, 103–111. 10.1016/S0165-0270(98)00023-59696315.
- [6] Dostovic Z., Dostovic E., Smajlovic D., Ibrahimagic O. C., Avdic L., (2016). Brain edema after ischaemic stroke. *Med. Arch.* 70, 339–341. 10.5455/medarh.2016.70.339-34127994292.
- [7] Dreier J. P., Lemale C. L., Kola V., Friedman A., Schoknecht K., (2018). Spreading depolarization is not an epiphenomenon but the principal mechanism of the cytotoxic edema in various gray matter structures of the brain during stroke. *Neuropharmacology* 134, 189–207. 10.1016/j.neuropharm.2017.09.027.
- [8] Dugue G. P., Dumoulin A., Triller A., Dieudonne S., (2005). Target-dependent use of co-released inhibitory transmitters at central synapses. *J. Neurosci.* 25, 6490–6498. 10.1523/JNEUROSCI.1500-05.200516014710.
- [9] Dzhala V., Khalilov I., Ben Ari Y., Khazipov R., (2001). Neuronal mechanisms of the anoxia-induced network oscillations in the rat hippocampus in vitro. *J. Physiol.* 536, 521–531. 10.1111/j.1469-7793.2001.0521c.xd11600686.
- [10] Elkin B. S., Shaik M. A., Morrison B., (2010). Fixed negative charge and the Donnan effect: a description of the driving forces associated with brain tissue swelling and oedema. *Philos. Trans. A Math. Phys. Eng. Sci.* 368, 585–603. 10.1098/rsta.2009.022320047940.
- [11] Frank J. I., (1995). Large hemispheric infarction, deterioration, and intracranial pressure. *Neurology* 45, 1286–1290. 10.1212/WNL.45.7.12867617183.
- [12] Joshi I., Andrew R. D., (2001). Imaging anoxic depolarization during ischemia-like conditions in the mouse hemi-brain slice. *J. Neurophysiol.* 85, 414–424. 10.1152/jn.2001.85.1.41411152742.
- [13] Juzekaeva E., Nasretidinov A., Gainutdinov A., Sintsov M., Mukhtarov M., Khazipov R., (2017). Preferential initiation and spread of anoxic depolarization in layer 4 of rat barrel cortex. *Front. Cell Neurosci.* 11:390 10.3389/fncel.2017.0039029326550.
- [14] Khazipov R., Zaynutdinova D., Ogievetsky E., Valeeva G., Mitrukina O., Manent J. B., (2015). Atlas of the postnatal rat brain in stereotaxic coordinates. *Front. Neuroanat.* 9:161. 10.3389/fnana.2015.0016126778970.
- [15] Klatzo I., (1987). Pathophysiological aspects of brain edema. *Acta Neuropathologica* 72, 236–239. 10.1007/BF006910953564903.
- [16] Kumar V., Naik R. S., Hillert M., Klein J., (2006). Effects of chloride flux modulators in an in vitro model of brain edema formation. *Brain Res.* 1122, 222–229. 10.1016/j.brainres.2006.09.01217014830.
- [17] MacGregor D. G., Avshalumov M. V., Rice M. E., (2003). Brain edema induced by in vitro ischemia: causal factors and neuroprotection. *J. Neurochem.* 85, 1402–1411. 10.1046/j.1471-4159.2003.01772.x12787060.
- [18] Murphy T. H., Li P., Betts K., Liu R., (2008). Two-photon imaging of stroke onset in vivo reveals that NMDA-receptor independent ischemic depolarization is the major cause of rapid reversible damage to dendrites and spines. *J. Neurosci.* 28, 1756–1772. 10.1523/JNEUROSCI.5128-07.200818272696.
- [19] Obeidat A. S., Jarvis C. R., Andrew R. D., (2000). Glutamate does not mediate acute neuronal damage after spreading depression induced by O-2/glucose deprivation in the hippocampal slice. *J. Cereb. Blood Flow Metabol.* 20, 412–422. 10.1097/00004647-200002000-0002410698080.
- [20] Rader R. K., Lanthorn T. H., (1989). Experimental ischemia induces a persistent depolarization blocked by decreased calcium and NMDA antagonists. *Neurosci. Lett.* 99, 125–130. 10.1016/0304-3940(89)9027-02546106.
- [21] Risher W., Ard D., Yuan J., Kirov S. A., (2010). Recurrent spontaneous spreading depolarizations facilitate acute dendritic injury in the ischemic penumbra. *J. Neurosci.* 30, 9859–9868. 10.1523/JNEUROSCI.1917-10.201020660268.
- [22] Risher W., Croom D., Kirov S. A., (2012). Persistent astroglial swelling accompanies rapid reversible dendritic injury during stroke-induced spreading depolarizations. *Glia* 60, 1709–1720. 10.1002/glia.2239022821441.
- [23] Risher W.C., Andrew R.D., Kirov S. A., (2009). Real-time passive volume responses of astrocytes to acute osmotic and ischemic stress in cortical slices and in vivo revealed by two-photon microscopy. *Glia* 57, 207–221. 10.1002/glia.2074718720409.
- [24] Schwab S., Aschoff A., Spranger M., Albert F., Hacke W., (1996). The value of intracranial pressure monitoring in acute hemispheric stroke. *Neurology* 47, 393–398. 10.1212/WNL.47.2.3938757010.
- [25] Somjen G. G., (2001). Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol. Rev.* 81, 1065–1096. 10.1152/physrev.2001.81.3.106511427692.
- [26] Stokum J. A., Gerzanich V., Simard J. M., (2016). Molecular pathophysiology of cerebral edema. *J. Cereb. Blood Flow Metab.* 36, 513–538. 10.1177/0271678X1561717226661240.
- [27] Tanaka E., Yamamoto S., Kudo Y., Mihara S., Higashi H., (1997). Mechanisms underlying the rapid depolarization produced by deprivation of oxygen and glucose in rat hippocampal CA1 neurons in vitro. *J. Neurophysiol.* 78, 891–902. 10.1152/jn.1997.78.2.8919307122.

- [28] Vanharreveld A., (1957). Changes in volume of cortical neuronal elements during asphyxiation. *Am. J. Physiol.* 191, 233-242. 10.1152/ajplegacy.1957.191.2.233.
- [29] Young J. N., Aitken P. G., Somjen G. G., (1991). Calcium, magnesium, and long-term recovery from hypoxia in hippocampal tissue slices. *Brain Res.* 548, 343-345. 10.1016/0006-8993(91)91146-R1868344.
- [30] Zheng H., Chen C., Zhang J., Hu Z., (2016). Mechanism and therapy of brain edema after intracerebral hemorrhage. *Cerebrovasc. Dis.* 42, 155-169. 10.1159/00044517027110940.