

c-Jun N-terminal kinases (JNKs) in myocardial and cerebral ischemia/reperfusion injury

Shvedova M., Anfinogenova Y., Atochina-Vasserman E., Schepetkin I., Atochin D.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Shvedova, Anfinogenova, Atochina-Vasserman, Schepetkin and Atochin. In this article, we review the literature regarding the role of c-Jun N-terminal kinases (JNKs) in cerebral and myocardial ischemia/reperfusion injury. Numerous studies demonstrate that JNK-mediated signaling pathways play an essential role in cerebral and myocardial ischemia/reperfusion injury. JNK-associated mechanisms are involved in preconditioning and post-conditioning of the heart and the brain. The literature and our own studies suggest that JNK inhibitors may exert cardioprotective and neuroprotective properties. The effects of modulating the JNK-dependent pathways in the brain and the heart are reviewed. Cardioprotective and neuroprotective mechanisms of JNK inhibitors are discussed in detail including synthetic small molecule inhibitors (AS601245, SP600125, IQ-1S, and SR-3306), ion channel inhibitor GsMTx4, JNK-interacting proteins, inhibitors of mixed-lineage kinase (MLK) and MLK-interacting proteins, inhibitors of glutamate receptors, nitric oxide (NO) donors, and anesthetics. The role of JNKs in ischemia/reperfusion injury of the heart in diabetes mellitus is discussed in the context of comorbidities. According to reviewed literature, JNKs represent promising therapeutic targets for protection of the brain and the heart against ischemic stroke and myocardial infarction, respectively. However, different members of the JNK family exert diverse physiological properties which may not allow for systemic administration of non-specific JNK inhibitors for therapeutic purposes. Currently available candidate JNK inhibitors with high therapeutic potential are identified. The further search for selective JNK3 inhibitors remains an important task.

<http://dx.doi.org/10.3389/fphar.2018.00715>

Keywords

Brain, C-Jun-N-terminal kinase, Heart, Ischemia/reperfusion injury, JNK inhibitor, Stroke

References

- [1] Aoki, H., Kang, P. M., Hampe, J., Yoshimura, K., Noma, T., Matsuzaki, M., et al. (2002). Direct activation of mitochondrial apoptosis machinery by c-Jun N-terminal kinase in adult cardiac myocytes. *J. Biol. Chem.* 277, 10244-10250. doi: 10.1074/jbc.M112355200
- [2] Armstrong, S. C. (2004). Protein kinase activation and myocardial ischemia-reperfusion injury. *Cardiovasc. Res.* 61, 427-436. doi: 10.1016/j.cardiores.2003.09.031
- [3] Atochin, D. N., Schepetkin, I. A., Khlebnikov, A. I., Seledtsov, V. I., Swanson, H., Quinn, M. T., et al. (2016). A novel dual NO-donating oxime and c-Jun N-terminal kinase inhibitor protects against cerebral ischemia-reperfusion injury in mice. *Neurosci. Lett.* 618, 45-49. doi: 10.1016/j.neulet.2016.02.033

- [4] Bahnasi, Y. M., Wright, H. M., Milligan, C. J., Dedman, A. M., Zeng, F., Hopkins, P. M., et al. (2008). Modulation of TRPC5 cation channels by halothane, chloroform and propofol. *Br. J. Pharmacol.* 153, 1505-1512. doi: 10.1038/sj.bjp.0707689
- [5] Bain, J., McLauchlan, H., Elliott, M., and Cohen, P. (2003). The specificities of protein kinase inhibitors: an update. *Biochem. J.* 371(Pt 1), 199-204. doi: 10.1042/BJ20021535
- [6] Barr, R. K., Boehm, I., Attwood, P. V., Watt, P. M., and Bogoyevitch, M. A. (2004). The critical features and the mechanism of inhibition of a kinase interaction motif-based peptide inhibitor of JNK. *J. Biol. Chem.* 279, 36327-36338. doi: 10.1074/jbc.M402181200
- [7] Barthwal, M. K., Sathyanarayana, P., Kundu, C. N., Rana, B., Pradeep, A., Sharma, C., et al. (2003). Negative regulation of mixed lineage kinase 3 by Akt/PKB leads to cell survival. *J. Biol. Chem.* 278, 3897-3902. doi: 10.1074/jbc.M211598200
- [8] Bedin, M., Gaben, A. M., Saucier, C., and Mester, J. (2004). Geldanamycin, an inhibitor of the chaperone activity of HSP90, induces MAPK-independent cell cycle arrest. *Int. J. Cancer* 109, 643-652. doi: 10.1002/ijc.20010
- [9] Beeler, N., Riederer, B. M., Waeber, G., and Abderrahmani, A. (2009). Role of the JNK-interacting protein 1/islet brain 1 in cell degeneration in Alzheimer disease and diabetes. *Brain Res. Bull.* 80, 274-281. doi: 10.1016/j.brainresbull.2009.07.006
- [10] Bode, A. M., and Dong, Z. (2007). The functional contrariety of JNK. *Mol. Carcinog.* 46, 591-598. doi: 10.1002/mc.20348
- [11] Bogoyevitch, M. A., and Kobe, B. (2006). Uses for JNK: the many and varied substrates of the c-Jun N-terminal kinases. *Microbiol. Mol. Biol. Rev.* 70, 1061-1095. doi: 10.1128/MMBR.00025-06
- [12] Borsello, T., Clarke, P. G., Hirt, L., Vercelli, A., Repici, M., Schorderet, D. F., et al. (2003). A peptide inhibitor of c-Jun N-terminal kinase protects against excitotoxicity and cerebral ischemia. *Nat. Med.* 9, 1180-1186. doi: 10.1038/nm911
- [13] Bozyczko-Coyne, D., O'Kane, T. M., Wu, Z. L., Dobrzanski, P., Murthy, S., Vaught, J. L., et al. (2001). CEP-1347/KT-7515, an inhibitor of SAPK/JNK pathway activation, promotes survival and blocks multiple events associated with A β -induced cortical neuron apoptosis. *J. Neurochem.* 77, 849-863. doi: 10.1046/j.1471-4159.2001.00294.x
- [14] Brecht, S., Kirchhof, R., Chromik, A., Willesen, M., Nicolaus, T., Raivich, G., et al. (2005). Specific pathophysiological functions of JNK isoforms in the brain. *Eur. J. Neurosci.* 21, 363-377. doi: 10.1111/j.1460-9568.2005.03857.x
- [15] Carboni, S., Antonsson, B., Gaillard, P., Gotteland, J. P., Gillon, J. Y., and Vitte, P. A. (2005). Control of death receptor and mitochondrial-dependent apoptosis by c-Jun N-terminal kinase in hippocampal CA1 neurons following global transient ischaemia. *J. Neurochem.* 92, 1054-1060. doi: 10.1111/j.1471-4159.2004.02925.x
- [16] Carboni, S., Boschert, U., Gaillard, P., Gotteland, J. P., Gillon, J. Y., et al. (2008). AS601245, a c-Jun NH2-terminal kinase (JNK) inhibitor, reduces axon/dendrite damage and cognitive deficits after global cerebral ischaemia in gerbils. *Br. J. Pharmacol.* 153, 157-163. doi: 10.1038/sj.bjp.0707574
- [17] Carboni, S., Hiver, A., Szyndralewicz, C., Gaillard, P., Gotteland, J. P., and Vitte, P. A. (2004). AS601245 (1, 3-benzothiazol-2-yl (2-[[2-(3-pyridinyl) ethyl] amino]-4 pyrimidinyl) acetonitrile): a c-Jun NH2-terminal protein kinase inhibitor with neuroprotective properties. *J. Pharmacol. Exp. Ther.* 310, 25-32. doi: 10.1124/jpet.103.064246
- [18] Carlsson, Y., Leverin, A. L., Hedtj rn, M., Wang, X., Mallard, C., and Hagberg, H. (2009). Role of mixed lineage kinase inhibition in neonatal hypoxia-ischemia. *Dev. Neurosci.* 31, 420-426. doi: 10.1159/000232560
- [19] Chambers, J. W., Pachori, A., Howard, S., Iqbal, S., and LoGrasso, P. V. (2013). Inhibition of JNK mitochondrial localization and signaling is protective against ischemia-reperfusion injury in rats. *J. Biol. Chem.* 288, 4000-4011. doi: 10.1074/jbc.M112.406777
- [20] Chaudhury, H., Zakkar, M., Boyle, J., Cuhlmann, S., van der Heiden, K., Luong le, A., et al. (2010). c-Jun N-terminal kinase primes endothelial cells at atheroprone sites for apoptosis. *Arterioscler. Thromb. Vasc. Biol.* 30, 546-553. doi: 10.1161/ATVBAHA.109.201368
- [21] Chen, Q., Xu, T., Li, D., Pan, D., Wu, P., Luo, Y., et al. (2016). JNK/PI3K/Akt signaling pathway is involved in myocardial ischemia-reperfusion injury in diabetic rats: effects of salvianolic acid A intervention. *Am. J. Transl. Res.* 8, 2534-2548
- [22] Chen, Z., Zhang, X., Liu, Y., and Liu, Z. (2016). Morphine postconditioning protects against reperfusion injury via inhibiting JNK/p38 MAPK and mitochondrial permeability transition pores signaling pathways. *Cell Physiol. Biochem.* 39, 61-70. doi: 10.1159/000445605
- [23] Clerk, A., Fuller, S. J., Michael, A., and Sugden, P. H. (1998). Stimulation of "stress-regulated" mitogen-activated protein kinases (stress-activated protein kinases/c-Jun N-terminal kinases and p38-mitogen-activated protein kinases) in perfused rat hearts by oxidative and other stresses. *J. Biol. Chem.* 273, 7228-7234. doi: 10.1074/jbc.273.13.7228

- [24] Cornett, P. M., Matta, J. A., and Ahern, G. P. (2008). General anesthetics sensitize the capsaicin receptor transient receptor potential V1. *Mol. Pharmacol.* 74, 1261-1268. doi: 10.1124/mol.108.049684
- [25] Davis, R. J. (2000). Signal transduction by the JNK group of MAP kinases. *Cell.* 103, 239-252. doi: 10.1016/S0092-8674(00)00116-1
- [26] Deloche, C., Lopez-Lazaro, L., Mouz, S., Perino, J., Abadie, C., and Combette, J. M. (2014). XG-102 administered to healthy male volunteers as a single intravenous infusion: a randomized, double-blind, placebo-controlled, dose-escalating study. *Pharmacol. Res. Perspect.* 2:e00020. doi: 10.1002/prp2.20
- [27] Di, J. H., Li, C., Yu, H. M., Zheng, J. N., and Zhang, G. Y. (2012). nNOS downregulation attenuates neuronal apoptosis by inhibiting nNOS-GluR6 interaction and GluR6 nitrosylation in cerebral ischemic reperfusion. *Biochem. Biophys. Res. Commun.* 420, 594-599. doi: 10.1016/j.bbrc.2012.03.039
- [28] Diaz-Cañestro, C., Merlini, M., Bonetti, N. R., Liberale, L., Wüst, P., Briand-Schumacher, S., et al. (2018). Sirtuin 5 as a novel target to blunt blood-brain barrier damage induced by cerebral ischemia/reperfusion injury. *Int. J. Cardiol.* 260, 148-155. doi: 10.1016/j.ijcard.2017.12.060
- [29] Dougherty, C. J., Kubasiak, L. A., Frazier, D. P., Li, H., Xiong, W. C., et al. (2004). Mitochondrial signals initiate the activation of c-Jun N-terminal kinase (JNK) by hypoxia-reoxygenation. *FASEB J.* 18, 1060-1070. doi: 10.1096/fj.04-1505com
- [30] Dougherty, C. J., Kubasiak, L. A., Prentice, H., Andreka, P., Bishopric, N. H., and Webster, K. A. (2002). Activation of c-Jun N-terminal kinase promotes survival of cardiac myocytes after oxidative stress. *Biochem. J.* 362(Pt 3), 561-571. doi: 10.1042/bj3620561
- [31] Du, Y., Li, C., Hu, W. W., Song, Y. J., and Zhang, G. Y. (2009). Neuroprotection of preconditioning against ischemic brain injury in rat hippocampus through inhibition of the assembly of GluR6-PSD95-mixed lineage kinase 3 signaling module via nuclear and non-nuclear pathways. *Neuroscience* 161, 370-380. doi: 10.1016/j.neuroscience.2009.03.050
- [32] Duplain, H. (2006). Salvage of ischemic myocardium: a focus on JNK. *Adv. Exp. Med. Biol.* 588, 157-164. doi: 10.1007/978-0-387-34817-9_14
- [33] Engelbrecht, A. M., Niesler, C., Page, C., and Lochner, A. (2004). p38 and JNK have distinct regulatory functions on the development of apoptosis during simulated ischaemia and reperfusion in neonatal cardiomyocytes. *Basic Res. Cardiol.* 99, 338-350. doi: 10.1007/s00395-004-0478-3
- [34] Erikson, J. M., Valente, A. J., Mummidi, S., Kandikattu, H. K., DeMarco, V. G., Bender, S. B., et al. (2017). Targeting TRAF3IP2 by genetic and interventional approaches inhibits ischemia/reperfusion-induced myocardial injury and adverse remodeling. *J. Biol. Chem.* 292, 2345-2358. doi: 10.1074/jbc.M116.764522
- [35] Farrokhnia, N., Roos, M. W., Terént, A., and Lennmyr, F. (2005). Differential early mitogen-activated protein kinase activation in hyperglycemic ischemic brain injury in the rat. *Eur. J. Clin. Invest.* 35, 457-463. doi: 10.1111/j.1365-2362.2005.01512.x
- [36] Ferrandi, C., Ballerio, R., Gaillard, P., Giachetti, C., Carboni, S., Vitte, P. A., et al. (2004). Inhibition of c-Jun N-terminal kinase decreases cardiomyocyte apoptosis and infarct size after myocardial ischemia and reperfusion in anaesthetized rats. *Br. J. Pharmacol.* 142, 953-960. doi: 10.1038/sj.bjp.0705873
- [37] Ferrer, I., Friguls, B., Dalfó, E., and Planas, A. M. (2003). Early modifications in the expression of mitogen-activated protein kinase (MAPK/ERK), stress-activated kinases SAPK/JNK and p38, and their phosphorylated substrates following focal cerebral ischemia. *Acta Neuropathol.* 105, 425-437. doi: 10.1007/s00401-002-061-2
- [38] Frazier, D. P., Wilson, A., Dougherty, C. J., Li, H., Bishopric, N. H., and Webster, K. A. (2007). PKC- α and TAK-1 are intermediates in the activation of c-Jun NH2-terminal kinase by hypoxia-reoxygenation. *Am. J. Physiol. Heart Circ. Physiol.* 292, H1675-H1684. doi: 10.1152/ajpheart.01132.2006
- [39] Fryer, R. M., Patel, H. H., Hsu, A. K., and Gross, G. J. (2001). Stress-activated protein kinase phosphorylation during cardioprotection in the ischemic myocardium. *Am. J. Physiol. Heart. Circ. Physiol.* 281, H1184-H1192. doi: 10.1152/ajpheart.2001.281.3.H1184
- [40] Gao, Y., Signore, A. P., Yin, W., Cao, G., Yin, X. M., Sun, F., et al. (2005). Neuroprotection against focal ischemic brain injury by inhibition of c-Jun N-terminal kinase and attenuation of the mitochondrial apoptosis-signaling pathway. *J. Cereb. Blood Flow Metab.* 25, 694-712. doi: 10.1038/sj.jcbfm.9600062
- [41] Gehringer, M., Muth, F., Koch, P., and Laufer, S. A. (2015). c-Jun N-terminal kinase inhibitors: a patent review (2010-2014). *Expert Opin. Ther. Pat.* 25, 849-872. doi: 10.1517/13543776.2015.1039984
- [42] Girn, H. R., Ahilathirunayagam, S., Mavor, A. I., and Homer-Vanniasinkam, S. (2007). Reperfusion syndrome: cellular mechanisms of microvascular dysfunction and potential therapeutic strategies. *Vasc. Endovascular Surg.* 41, 277-293. doi: 10.1177/1538574407304510
- [43] Guan, Q. H., Pei, D. S., Liu, X. M., Wang, X. T., Xu, T. L., and Zhang, G. Y. (2006). Neuroprotection against ischemic brain injury by SP600125 via suppressing the extrinsic and intrinsic pathways of apoptosis. *Brain Res.* 1092, 36-46. doi: 10.1016/j.brainres.2006.03.086

- [44] Guan, Q. H., Pei, D. S., Zhang, Q. G., Hao, Z. B., Xu, T. L., and Zhang, G. Y. (2005). The neuroprotective action of SP600125, a new inhibitor of JNK, on transient brain ischemia/reperfusion-induced neuronal death in rat hippocampal CA1 via nuclear and non-nuclear pathways. *Brain Res.* 1035, 51-59. doi: 10.1016/j.brainres.2004.11.050
- [45] Guo, X. X., An, S., Yang, Y., Liu, Y., Hao, Q., Tang, T., et al. (2018). Emerging role of the Jun N-terminal kinase interactome in human health. *Cell Biol. Int.* 42, 756-768. doi: 10.1002/cbin.10948
- [46] Gupta, S., Barrett, T., Whitmarsh, A. J., Cavanagh, J., Sluss, H. K., Dérijard, B., et al. (1996). Selective interaction of JNK protein kinase isoforms with transcription factors. *EMBO J.* 15, 2760-2770
- [47] Hausenloy, D. J., and Yellon, D. M. (2006). Survival kinases in ischemic preconditioning and postconditioning. *Cardiovasc. Res.* 70, 240-253. doi: 10.1016/j.cardiores.2006.01.017
- [48] Hausenloy, D. J., and Yellon, D. M. (2007). Preconditioning and postconditioning: united at reperfusion. *Pharmacol. Ther.* 116, 173-191. doi: 10.1016/j.pharmthera.2007.06.005
- [49] Hayashi, T., Sakai, K., Sasaki, C., Zhang, W. R., Warita, H., and Abe, K. (2000). c-Jun N-terminal kinase (JNK) and JNK interacting protein response in rat brain after transient middle cerebral artery occlusion. *Neurosci. Lett.* 284, 195-199. doi: 10.1016/S0304-3940(00)01024-7
- [50] He, H., Li, H. L., Lin, A., and Gottlieb, R. A. (1999). Activation of the JNK pathway is important for cardiomyocyte death in response to simulated ischemia. *Cell Death Differ.* 6, 987-991. doi: 10.1038/sj.cdd.4400572
- [51] Hreniuk, D., Garay, M., Gaarde, W., Monia, B. P., McKay, R. A., and Cioffi, C. L. (2001). Inhibition of c-Jun N-terminal kinase 1, but not c-Jun N-terminal kinase 2, suppresses apoptosis induced by ischemia/reoxygenation in rat cardiac myocytes. *Mol. Pharmacol.* 59, 867-874. doi: 10.1124/mol.59.4.867
- [52] Hu, M., Ye, P., Liao, H., Chen, M., and Yang, F. (2016). Metformin protects H9C2 cardiomyocytes from high-glucose and hypoxia/reoxygenation injury via inhibition of reactive oxygen species generation and inflammatory responses: role of AMPK and JNK. *J. Diabetes Res.* 2016:2961954. doi: 10.1155/2016/2961954
- [53] Hu, S. Q., Ye, J. S., Zong, Y. Y., Sun, C. C., Liu, D. H., Wu, Y. P., et al. (2012). S-nitrosylation of mixed lineage kinase 3 contributes to its activation after cerebral ischemia. *J. Biol. Chem.* 287, 2364-2377. doi: 10.1074/jbc.M111.227124
- [54] Hu, W. W., Du, Y., Li, C., Song, Y. J., and Zhang, G. Y. (2008). Neuroprotection of hypothermia against neuronal death in rat hippocampus through inhibiting the increased assembly of GluR6-PSD95-MLK3 signaling module induced by cerebral ischemia/reperfusion. *Hippocampus* 18, 386-397. doi: 10.1002/hipo.20402
- [55] Ip, Y. T., and Davis, R. J. (1998). Signal transduction by the c-Jun N-terminal kinase (JNK)-from inflammation to development. *Curr. Opin. Cell Biol.* 10, 205-219. doi: 10.1016/S0955-0674(98)80143-9
- [56] Irving, E. A., and Bamford, M. (2002). Role of mitogen- and stress-activated kinases in ischemic injury. *J. Cereb. Blood Flow Metab.* 22, 631-647. doi: 10.1097/00004647-200206000-00001
- [57] Jang, S., and Javadov, S. (2014). Inhibition of JNK aggravates the recovery of rat hearts after global ischemia: the role of mitochondrial JNK. *PLoS ONE* 9:e113526. doi: 10.1371/journal.pone.0113526
- [58] Javadov, S., Jang, S., and Agostini, B. (2014). Crosstalk between mitogen-activated protein kinases and mitochondria in cardiac diseases: therapeutic perspectives. *Pharmacol. Ther.* 144, 202-225. doi: 10.1016/j.pharmthera.2014.05.013
- [59] Ji, F. T., Liang, J. J., Miao, L. P., Wu, Q., and Cao, M. H. (2015). Propofol post-conditioning protects the blood brain barrier by decreasing matrix metalloproteinase-9 and aquaporin-4 expression and improves the neurobehavioral outcome in a rat model of focal cerebral ischemia-reperfusion injury. *Mol. Med. Rep.* 12, 2049-2055. doi: 10.3892/mmr.2015.3585
- [60] Ji, Y., Teng, L., Zhang, R., Sun, J., and Guo, Y. (2017). NRG-1 β exerts neuroprotective effects against ischemia reperfusion-induced injury in rats through the JNK signaling pathway. *Neuroscience* 362, 13-24. doi: 10.1016/j.neuroscience.2017.08.032
- [61] Jin, Q., Li, R., Hu, N., Xin, T., Zhu, P., Hu, S., et al. (2017). DUSP1 alleviates cardiac ischemia-reperfusion injury by suppressing the Mff-required mitochondrial fission and Bnip3-related mitophagy via the JNK pathways. *Redox Biol.* 14, 576-587. doi: 10.1016/j.redox.2017.11.004
- [62] Johnson, G. L., and Nakamura, K. (2007). The c-jun kinase/stress-activated pathway: regulation, function and role in human disease. *Biochim. Biophys. Acta* 1773, 1341-1348. doi: 10.1016/j.bbamcr.2006.12.009
- [63] Kaiser, R. A., Liang, Q., Bueno, O., Huang, Y., Lackey, T., Klevitsky, R., et al. (2005). Genetic inhibition or activation of JNK1/2 protects the myocardium from ischemia-reperfusion-induced cell death in vivo. *J. Biol. Chem.* 280, 32602-32608. doi: 10.1074/jbc.M500684200
- [64] Kaminsky, V. O., and Zhivotovsky, B. (2014). Free radicals in cross talk between autophagy and apoptosis. *Antioxid. Redox Signal.* 21, 86-102. doi: 10.1089/ars.2013.5746
- [65] Kase, H., Iwahashi, K., and Matsuda, Y. (1986). K-252a, a potent inhibitor of protein kinase C from microbial origin. *J. Antibiot.* 39, 1059-1065. doi: 10.7164/antibiotics.39.1059
- [66] Keyse, S. M. (2008). Dual-specificity MAP kinase phosphatases (MKPs) and cancer. *Cancer Metastasis Rev.* 27, 253-261. doi: 10.1007/s10555-008-9123-1

- [67] Khalid, S., Drasche, A., Thurner, M., Hermann, M., Ashraf, M. I., Fresser, F., et al. (2016). cJun N-terminal kinase (JNK) phosphorylation of serine 36 is critical for p66Shc activation. *Sci. Rep.* 6:20930. doi: 10.1038/srep20930
- [68] Khan, S. I., Malhotra, R. K., Rani, N., Sahu, A. K., Tomar, A., Garg, S., et al. (2017). Febuxostat modulates MAPK/NF- κ Bp65/TNF- α signaling in cardiac ischemia-reperfusion injury. *Oxid. Med. Cell Longev.* 2017:8095825. doi: 10.1155/2017/8095825
- [69] Khandoudi, N., Delerive, P., Berrebi-Bertrand, I., Buckingham, R. E., Staels, B., and Bril, A. (2002). Rosiglitazone, a peroxisome proliferator-activated receptor-gamma, inhibits the Jun NH(2)-terminal kinase/activating protein 1 pathway and protects the heart from ischemia-reperfusion injury. *Diabetes* 51, 1507-1514. doi: 10.2337/diabetes.51.5.1507
- [70] Kim, A. H., Khursigara, g., Sun, X., Franke, T. F., and Chao, M. V. (2001). Akt phosphorylates and negatively regulates apoptosis signal-regulating kinase 1. *Mol. Cell. Biol.* 21, 893-901. doi: 10.1128/MCB.21.3.893-901.2001
- [71] Kim, S., Jeong, J., Jung, H. S., Kim, B., Kim, Y. E., et al. (2017). Anti-inflammatory effect of glucagon like peptide-1 receptor agonist, exendin-4, through modulation of IB1/JIP1 expression and JNK signaling in stroke. *Exp. Neurobiol.* 26, 227-239. doi: 10.5607/en.2017.26.4.227
- [72] Knight, R. J., and Buxton, D. B. (1996). Stimulation of c-Jun kinase and mitogen-activated protein kinase by ischemia and reperfusion in the perfused rat heart. *Biochem. Biophys. Res. Commun.* 218, 83-88. doi: 10.1006/bbrc.1996.0016
- [73] Koch, P., Gehringer, M., and Laufer, S. A. (2015). Inhibitors of c-Jun N-terminal kinases: an update. *J. Med. Chem.* 58, 72-95. doi: 10.1021/jm501212r
- [74] Koga, S., Kojima, S., Kishimoto, T., Kuwabara, S., and Yamaguchi, A. (2012). Over-expression of map kinase phosphatase-1 (MKP-1) suppresses neuronal death through regulating JNK signaling in hypoxia/reoxygenation. *Brain Res.* 1436, 137-146. doi: 10.1016/j.brainres.2011.12.004
- [75] Krenitsky, V. P., Delgado, M., Nadolny, L., Sahasrabudhe, K., Ayala, L., Clareen, S. S., et al. (2012). Aminopurine based JNK inhibitors for the prevention of ischemia reperfusion injury. *Bioorg. Med. Chem. Lett.* 22, 1427-1432. doi: 10.1016/j.bmcl.2011.12.028
- [76] Kuan, C. Y., and Burke, R. E. (2005). Targeting the JNK signaling pathway for stroke and Parkinson's diseases therapy. *Curr. Drug Targets CNS Neurol. Disord.* 4, 63-67. doi: 10.2174/1568007053005145
- [77] Kuan, C. Y., Whitmarsh, A. J., Yang, D. D., Liao, G., Schloemer, A. J., Dong, C., et al. (2003). A critical role of neural-specific JNK3 for ischemic apoptosis. *Proc. Natl. Acad. Sci. U S A.* 100, 15184-15189. doi: 10.1073/pnas.2336254100
- [78] Laderoute, K. R., and Webster, K. A. (1997). Hypoxia/reoxygenation stimulates Jun kinase activity through redox signaling in cardiac myocytes. *Circ. Res.* 80, 336-344. doi: 10.1161/01.RES.80.3.336
- [79] Li, D., Li, X., Wu, J., Li, J., Zhang, L., Xiong, T., et al. (2015). Involvement of the JNK/FOXO3a/Bim pathway in neuronal apoptosis after hypoxic-ischemic brain damage in neonatal rats. *PLoS ONE* 10:e0132998. doi: 10.1371/journal.pone.0132998
- [80] Li, H. H., Du, J., Fan, Y. N., Zhang, M. L., Liu, D. P., Li, L., et al. (2011). The ubiquitin ligase MuRF1 protects against cardiac ischemia-reperfusion injury by its proteasome-dependent degradation of phospho-c-Jun. *Am. J. Pathol.* 178, 1043-1058. doi: 10.1016/j.ajpath.2010.11.049
- [81] Li, L., Xu, T., Du, Y., Pan, D., Wu, W., Zhu, H., et al. (2016). Salvianolic acid A attenuates cell apoptosis, oxidative stress, Akt and NF- κ B activation in angiotensin-II induced murine peritoneal macrophages. *Curr. Pharm. Biotechnol.* 17, 283-290. doi: 10.2174/138920101703160206150535
- [82] Li, X. M., Ma, Y. T., Yang, Y. N., Zhang, J. F., Chen, B. D., Liu, F., et al. (2009). Ischemic postconditioning protects hypertrophic myocardium by ERK1/2 signaling pathway: experiment with mice. *Chin. Med. J.* 89, 846-850
- [83] Liu, D. H., Yuan, F. G., Hu, S. Q., Diao, F., Wu, Y. P., Zong, Y. Y., et al. (2013). Endogenous nitric oxide induces activation of apoptosis signal-regulating kinase 1 via S-nitrosylation in rat hippocampus during cerebral ischemia-reperfusion. *Neuroscience* 229, 36-48. doi: 10.1016/j.neuroscience.2012.10.055
- [84] Liu, H. T., Zhang, H. F., Si, R., Zhang, Q. J., Zhang, K. R., Guo, W. Y., et al. (2007). Insulin protects isolated hearts from ischemia-reperfusion injury: cross-talk between PI3-K/Akt and JNKs. *Acta Physiol. Sin.* 59, 651-659
- [85] Liu, X. H., Zhang, Z. Y., Sun, S., and Wu, X. D. (2008). Ischemic postconditioning protects myocardium from ischemia-reperfusion injury through attenuating endoplasmic reticulum stress. *Shock* 30, 422-427. doi: 10.1097/SHK.0b013e318164ca29
- [86] Liu, X., Xu, F., Fu, Y., Liu, F., Sun, S., and Wu, X. (2006). Calreticulin induces delayed cardioprotection through mitogen-activated protein kinases. *Proteomics* 6, 3792-3800. doi: 10.1002/pmic.200500906
- [87] Liu, Y., Huang, Y., Xu, Y., Qu, P., and Wang, M. (2018). Memantine protects against ischemia/reperfusion-induced brain endothelial permeability. *IUBMB Life* 70, 336-343. doi: 10.1002/iub.1729
- [88] Luo, F., Shi, J., Shi, Q., Xu, X., Xia, Y., and He, X. (2016). Mitogen-activated protein kinases and hypoxic/ischemic nephropathy. *Cell Physiol. Biochem.* 39, 1051-1067. doi: 10.1159/000447812

- [89] Luo, S. Y., Li, R., Le, Z. Y., Li, Q. L., and Chen, Z. W. (2017). Anfibatide protects against rat cerebral ischemia-reperfusion injury via TLR4/JNK/caspase-3 pathway. *Eur. J. Pharmacol.* 807, 127-137. doi: 10.1016/j.ejphar.2017.04.002
- [90] Ma, Y., Sun, S., Zhang, J., Chen, Z., Guo, F., Du, Y., et al. (2016). Phosphorylation of JNK increases in the cortex of rat subjected to diabetic cerebral ischemia. *Neurochem. Res.* 41, 787-794. doi: 10.1007/s11064-015-173-4
- [91] Maroney, A. C., Glicksman, M. A., Basma, A. N., Walton, K. M., Knight, E. Jr., Murphy, C. A., et al. (1998). Motoneuron apoptosis is blocked by CEP-1347 (KT 7515), a novel inhibitor of the JNK signaling pathway. *J. Neurosci.* 18, 104-111. doi: 10.1523/JNEUROSCI.18-01-00104.1998
- [92] Maslov, L. N., Mrochek, A. G., Schepetkin, I. A., Headrick, J. P., Hanus, L., Barzakh, E. I., et al. (2013). Protein kinases role in adaptive phenomenon of heart ischemic postconditioning development. *Russ. J. Physiol.* 99, 433-452
- [93] Messner, F., Grahmmer, J., Hautz, T., Brandacher, G., and Schneeberger, S. (2016). Ischemia/reperfusion injury in vascularized tissue allotransplantation: tissue damage and clinical relevance. *Curr. Opin. Organ Transplant.* 21, 503-509. doi: 10.1097/MOT.0000000000000343
- [94] Messoussi, A., Feneyrolles, C., Bros, A., Deroide, A., Daydé-Cazals, B., Chev, é, G., et al. (2014). Recent progress in the design, study, and development of c-Jun N-terminal kinase inhibitors as anticancer agents. *Chem. Biol.* 21, 1433-1443. doi: 10.1016/j.chembiol.2014.09.007
- [95] Milano, G., Morel, S., Bonny, C., Samaja, M., von Segesser, L. K., Nicod, P., et al. (2007). A peptide inhibitor of c-Jun NH2-terminal kinase reduces myocardial ischemia-reperfusion injury and infarct size in vivo. *Am. J. Physiol. Heart Circ. Physiol.* 292, H1828-H1835. doi: 10.1152/ajpheart.01117.2006
- [96] Monassier, J. P. (2008). Reperfusion injury in acute myocardial infarction. From bench to cath lab. Part I: Basic considerations. *Arch. Cardiovasc. Dis.* 101, 491-500. doi: 10.1016/j.acvd.2008.06.014
- [97] Morrison, A., Yan, X., Tong, C., and Li, J. (2011). Acute rosiglitazone treatment is cardioprotective against ischemia-reperfusion injury by modulating AMPK, Akt, and JNK signaling in nondiabetic mice. *Am. J. Physiol. Heart Circ. Physiol.* 301, H895-H902. doi: 10.1152/ajpheart.00137.2011
- [98] Murata, Y., Fujiwara, N., Seo, J. H., Yan, F., Liu, X., Terasaki, Y., et al. (2012). Delayed inhibition of c-Jun N-terminal kinase worsens outcomes after focal cerebral ischemia. *J. Neurosci.* 32, 8112-8115. doi: 10.1523/JNEUROSCI.0219-12.2012
- [99] Nakano, A., Baines, C. P., Kim, S. O., Pelech, S. L., Downey, J. M., Cohen, M. V., et al. (2000). Ischemic preconditioning activates MAPKAPK2 in the isolated rabbit heart: evidence for involvement of p38 MAPK. *Circ. Res.* 86, 144-151. doi: 10.1161/01.RES.86.2.144
- [100] Nijboer, C. H., van der Kooij, M. A., van Bel, F., Ohi, F., Heijnen, C. J., and Kavelaars, A. (2010). Inhibition of the JNK/AP-1 pathway reduces neuronal death and improves behavioral outcome after neonatal hypoxic-ischemic brain injury. *Brain Behav. Immun.* 24, 812-821. doi: 10.1016/j.bbi.2009.09.008
- [101] Oshikawa, J., Kim, S. J., Furuta, E., Caliceti, C., Chen, G. F., McKinney, R. D., et al. (2012). Novel role of p66Shc in ROS-dependent VEGF signaling and angiogenesis in endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.* 302, H724-H732. doi: 10.1152/ajpheart.00739.2011
- [102] Pan, J., Zhang, Q. G., and Zhang, G. Y. (2005). The neuroprotective effects of K252a through inhibiting MLK3/MKK7/JNK3 signaling pathway on ischemic brain injury in rat hippocampal CA1 region. *Neuroscience* 131, 147-159. doi: 10.1016/j.neuroscience.2004.09.031
- [103] Pei, D. S., Guan, Q. H., Sun, Y. F., Zhang, Q. X., Xu, T. L., and Zhang, G. Y. (2005). Neuroprotective effects of GluR6 antisense oligodeoxynucleotides on transient brain ischemia/reperfusion-induced neuronal death in rat hippocampal CA1 region. *J. Neurosci. Res.* 82, 642-649. doi: 10.1002/jnr.20669
- [104] Pei, D. S., Song, Y. J., Yu, H. M., Hu, W. W., Du, Y., and Zhang, G. Y. (2008). Exogenous nitric oxide negatively regulates c-Jun N-terminal kinase activation via inhibiting endogenous NO-induced S-nitrosylation during cerebral ischemia and reperfusion in rat hippocampus. *J. Neurochem.* 106, 1952-1963. doi: 10.1111/j.1471-4159.2008.05531.x
- [105] Pei, D. S., Wang, X. T., Liu, Y., Sun, Y. F., Guan, Q. H., Wang, W., et al. (2006). Neuroprotection against ischemic brain injury by a GluR6-9c peptide containing the TAT protein transduction sequence. *Brain.* 129(Pt 2), 465-479. doi: 10.1093/brain/awh700
- [106] Ping, P., Zhang, J., Huang, S., Cao, X., Tang, X. L., Li, R. C., et al. (1999). PKC-dependent activation of p46/p54 JNKs during ischemic preconditioning in conscious rabbits. *Am. J. Physiol.* 277(Pt 2), H1771-H1785. doi: 10.1152/ajpheart.1999.277.5.H1771
- [107] Qi, D., Hu, X., Wu, X., Merk, M., Leng, L., Bucala, R., et al. (2009). Cardiac macrophage migration inhibitory factor inhibits JNK pathway activation and injury during ischemia/reperfusion. *J. Clin. Invest.* 119, 3807-3816. doi: 10.1172/JCI39738
- [108] Rami, A., Langhagen, A., and Steiger, S. (2008). Focal cerebral ischemia induces upregulation of Beclin 1 and autophagy-like cell death. *Neurobiol. Dis.* 29, 132-141. doi: 10.1016/j.nbd.2007.08.005

- [109] Rose, B. A., Force, T., and Wang, Y. (2010). Mitogen-activated protein kinase signaling in the heart: angels versus demons in a heart-breaking tale. *Physiol. Rev.* 90, 1507-1546 doi: 10.1152/physrev.00054.2009
- [110] Rosso, C., Pires, C., Corvol, J. C., Baronnet, F., Crozier, S., Leger, A., et al. (2015). Hyperglycaemia, insulin therapy and critical penumbral regions for prognosis in acute stroke: further insights from the INSULINFARCT trial. *PLoS ONE* 10:e0120230. doi: 10.1371/journal.pone.0120230
- [111] Schepetkin, I. A., Kirpotina, L. N., Khlebnikov, A. I., Hanks, T. S., Kochetkova, I., et al. (2012). Identification and characterization of a novel class of c-Jun N-terminal kinase inhibitors. *Mol. Pharmacol.* 81, 832-845. doi: 10.1124/mol.111.077446
- [112] Shang, L., Ananthakrishnan, R., Li, Q., Quadri, N., Abdillahi, M., Zhu, Z., et al. (2010). RAGE modulates hypoxia/reoxygenation injury in adult murine cardiomyocytes via JNK and GSK-3 β signaling pathways. *PLoS ONE*. 5:e10092. doi: 10.1371/journal.pone.0010092
- [113] Shao, S., Xu, M., Zhou, J., Ge, X., Chen, G., Guo, L., et al. (2017). Atorvastatin attenuates ischemia/reperfusion-induced hippocampal neurons injury via Akt-nNOS-JNK signaling pathway. *Cell Mol. Neurobiol.* 37, 753-762. doi: 10.1007/s10571-016-0412-x
- [114] Shao, Z., Bhattacharya, K., Hsich, E., Park, L., Walters, B., Germann, U., et al. (2006). c-Jun N-terminal kinases mediate reactivation of Akt and cardiomyocyte survival after hypoxic injury in vitro and in vivo. *Circ. Res.* 98, 111-118. doi: 10.1161/01.RES.0000197781.20524.b9
- [115] Sharma, V., Bell, R. M., and Yellon, D. M. (2012). Targeting reperfusion injury in acute myocardial infarction: a review of reperfusion injury pharmacotherapy. *Expert Opin. Pharmacother.* 13, 1153-1175. doi: 10.1517/14656566.2012.685163
- [116] Shi, S., Li, Q. S., Li, H., Zhang, L., Xu, M., Cheng, J. L., et al. (2009). Anti-apoptotic action of hydrogen sulfide is associated with early JNK inhibition. *Cell Biol. Int.* 33, 1095-1101. doi: 10.1016/j.cellbi.2009.06.029
- [117] Shu, Y., Yang, Y., and Zhang, P. (2016). Neuroprotective effects of penehyclidine hydrochloride against cerebral ischemia-reperfusion injury in mice. *Brain Res. Bull.* 121, 115-123. doi: 10.1016/j.brainresbull.2016.01.008
- [118] Shvedova, M. V., Anfinogenova, Y., Popov, S. V., Shchepetkin, I. A., and Atochin, D. N. (2016). C-Jun N-terminal kinases and their modulators in myocardial ischemia/reperfusion injury. *Siberian Med. J.* 31, 7-15
- [119] Shvedova, M. V., Anfinogenova, Y., Schepetkin, I. A., and Atochin, D. N. (2017). The role of JNK-kinases and their inhibitors in neuroprotection in ischemic brain injury. *Russ. J. Physiol.* 103, 268-283
- [120] Sinha, S., Sinharoy, P., Bratz, I. N., and Damron, D. S. (2015). Propofol causes vasodilation in vivo via TRPA1 ion channels: role of nitric oxide and BKCa channels. *PLoS ONE* 10:e0122189. doi: 10.1371/journal.pone.0122189
- [121] Soga, M., Matsuzawa, A., and Ichijo, H. (2012). Oxidative stress induced diseases via the ASK1 signaling pathway. *Int. J. Cell Biol.* 2012:439587. doi: 10.1155/2012/439587
- [122] Song, Z. F., Ji, X. P., Li, X. X., Wang, S. J., Wang, S. H., et al. (2008). Inhibition of the activity of poly (ADP-ribose) polymerase reduces heart ischaemia/reperfusion injury via suppressing JNK-mediated AIF translocation. *J. Cell Mol. Med.* 12, 1220-1228. doi: 10.1111/j.1582-4934.2008.00183.x
- [123] Sun, H. Y., Wang, N. P., Halkos, M., Kerendi, F., Kin, H., Guyton, R. A., et al. (2006). Postconditioning attenuates cardiomyocyte apoptosis via inhibition of JNK and p38 mitogen-activated protein kinase signaling pathways. *Apoptosis* 11, 1583-1593. doi: 10.1007/s10495-006-9037-8
- [124] Sun, L., Isaak, C. K., Zhou, Y., Petkau, J. C., O, K., Liu, Y., et al. (2012). Salidroside and tyrosol from *Rhodiola* protect H9c2 cells from ischemia/reperfusion-induced apoptosis. *Life Sci.* 91, 151-158. doi: 10.1016/j.lfs.2012.06.026
- [125] Talmor, D., Applebaum, A., Rudich, A., Shapira, Y., and Tirosh, A. (2000). Activation of mitogen-activated protein kinases in human heart during cardiopulmonary bypass. *Circ. Res.* 86, 1004-1007. doi: 10.1161/01.RES.86.9.1004
- [126] Tian, H., Zhang, Q. G., Zhu, G. X., Pei, D. S., Guan, Q. H., and Zhang, G. Y. (2005). Activation of c-Jun NH2-terminal kinase 3 is mediated by the GluR6.PSD-95.MLK3 signaling module following cerebral ischemia in rat hippocampus. *Brain Res.* 1061, 57-66. doi: 10.1016/j.brainres.2005.09.001
- [127] Vassalli, G., Milano, G., and Moccetti, T. (2012). Role of mitogen-activated protein kinases in myocardial ischemia-reperfusion injury during heart transplantation. *J. Transplant.* 2012:928954. doi: 10.1155/2012/928954
- [128] Vlahopoulos, S., and Zoumpourlis, V. C. (2004). JNK: a key modulator of intracellular signaling. *Biochemistry* 69, 844-854. doi: 10.1023/B:BIRY.0000040215.02460.45
- [129] Waetzig, V., and Herdegen, T. (2005). Context-specific inhibition of JNKs: overcoming the dilemma of protection and damage. *Trends Pharmacol. Sci.* 26, 455-461. doi: 10.1016/j.tips.2005.07.006
- [130] Walshe, C. M., Laffey, J. G., Kevin, L., and O'Toole, D. (2015). Sepsis protects the myocardium and other organs from subsequent ischaemic/reperfusion injury via a MAPK-dependent mechanism. *Intensive Care Med.* Exp. 3:1. doi: 10.1186/s40635-014-0035-9

- [131] Wang, J., Ma, Y., Sachs, F., Li, J., and Suchyna, T. M. (2016). GsMTx4-D is a cardioprotectant against myocardial infarction during ischemia and reperfusion. *J. Mol. Cell Cardiol.* 98, 83-94. doi: 10.1016/j.yjmcc.2016.07.005
- [132] Wang, J., Yang, L., Rezaie, A. R., and Li, J. (2011). Activated protein C protects against myocardial ischemic/reperfusion injury through AMP-activated protein kinase signaling. *J. Thromb. Haemost.* 9, 1308-1317. doi: 10.1111/j.1538-7836.2011.04331.x
- [133] Wang, S., Yin, J., Ge, M., Dai, Z., Li, Y., Si, J., et al. (2016). Transforming growth- β 1 contributes to isoflurane postconditioning against cerebral ischemia-reperfusion injury by regulating the c-Jun N-terminal kinase signaling pathway. *Biomed. Pharmacother.* 78, 280-290. doi: 10.1016/j.biopha.2016.01.030
- [134] Wang, X. T., Pei, D. S., Xu, J., Guan, Q. H., Sun, Y. F., Liu, X. M., et al. (2007). Opposing effects of Bad phosphorylation at two distinct sites by Akt1 and JNK1/2 on ischemic brain injury. *Cell Signal.* 19, 1844-1856. doi: 10.1016/j.cellsig.2007.04.005
- [135] Wang, Z., Huang, H., He, W., Kong, B., Hu, H., Fan, Y., et al. (2016). Regulator of G-protein signaling 5 protects cardiomyocytes against apoptosis during in vitro cardiac ischemia-reperfusion in mice by inhibiting both JNK1/2 and P38 signaling pathways. *Biochem. Biophys. Res. Commun.* 473, 551-557. doi: 10.1016/j.bbrc.2016.03.114
- [136] Wei, C., Zhao, Y., Wang, L., Peng, X., Li, H., Zhao, Y., et al. (2015). H₂S restores the cardioprotection from ischemic post-conditioning in isolated aged rat hearts. *Cell Biol. Int.* 39, 1173-1176. doi: 10.1002/cbin.10507
- [137] Wei, J., Wang, W., Chopra, I., Li, H. F., Dougherty, C. J., Adi, J., et al. (2011). c-Jun N-terminal kinase (JNK-1) confers protection against brief but not extended ischemia during acute myocardial infarction. *J. Biol. Chem.* 286, 13995-14006. doi: 10.1074/jbc.M110.211334
- [138] Wen, J., Watanabe, K., Ma, M., Yamaguchi, K., Tachikawa, H., Kodama, M., et al. (2006). Edaravone inhibits JNK-c-Jun pathway and restores anti-oxidative defense after ischemia-reperfusion injury in aged rats. *Biol. Pharm. Bull.* 29, 713-718. doi: 10.1248/bpb.29.713
- [139] Wen, X. R., Li, C., Zong, Y. Y., Yu, C. Z., Xu, J., Han, D., et al. (2008). Dual inhibitory roles of geldanamycin on the c-Jun NH₂-terminal kinase 3 signal pathway through suppressing the expression of mixed-lineage kinase 3 and attenuating the activation of apoptosis signal-regulating kinase 1 via facilitating the activation of Akt in ischemic brain injury. *Neuroscience* 156, 483-497. doi: 10.1016/j.neuroscience.2008.08.006
- [140] Wen, X. R., Tang, M., Qi, D. S., Huang, X. J., Liu, H. Z., Zhang, F., et al. (2016). Butylphthalide suppresses neuronal cells apoptosis and inhibits JNK-caspase-3 signaling pathway after brain ischemia/reperfusion in rats. *Cell Mol. Neurobiol.* 36, 1087-1095. doi: 10.1007/s10571-015-0302-7
- [141] Wiegler, K., Bonny, C., Coquoz, D., and Hirt, L. (2008). The JNK inhibitor XG-102 protects from ischemic damage with delayed intravenous administration also in the presence of recombinant tissue plasminogen activator. *Cerebrovasc. Dis.* 26, 360-366. doi: 10.1159/000151639
- [142] Wiltshire, C., Gillespie, D. A., and May, G. H. (2004). Sab (SH3BP5), a novel mitochondria-localized JNK-interacting protein. *Biochem. Soc. Trans.* 32(Pt 6), 1075-1077. doi: 10.1042/BST0321075
- [143] Wu, J., Li, J., Zhang, N., and Zhang, C. (2011). Stem cell-based therapies in ischemic heart diseases: a focus on aspects of microcirculation and inflammation. *Basic Res. Cardiol.* 106, 317-324. doi: 10.1007/s00395-01-0168-x
- [144] Xie, P., Guo, S., Fan, Y., Zhang, H., Gu, D., et al. (2009). Atrogin-1/MAFbx enhances simulated ischemia/reperfusion-induced apoptosis in cardiomyocytes through degradation of MAPK phosphatase-1 and sustained JNK activation. *J. Biol. Chem.* 284, 5488-5496. doi: 10.1074/jbc.M806487200
- [145] Xu, H., Yao, Y., Su, Z., Yang, Y., Kao, R., Martin, C. M., et al. (2011). Endogenous HMGB1 contributes to ischemia-reperfusion-induced myocardial apoptosis by potentiating the effect of TNF- α /JNK. *Am. J. Physiol. Heart Circ. Physiol.* 300, H913-H921. doi: 10.1152/ajpheart.00703.2010
- [146] Xu, J., Qin, X., Cai, X., Yang, L., Xing, Y., Li, J., et al. (2015). Mitochondrial JNK activation triggers autophagy and apoptosis and aggravates myocardial injury following ischemia/reperfusion. *Biochim. Biophys. Acta* 1852, 262-270. doi: 10.1016/j.bbdis.2014.05.012
- [147] Yang, S. I., Yuan, Y., Jiao, S., Luo, Q. I., and Yu, J. (2016). Calcitonin gene-related peptide protects rats from cerebral ischemia-reperfusion injury via a mechanism of action in the MAPK pathway. *Biomed. Rep.* 4, 699-703. doi: 10.3892/br.2016.658
- [148] Yao, B., Wang, S., Xiao, P., Wang, Q., Hea, Y., and Zhang, Y. (2017). MAPK signaling pathways in eye wounds: multifunction and cooperation. *Exp. Cell Res.* 359, 10-16. doi: 10.1016/j.yexcr.2017.06.024
- [149] Yu, H. M., Xu, J., Li, C., Zhou, C., Zhang, F., Han, D., et al. (2008). Coupling between neuronal nitric oxide synthase and glutamate receptor 6-mediated c-Jun N-terminal kinase signaling pathway via S-nitrosylation contributes to ischemia neuronal death. *Neuroscience* 155, 1120-1132. doi: 10.1016/j.neuroscience.2008.03.061
- [150] Yue, T. L., Wang, C., Gu, J. L., Ma, X. L., Kumar, S., Lee, J. C., et al. (2000). Inhibition of extracellular signal-regulated kinase enhances Ischemia/Reoxygenation-induced apoptosis in cultured cardiac myocytes and exaggerates reperfusion injury in isolated perfused heart. *Circ. Res.* 86, 692-699. doi: 10.1161/01.RES.86.6.692

- [151] Zaha, V. G., Qi, D., Su, K. N., Palmeri, M., Lee, H. Y., Hu, X., et al. (2016). AMPK is critical for mitochondrial function during reperfusion after myocardial ischemia. *J. Mol. Cell Cardiol.* 91, 104-113. doi: 10.1016/j.yjmcc.2015.12.032
- [152] Zeke, A., Misheva, M., Reményi, A., and Bogoyevitch, M. A. (2016). JNK Signaling: regulation and functions based on complex protein-protein partnerships. *Microbiol. Mol. Biol. Rev.* 80, 793-835. doi: 10.1128/MMBR.00043-14
- [153] Zhang, G. M., Wang, Y., Li, T. D., Li, X. Y., Su, S. P., Sun, Y. Y., et al. (2014). Post-conditioning with gradually increased reperfusion provides better cardioprotection in rats. *World J. Emerg. Med.* 5, 128-134. doi: 10.5847/wjem.j.issn.1920-8642.2014.02.009
- [154] Zhang, G. M., Wang, Y., Li, T. D., Zhang, D. W., Liu, X. H., and Yang, F. F. (2009). Change of JNK MAPK and its influence on cardiocyte apoptosis in ischemic postconditioning. *J. Zhejiang Univ. Med. Sci.* 38, 611-619
- [155] Zhang, H., Wu, W., Du, Y., Santos, S. J., Conrad, S. E., Watson, J. T., et al. (2004). Hsp90/p50cdc37 is required for mixed-lineage kinase (MLK) 3 signaling. *J. Biol. Chem.* 279, 19457-19463. doi: 10.1074/jbc.M311377200
- [156] Zhang, J., Li, X. X., Bian, H. J., Liu, X. B., Ji, X. P., and Zhang, Y. (2009). Inhibition of the activity of Rho-kinase reduces cardiomyocyte apoptosis in heart ischemia/reperfusion via suppressing JNK-mediated AIF translocation. *Clin. Chim. Acta* 401, 76-80. doi: 10.1016/j.cca.2008.11.016
- [157] Zhen, Y., Ding, C., Sun, J., Wang, Y., Li, S., and Dong, L. (2016). Activation of the calcium-sensing receptor promotes apoptosis by modulating the JNK/p38 MAPK pathway in focal cerebral ischemia-reperfusion in mice. *Am. J. Transl. Res.* 8, 911-921
- [158] Zheng, G. Y., Chen, X. C., Du, J., Liu, C. Y., Fang, F., Zhang, J., et al. (2006). Inhibitory action of propyl gallate on the activation of SAPK/JNK and p38MAPK induced by cerebral ischemia-reperfusion in rats. *Acta Pharm. Sin.* 41, 548-554
- [159] Zhu, H., Zhang, Y., Shi, Z., Lu, D., Li, T., Ding, Y., et al. (2016). The Neuroprotection of liraglutide against ischaemia-induced apoptosis through the activation of the PI3K/AKT and MAPK pathways. *Sci. Rep.* 6:26859. doi: 10.1038/srep26859
- [160] Zhu, J., Yi, X., Zhang, J., Chen, S., and Wu, Y. (2017). Chemical profiling and antioxidant evaluation of Yangxinshi Tablet by HPLC-ESI-Q-TOF-MS/MS combined with DPPH assay. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 1060, 262-271. doi: 10.1016/j.jchromb.2017.06.022
- [161] Zinkel, S., Gross, A., and Yang, E. (2006). BCL2 family in DNA damage and cell cycle control. *Cell Death Differ.* 13, 1351-1359. doi: 10.1038/sj.cdd.4401987