


**KAZAN FEDERAL UNIVERSITY
A. E. ARBUZOV INSTITUTE OF ORGANIC AND PHYSICAL
CHEMISTRY
RUSSIAN SCIENTIFIC FOUNDATION
RUSSIAN FOUNDATION OF BASIC RESEARCH
RUSSIAN ACADEMY OF SCIENCES
FEDERAL AGENCY OF SCIENTIFIC ORGANIZATIONS
D.I. MENDELEEV RUSSIAN CHEMICAL SOCIETY**



**VIIIth INTERNATIONAL SYMPOSIUM
«DESIGN AND SYNTHESIS OF SUPRAMOLECULAR
ARCHITECTURES»**

**IInd YOUTH SCHOOL ON SUPRAMOLECULAR AND
COORDINATION CHEMISTRY**

**April 25-29, 2016
Kazan, Russia**

INTERNATIONAL ADVISORY COMMITTEE

Prof. Alexander Konovalov - Chairman

Prof. Vladimir Arslanov (Russia)

Prof. Antony Coleman (France)

Prof. Vladimir Fedin (Russia)

Prof. Olga Fedorova (Russia)

Prof. Yulia Gorbunova (Russia)

Prof. Sergei Gromov (Russia)

Prof. Sylvie Ferlay (France)

Prof. Mir Wais Hosseini (France)

Prof. Vitaly Kalchenko (Ukraine)

Prof. Burkhard Koenig (Germany)

Prof. Janusz Lipkowski (Poland)

Prof. David Reinhoudt (Netherlands)

Prof. Kinga Suwinska (Poland)

ORGANIZING COMMITTEE

Prof. Igor Antipin - Chairman

Prof. Svetlana Solovieva - Secretary

Dr. Vladimir Burilov

Dr. Yulia Elistratova

Prof. Valery Gorbachuk

Nataly Ivanova

Dr. Alla Mirgorodskaya

Dr. Diana Mironova

Dr. Anton Muravev

Prof. Asiya Mustafina

Dr. Alexander Ovsyannikov

Dr. Farida Valeeva

Prof. Lucia Zakharova

Dr. Albina Ziganshina

Dr. Marat Ziganshin

ELUCIDATION OF HYBRID MULTI-PORPHYRIN SUPRAMOLECULAR ASSEMBLIES BY 2D NMR SPECTROSCOPY

I. A. Khodov, N. Zh. Mamardashvili

Institute of Solution Chemistry of Russian Academy of Science, Ivanovo, Russia

iakh@isc-ras.ru

Two-dimensional nuclear magnetic resonance (NMR) spectroscopy studies of supramolecular assemblies were carried out. The chemical structure of supramolecular assemblies were confirmed by the two-dimensional NMR. It was found that two-dimensional rotating-frame Overhauser spectroscopy and double-quantum filtered correlation spectroscopy for this system are not sufficiently informative. Moreover, the explanation of experimental data is complicated by the fact that spectral signals of the supramolecular assemblies molecules at the same time depend on two factors: a variation of the electron density distribution and a change in the shielding upon complex formation. Thus questions about the chemical structure of the complexes formed and their size remains open. Therefore, to obtain true evidence of the fact of information about their size and complex formation, other way should be applied. We have used an approach to elucidate the chemical structures for complex that is based on the analysis of two-dimensional high-resolution diffusion-ordered NMR spectroscopy.

1. I.A. Khodov, G.A. Alper, G.M. Mamardashvili, N.Z. Mamardashvili . *J. Mols. Str.*, 2015, **1090**, 174-180.

Acknowledgements This work was supported by Russian Scientific Foundation, project № 14-13-00232.